
13 unflagged sessions: accepted signal-like 266 ROIs; rejected 3234 ROIs, of which 21% [12, 33] (mean over sessions) pass the axon transient test (activity-like) and 1230 form the strict control (fail the test, skew < 0.3, not touching accepted ROIs)



i

session	accepted sig.-like	rejected	activity-like	strict control
RB22 10-23 1427	6	124	44%	41
RB22 10-27 1138	12	340	13%	164
RB23 10-23 1642	29	386	9%	156
RB23 10-27 1423	11	85	14%	27
RB24 10-21 1630	20	318	31%	104
RB24 10-23 1538	13	231	22%	115
RB24 10-27 1456	10	139	15%	87
RB24 10-27 1534	8	90	16%	47
RB24 10-29 1752	8	153	18%	67
RB24 10-30 1744	8	143	29%	56
RB24 11-04 1651	50	397	19%	107
RB24 11-05 1423	47	393	21%	108
RB24 11-05 1520	44	435	20%	151

All ROIs of the session's suite2p plane0 output with the same F / Fneu and dF/F recipe; rejected = iscell 0, not a hidden constituent of a merged ROI (suite2p inmerge > 0), median Fc > 5 counts. Activity-like = passes the transient test used for the accepted axons (slow variance >= 4 x white-noise expectation, upward events >= 2 x downward events) applied on the 5 Hz grid (approximate: accepted ROIs pass 40-90%). Coupling = r (lag 0, band-passed 1-30 s) with whisking and its circular-shift p < 0.01 (shift >= 120 s). Panel g: the same ROIs (first 220 rejected ROIs per session) with dF/F recomputed from F alone or from the neuropil trace alone. 13 unflagged sessions, 3 mice.