Results ledger, continued: Figs 9-12 (whisking shuffle tests, loom heatmaps, lick task vs passive, rejected ROIs)

what was tested	effect [95% CI] unit	sessions / mice	verdict
Fig. 9 Whisking vs axon calcium (shuffle tests)			
peak r axon mean vs whisking, shuffle-tested per session	+0.48 [0.26, 0.64] r	13 / 3	SUPPORTED
fraction of signal-like ROIs with r > 0 and p < 0.01 against time-shifted whisking	+0.77 [0.52, 0.95] of ROIs	13 / 3	AMBIGUOUS
partial r of axon mean with whisk (the other two behaviours regressed out, lag 0)	+0.13 [0.01, 0.28] r	13 / 3	AMBIGUOUS
partial r of axon mean with face (the other two behaviours regressed out, lag 0)	+0.12 [-0.01, 0.26] r	13 / 3	AMBIGUOUS
partial r of axon mean with pupil (the other two behaviours regressed out, lag 0)	+0.24 [0.09, 0.36] r	13 / 3	AMBIGUOUS
Fig. 10 Looming discs: heatmaps and per-session responses			
unflagged sessions with a clear expanding-loom response (z > 2.33 raw AND after movement removal)	+0.20 of sessions	5 / 3	AMBIGUOUS
unflagged sessions with a clear receding-loom response (z > 2.33 raw AND after movement removal)	+0.00 of sessions	4 / 3	NOT SEEN
Fig. 11 Lick task vs passive square, engagement (behaviours regressed)			
axon response to the square, cue − pre, lick-task sessions (raw)	+0.007 [-0.011, 0.026] dF/F	4 / 1	NOT SEEN
axon response to the square, cue − pre, lick-task sessions (whisking, face, pupil, ball regressed)	-0.004 [-0.020, 0.014] dF/F	4 / 1	NOT SEEN
axon response to the square, cue − pre, passive sessions (raw)	+0.004 [-0.007, 0.010] dF/F	3 / 2	NOT SEEN
axon response to the square, cue − pre, passive sessions (whisking, face, pupil, ball regressed)	+0.005 [-0.007, 0.013] dF/F	3 / 2	NOT SEEN
square response cue − pre: lick-task minus passive sessions (raw)	+0.005 dF/F	7 / 2	NOT SEEN
square response cue − pre: lick-task minus passive sessions (behaviours regressed)	-0.007 dF/F	7 / 2	NOT SEEN
axon response to the square, post − pre, lick-task sessions (raw)	+0.052 [0.009, 0.106] dF/F	4 / 1	AMBIGUOUS
axon response to the square, post − pre, lick-task sessions (whisking, face, pupil, ball regressed)	+0.015 [-0.012, 0.048] dF/F	4 / 1	NOT SEEN
axon response to the square, post − pre, passive sessions (raw)	+0.011 [0.010, 0.012] dF/F	3 / 2	NOT SEEN
axon response to the square, post − pre, passive sessions (whisking, face, pupil, ball regressed)	+0.016 [0.003, 0.032] dF/F	3 / 2	NOT SEEN
square response post − pre: lick-task minus passive sessions (raw)	+0.041 dF/F	7 / 2	NOT SEEN
square response post − pre: lick-task minus passive sessions (behaviours regressed)	-0.002 dF/F	7 / 2	NOT SEEN
axon activity in hit-trial vs miss-trial cycles (raw)	+0.024 [-0.001, 0.054] dF/F	4 / 1	NOT SEEN
axon activity in hit-trial vs miss-trial cycles (behaviours regressed)	-0.007 [-0.012, -0.000] dF/F	4 / 1	NOT SEEN
axon activity vs hit rate of the surrounding 10 trials (raw)	+0.001 [-0.003, 0.004] dF/F per SD	4 / 1	NOT SEEN
axon activity vs hit rate of the surrounding 10 trials (behaviours regressed)	-0.000 [-0.006, 0.005] dF/F per SD	4 / 1	NOT SEEN
share of axon variance explained by whisking, face, pupil, ball (in-sample, lags -1..+3 s), lick-task sessions	+0.49 [0.37, 0.61] R2	4 / 1	AMBIGUOUS
share of axon variance explained by whisking, face, pupil, ball (in-sample, lags -1..+3 s), passive sessions	+0.49 [0.45, 0.52] R2	3 / 2	SUPPORTED
Fig. 12 What the rejected ROIs are (control population)			
fraction of rejected ROIs that pass the axon transient criterion (5 Hz grid version)	+0.21 [0.12, 0.33] fraction of rejected ROIs	13 / 3	AMBIGUOUS
strict control ROIs significantly positively coupled to whisking	35 [27, 44] % of ROIs	13 / 3	AMBIGUOUS
strict control ROIs significantly negatively coupled to whisking	44 [29, 59] % of ROIs	13 / 3	AMBIGUOUS
rejected ROIs significantly NEGATIVELY coupled to whisking with dF/F from F alone (no neuropil subtraction)	n/a	13 / 3	AMBIGUOUS

Verdict rule: SUPPORTED = 95% CI (hierarchical bootstrap over animals / sessions) excludes 0 and most sessions agree; any result carried by one animal is downgraded to AMBIGUOUS automatically; NOT SEEN = CI includes 0 and no session-level support. Effects are in different units and are not comparable with each other. "Unit" columns: r = correlation, z = session z, dF/F = high-passed, SD = SD of the metric over the session's windows. n = 3 mice at most: every number describes these animals.