

① a  $M = -19$

$$d = 10^9 \text{ lightyears} = 3 \times 10^8 \text{ pc}$$

(where  $1 \text{ ly} = 3.3 \text{ pc}$ )

$$m - M = 5 \log \left( \frac{3 \times 10^8}{10} \right) = 5 \log (3 \times 10^7) = 5(\log 3 + \log 7)$$

$$= 5(0.5 + 7) = 37.5$$

$$m - M = 37.5$$

$$m - (-19) = 37.5$$

$$\boxed{m = 18.5 \text{ magnitudes}}$$

b  $\lambda_0 = 0.65 \text{ microns}$

$$z = \frac{1}{a} - 1 = \frac{1}{0.9} - 1 = 1.11 - 1 = 0.11$$

$$z = \frac{\lambda - \lambda_0}{\lambda_0}$$

$$0.11 \lambda_0 = \lambda - \lambda_0$$

$$0.11 \lambda_0 + \lambda_0 = \lambda$$

$$\lambda_0 (1.11) = \lambda$$

$$\lambda = 0.65 (1.11) = \boxed{0.72 \text{ microns}}$$

OR

$$\frac{a_{\text{now}}}{a_{\text{then}}} = \frac{\lambda_{\text{obs}}}{\lambda_{\text{emit}}}$$

$$\frac{1}{0.9} = \frac{\lambda_{\text{obs}}}{0.65 \text{ microns}}$$

$$\boxed{\lambda_{\text{obs}} = 0.72 \text{ microns}}$$

II  $z=1.5$

$m-M=42.5$

$a = \frac{1}{1+z} = \frac{1}{1+1.5} = 0.4$

$m-M = 5 \log\left(\frac{d}{10}\right)$

$42.5 = 5 \log\left(\frac{d}{10}\right)$

$d = 3 \times 10^9 \text{ pc} = 10^{10} \text{ lightyears}$


point "now"

$a=1 \quad t=12.5 \text{ billion yrs}$

point "then"

$a=0.4 \quad t=2.5 \text{ billion yrs}$

↑ on  $10^{10}$  billion yrs IN THE PAST




universe is accelerating

We know this because at  $t=2.5$  billion years the expansion of the empty universe had a rate of  $z=4$  ( $a=0.2$  at 2.5 billion on the empty line), while the observed point has a slower expansion at the same time,  $z=1.5$ . Since all points must meet at "now" the observed universe expansion needs to speed up to end up matching the empty universe "now".

③ a We need dark energy (DE) to explain why the observations of SNe Ia appear fainter at high  $z$  than would be expected in a pure  $\Omega_m = 1$  universe. This assumes that Ia's are **STANDARD CANDLES!** If distance SNe Ia's are fainter than nearby ones, then the fact that high  $z$  Ia's appear faint may be due merely to the fact that they ARE faint. In that case, DE need not be invoked to explain the observed points.

b from  $\Delta(m-M)$  plot at  $z=1$  the magnitude difference between the  $\Omega_m = 0.25 \Omega_\Lambda = 0.75$  line and the  $\Omega_m = 1$  line is 0.6 magnitudes



$$0.6 = 2.5 \log \left( \frac{b_{z=1}}{b_{z=0}} \right)$$

$$b_{z=1} = 0.58 b_{z=0}$$

$\Rightarrow$  nearby SNe are about twice as bright as the distance SNe