



Supine Hamstring Stretch - Strap

How: use a stretch strap, bath towel, or bed sheet. Wrap it around the arch of your foot, then lay down on your back in a relaxed and comfortable position. Keeping your leg completely relaxed, use your arms to pull the strap. Feel: you will feel a gentle stretch behind your thigh and/or knee. Compensation : try to keep your knee straight as you pull on the strap to ensure a good stretch.

Details:

No specific exercise parameters provided.



Tandem Balance

How: get set-up standing with your feet in a heel-to-toe position. Maintain your balance, switch feet and perform again. Feel: you should feel like your balance is being challenged. If you're having difficulty maintaining your balance, try to focus on an object with your eyes in the distance or near you. Compensation: try not to lose your posture or excessively lean forward/backward or side to side. If you lose your balance it is ok, take a step and get back to starting position. Do not allow yourself to fall or take the risk of injuring yourself.

Details:

No specific exercise parameters provided.



Sit To Stand

How: place a box, chair, or elevated surface behind you. Sink your hips down and backwards with good squat form, and completely sit down. Then, shift your weight forward and stand back up. Feel: you should feel your thigh and butt muscles working. Compensation: try your best not to fall into the chair. Try to control yourself as your lower into the chair. Do not use momentum to stand up, if able.

Details:

No specific exercise parameters provided.



BIG: Standing Hip Abduction

Stand upright while holding onto a stable surface for support (kitchen sink works well). Slowly move the working leg out to the side while keeping the trunk upright and the pelvis level. Return to the starting position with control and repeat. don't need ankle weights, but can for added difficulty. moving slow and steady, no leaning. the less dependent you are on your hold, making you balance more with your legs the better.

Reps: 20

Hold: 3-5 seconds.

Daily: 1-2 times a day.

Ankle weights: in clinic we typically use 3-5 lbs.

Details:

No specific exercise parameters provided.



BIG: Seated Knee Extension (Long Arc Quad LAQ)

Sit at the edge of a table or chair with feet hanging freely (if possible). Slowly straighten your knee until your leg is fully extended, then lower it back down with control. Keep your trunk upright and avoid leaning back. Ankle weights are not always necessary, but often used in clinic.

Hold for 3 seconds.

Reps: 20

Daily: 2-3 times per day.

Details:

No specific exercise parameters provided.



BIG: Standing Knee Bend / Standing Hamstring Curl (Supported)

Stand tall while holding onto a stable surface for balance. Slowly bend your knee, bringing your heel toward your buttocks while keeping your thighs aligned and your hips steady. Lower the leg back down with control and repeat. ankle weights are not always necessary, but often used in clinic.

Hold for 3 seconds.

Reps: 20

Daily: 2-3 times per day.

Details:

No specific exercise parameters provided.



BIG: Toe and Heel Raises

Stand tall while lightly holding onto a stable surface for balance. the kitchen sink is a good place to hold on to. do not fall backwards when raising your toes.

Heel raises: lift your heels off the ground, rising up onto the balls of your feet. Hold briefly, then lower slowly.

Toe raises: lift your toes and the front of your foot while keeping your heels on the ground. Hold briefly, then lower with control.

Repeat in a smooth, controlled motion without leaning forward or backward.

Hold: pause for a second while on your toes and when on your heels.

Reps: 20

Daily: 2-3 times per day.

Details:

No specific exercise parameters provided.