

Computational Identification of Unknown Organic Contaminants and Molecules in Remote Environments via Molecular Networking

Raelyn Brooks

April 30, 2026

1 Hypothesis

This project aims to identify contaminants of emerging concern (CECs) in surface water samples from remote, high-altitude environments, including Mt. Everest, the Tibetan Plateau, Qinghai Lake, and Mt. Kilimanjaro, using high-resolution mass spectrometry (HRMS). While existing database-driven, targeted approaches can successfully identify known contaminants, most current software and reference libraries are not well optimized for complex environmental matrices, leaving significant gaps in identification coverage. I hypothesize that a substantial fraction of these unidentified molecules can be assigned through non-targeted analysis, particularly molecular networking, thereby extending existing contaminant databases. Together, these approaches will improve contaminant identification in remote environmental systems where current reference frameworks are incomplete.




Figure 1: Comparison of non-targeted analysis (NTA) and targeted analysis (TA) approaches for identification of organic contaminants [22].

2 Methods

2.1 High-Resolution Mass Spectrometry Data Acquisition

All samples will be analyzed using liquid chromatography coupled with high-resolution mass spectrometry (LC-HRMS) using an Agilent 1260 Infinity II HPLC and Agilent 6546 QTOF-MS. Chromatographic separation will be performed using a C18 reverse-phase column, allowing for the retention and separation of semi-polar to nonpolar organic compounds.

LCMS-grade water and LCMS-grade methanol will be used for all mobile phases to ensure minimal background contamination. Data will be acquired in both positive and negative electrospray ionization (ESI) modes to maximize compound coverage. For positive ionization mode, the mobile phase will consist of 0.1% formic acid in LCMS-grade water (aqueous phase) and methanol with 0.1% formic acid (organic phase) [9]. For negative ionization mode, the mobile phase will consist of 10 mM ammonium acetate in LCMS-grade water (aqueous phase) and 10 mM ammonium acetate in methanol (organic phase) [9]. Gradient elution will be used in both modes to separate compounds across a range of polarities.

The analytical workflow will begin with MS1 full-scan acquisition to detect all ionizable features present in the samples. From these data, exclusion and inclusion lists will be generated (Figure 2).

The exclusion list will contain background contaminants high-abundance ions that do not require repeated fragmentation, while the inclusion list will prioritize features of interest, particularly those matching entries in existing chemical databases [16].

Subsequent analyses will employ MS/MS acquisition using an iterative data-dependent acquisition (DDA) strategy. In DDA, precursor ions are selected in real time based on their intensity for fragmentation. In this study, each sample will be analyzed across 3–4 iterative runs. After each run, previously fragmented ions are added to an exclusion list, allowing subsequent runs to target new, lower-abundance features. This iterative process increases MS/MS coverage by systematically expanding fragmentation across the detected feature space, improving the likelihood of identifying both known and unknown compounds in complex environmental mixtures.




Figure 2: Example of exclusion and inclusion of features to establish our lists. [16]

2.2 Data Processing and Feature Extraction

Raw LC-HRMS data will be processed using MS-DIAL v5.5.250530. Data processing will include peak detection, alignment across samples, and feature annotation. A representative workflow is shown in Figure 3; although this figure illustrates data-independent acquisition (DIA), the overall processing steps are similar to those used for data-dependent acquisition (DDA) in this study.




Figure 3: This workflow illustrates data independent acquisition (DIA) using MS Dial while different from our data dependent acquisition (DDA) we are seeking something similar. [24]

Detected signals will initially be treated as features, defined by their mass-to-charge ratio (m/z), retention time, and associated MS/MS spectra. A feature is only considered a confirmed compound once sufficient evidence supports its identification.

Quality control steps, including blank subtraction and filtering of low-confidence signals, will be applied to remove background contaminants [22]. The resulting feature table will represent the set of detected molecular features across all samples and will serve as the basis for downstream identification and analysis.

2.3 Suspect Screening for Known Contaminants

To identify contaminants of emerging concern (CECs), suspect screening will be performed using curated chemical databases, such as the Agilent environmental contaminant database. Figure 4 illustrates how MS/MS spectral databases are constructed and used in conjunction with LC-HRMS

data to identify compounds based on fragmentation patterns.

Features will be matched to database entries using a combination of accurate mass, MS/MS spectral similarity, isotopic pattern, and retention time [14]. Retention time serves as an important orthogonal parameter that strengthens confidence in candidate matches.




Figure 4: This figure illustrates the MS/MS database creation and how LCMS is used to identify compounds based on their fragmentation patterns [14].

Candidate annotations will be evaluated using the Schymanski framework, which classifies identification confidence into levels based on available evidence (Figure 5) [23]. These levels reflect the degree of confidence in a proposed identification rather than representing definitive assignments. This approach enables identification of known contaminants while simultaneously highlighting gaps in existing chemical databases where features cannot be confidently matched [18].




Figure 5: Confidence levels example for molecular feature identification based on the Schymanski framework. [23]

2.4 Molecular Networking and Non-Targeted Analysis

To extend beyond database-driven identification, molecular networking will be used to characterize unknown features not identified through suspect screening. Molecular networking is a computational approach that groups compounds based on similarities in their MS/MS fragmentation patterns, enabling structurally related molecules to be identified even when exact matches are not present in databases [25].

Existing tools such as Global Natural Products Social Molecular Networking (GNPS) are widely used for this purpose; however, GNPS is primarily designed for natural product datasets and is not optimized for environmental contaminant analysis [21, 25]. As shown in Figure 6, this can lead to gaps in annotation when applied to environmental samples. Therefore, a custom computational workflow will be developed to better suit the characteristics of environmental HRMS data.

Spectral similarity between features will be quantified using cosine similarity scoring. In this approach, each MS/MS spectrum is represented as a vector, where fragment ions (m/z values) are paired with their corresponding intensities [2]. Before comparison, spectra are typically filtered and aligned so that only matching or near-matching fragment ions are considered [6]. The cosine similarity score between two spectra is calculated as:

$$\text{Cosine Similarity} = \frac{\sum(A_i \cdot B_i)}{\sqrt{\sum(A_i^2)} \cdot \sqrt{\sum(B_i^2)}}$$

where A_i and B_i represent the intensities of matching fragment ions in the two spectra. This calculation measures how similar two fragmentation patterns are, with scores ranging from 0 (no similarity) to 1 (identical spectra).

This method considers:

- The presence of shared fragment ions (m/z alignment)
- The relative intensities of those fragments

A cosine similarity threshold of greater than or equal to 0.70 will be used to define meaningful structural similarity between an unknown feature and a known compound (known pesticides or contaminants). Features exceeding this threshold will be grouped and used to construct a molecular similarity graph, enabling identification of structurally related unknowns.




Figure 6: This example of Global Natural Product Social (GNPS) molecular networking illustrates the relationships between different molecular features but this is not fitted for environmental samples which causes gaps [21].

2.5 Structural Interpretation and Data Analysis

Structural interpretation will focus on analyzing mass differences and fragmentation patterns within molecular networks to propose tentative structures for unknown compounds [28]. These mass shifts are of particular interest because they often correspond to known environmental transformation processes, such as oxidation, reduction, or functional group modification. For example:

- A mass increase of +16 Da may indicate the addition of an oxygen atom, commonly associated with hydroxylation reactions.
- A mass shift of +14 Da may suggest methylation.
- A mass change of +16 Da combined with loss of 2 hydrogen atoms may indicate the formation of a carbonyl group through oxidation.

These transformations are well-documented in environmental chemistry and provide insight into degradation pathways and secondary pollutant formation [28].

Fragmentation patterns will be analyzed alongside these mass differences to strengthen structural hypotheses [21]. A molecular networking framework (Figure 7) will be used to visualize relationships between features and support the identification of unknown compounds based on their similarity to known structures.

This integrated workflow enables (1) comprehensive MS/MS data acquisition, (2) identification of known contaminants using existing libraries, and (3) extension beyond current databases through molecular networking, directly supporting the three specific aims of this study.




Figure 7: This diagram illustrates how a molecular network can be constructed from MS/MS data, enabling the identification of unknown compounds based on their fragmentation patterns while this is ion identity molecular networking a similar graph will be constructed [21].

References

- [1] Francesco Asnicar, Andrew Maltez Thomas, Andrea Passerini, Levi Waldron, and Nicola Segata. Machine learning for microbiologists. *Nature Reviews Microbiology*, 22(4):191–205, Apr 2024.
- [2] Wout Bittremieux, Robin Schmid, Florian Huber, Justin J. J. van der Hooft, Mingxun Wang, and Pieter C. Dorrestein. Comparison of cosine, modified cosine, and neutral loss based spectrum alignment for discovery of structurally related molecules. *Journal of the American Society for Mass Spectrometry*, 33(9):1733–1744, 2022.
- [3] Jules M. Blais, Michael R. Rosen, and John P. Smol, editors. *Environmental Contaminants: Using Natural Archives to Track Sources and Long-Term Trends of Pollution*. Springer, Dordrecht, 2015.
- [4] Jordi. Catalan, editor. *Tracking Long-Range Atmospheric Transport of Trace Metals, Polycyclic Aromatic Hydrocarbons, and Organohalogen Compounds Using Lake Sediments of Mountain Regions*. Springer, Dordrecht, 2015.
- [5] Alex Chao, Jeffrey M. Minucci, Troy M. Ferland, E. Tyler Carr, Greg Janesch, Safia Rizwan, Heather D. Whitehead, Tommy Cathey, Shirley Pu, Laura D. Brunelle, Angela L. Batt, Jon R. Sobus, and Antony J. Williams. Prioritizing chemical candidates from non-targeted analysis using metadata, spectral similarity, and hazard scoring within INTERPRET NTA. *Analytical Chemistry*, 97(29):15904–15912, 2025.
- [6] Nicola Dalla Valle, Mar Garcia-Aloy, Peter Robatscher, Pietro Franceschi, and Michael Oberhuber. Improving spectral similarity and molecular network reliability through noise signal filtering in ms/ms spectra. *Analytical Chemistry*, 97(29):15873–15882, Jul 2025. PMID: 40673560; PMCID: PMC12311886.
- [7] Gillian L. Daly and Frank Wania. Organic contaminants in mountains. *Environmental Science & Technology*, 39(2):385–398, 2005.
- [8] Madeleine Ernst, Kyo Bin Kang, Andrés Mauricio Caraballo-Rodríguez, Louis-Felix Nothias, Joe Wandy, Christopher Chen, Mingxun Wang, et al. MolNetEnhancer: Enhanced molecular networks by integrating metabolome mining and annotation tools. *Metabolites*, 9(7):144, 2019.
- [9] Wenkai Fan. Suspect screening and semi-quantification of organic contaminants in snow and water samples on the qinghai–tibet plateau and high mountain regions. Undergraduate honors thesis, The College of Wooster, Wooster, OH, 2025. Unpublished thesis.
- [10] Peter K.T. Newton S.R. et al. Fisher, C.M. Approaches for assessing performance of high-resolution mass spectrometry–based non-targeted analysis methods. *Anal Bioanal Chem*, 414:6455–6471, 2022.

- [11] Rebekah Gray, Eve Painter, Jameson W. Sprankle, Alex D. Crawford, Jillian J. Morrison, Marian Frazier, and Jennifer A. Faust. Suspect screening for pesticides in rain and snow using liquid chromatography high-resolution mass spectrometry. *Atmospheric Environment*, 291:119389, 2022.
- [12] B. M. J. Herbert, S. Villa, and C. J. Halsall. Chemical interactions with snow: Understanding the behavior and fate of semi-volatile organic compounds in snow. *Ecotoxicology and Environmental Safety*, 63(1):3–16, 2006.
- [13] Henriette Hupatz, Ingrid Rahu, Wen-Chao Wang, et al. Critical review on in silico methods for structural annotation of chemicals detected with lc/hrms non-targeted screening. *Analytical and Bioanalytical Chemistry*, 417(2):473–493, Jan 2025.
- [14] Tobias Kind, Hiroshi Tsugawa, Tomas Cajka, Yan Ma, Zijuan Lai, Sajjan S. Mehta, Gert Wohlgemuth, Dinesh Kumar Barupal, Megan R. Showalter, Masanori Arita, and Oliver Fiehn. Identification of small molecules using accurate mass MS/MS search. *Mass Spectrometry Reviews*, 37(4):513–532, 2018.
- [15] Aliz Kiss and Elke Fries. Occurrence of benzotriazoles in the rivers main, hengstbach, and hegbach (germany). *Environmental Science and Pollution Research*, 16(6):702–710, 2009.
- [16] Jeremy Koelmel, Nicholas Kroeger, Emily Gill, Candice Ulmer Holland, John Bowden, Rainey Patterson, Richard Yost, and Timothy Garrett. Expanding lipidome coverage using lc-ms/ms data-dependent acquisition with automated exclusion list generation. *Journal of The American Society for Mass Spectrometry*, 28:1–10, 03 2017.
- [17] Zhiyuan Ma, Shunlong Meng, Xin Gao, Hui Wan, Longxiang Fang, Lei Zhu, Zhonghua Li, Yuxiao Gao, Liping Qiu, Bo Rong, and Chao Song. Human activity driven spatial heterogeneity of pollutants in a highland river basin revealed by non-target screening and improved pbt evaluation: A case study from the yalong river on the qinghai-tibet plateau,china. *Journal of Hazardous Materials*, 496:139493, 2025.
- [18] K. E. Manz, A. Feerick, J. M. Braun, Y. L. Feng, A. Hall, J. Koelmel, C. Manzano, S. R. Newton, K. D. Pennell, B. J. Place, K. J. Godri Pollitt, C. Prasse, and J. A. Young. Non-targeted analysis (NTA) and suspect screening analysis (SSA): A review of examining the chemical exposome. *Journal of Exposure Science & Environmental Epidemiology*, 33(4):524–536, 2023.
- [19] Stephanie Nino-Suastegui, Eve Painter, Jameson W. Sprankle, Jillian J. Morrison, Jennifer A. Faust, and Rebekah Gray. Non-targeted analysis and suspect screening of organic contaminants in temperate snowfall using liquid chromatography high-resolution mass spectrometry. *Environmental Research*, 266:120494, 2025.

- [20] Florent Olivon, Gwendal Grelier, Fanny Roussi, Marc Litaudon, and David Touboul. MZmine 2 data preprocessing to enhance molecular networking reliability. *Analytical Chemistry*, 89(15):7836–7840, 2017.
- [21] Robin Schmid, Daniel Petras, Louis-Felix Nothias, Mingxun Wang, Allegra T. Aron, Annika Jagels, Hiroshi Tsugawa, et al. Ion identity molecular networking for mass spectrometry-based metabolomics in the GNPS environment. *Nature Communications*, 12:3832, 2021.
- [22] Bastian Schulze, Youngjoon Jeon, Sarit Kaserzon, Amy L. Heffernan, Pradeep Dewapriya, Jake O’Brien, Maria Jose Gomez Ramos, Sara Ghorbani Gorji, Jochen F. Mueller, Kevin V. Thomas, and Saer Samanipour. An assessment of quality assurance/quality control efforts in high resolution mass spectrometry non-target workflows for analysis of environmental samples. *TrAC Trends in Analytical Chemistry*, 133:116063, 2020.
- [23] Emma L. Schymanski, Junho Jeon, Rebekka Gulde, Kathrin Fenner, Matthias Ruff, Heinz P. Singer, and Juliane Hollender. Identifying small molecules via high resolution mass spectrometry: Communicating confidence. *Environmental Science & Technology*, 48(4):2097–2098, 2014.
- [24] Hiroshi Tsugawa, Tomas Cajka, Tobias Kind, Yan Ma, Brendan Higgins, Kazutaka Ikeda, Mitsuhiro Kanazawa, Jean VanderGheynst, and Masanori Arita. Ms-dial: Data-independent ms/ms deconvolution for comprehensive metabolome analysis. *Nature methods*, 12, 05 2015.
- [25] Mingxun Wang, Jeremy J. Carver, Vanessa V. Phelan, Laura M. Sanchez, Neha Garg, et al. Sharing and community curation of mass spectrometry data with global natural products social molecular networking. *Nature Biotechnology*, 34(8):828–837, 2016.
- [26] Yue Xu, Gan Zhang, Jun Li, Paromita Chakraborty, Hua Li, and Xiang Liu. Long-range atmospheric transport of persistent organochlorinated compounds to a remote mountain site in southwestern china. *Journal of Environmental Monitoring*, 13:3119–3127, 2011.
- [27] Weiting Ye, Jingcheng Li, and Xianfa Cai. MfGNN: Multi-scale feature-attentive graph neural networks for molecular property prediction. *Journal of Computational Chemistry*, 46(3):e70011, 2025.
- [28] Daniel Zahn, Hans Peter H. Arp, Kathrin Fenner, Anett Georgi, Jasmin Hafner, Sarah E. Hale, Juliane Hollender, Thomas Letzel, Emma L. Schymanski, Gabriel Sigmund, and Thorsten Reemtsma. Should transformation products change the way we manage chemicals? *Environmental Science & Technology*, 58(18):7710–7718, 2024.
- [29] Weimin Zhu, Yi Zhang, Duancheng Zhao, Jianrong Xu, and Ling Wang. HiGNN: A hierarchical informative graph neural network for molecular property prediction. *Journal of Chemical Information and Modeling*, 63(1):43–55, 2023.