

Computational Identification of Unknown Organic Contaminants and Molecules in Remote Environments via Molecular Networking

Raelyn Brooks

April 23, 2026

1 Goal

This project aims to characterize surface water samples from remote high-altitude environments by identifying both known and unknown organic contaminants using high-resolution mass spectrometry (HRMS). While database-driven approaches with lists of known contaminants enable identification of some contaminants, I hypothesize that a significant fraction of uncharacterized molecules can be identified through non-targeted analysis, particularly molecular networking. Together, these approaches will reveal the identity and patterns of contaminants in these remote regions.

2 Methods

2.1 Sample Collection and Preparation

Surface water samples collected from remote high-altitude environments—including Mt. Everest, Mt. Kilimanjaro, the Tibetan Plateau, and Qinghai Lake—will be analyzed for organic contaminants. These samples represent integrated records of atmospheric deposition. Prior to analysis, samples will be filtered to remove particulate matter and stored under controlled conditions to minimize degradation or contamination. Solid-phase extraction (SPE) will be used to concentrate dissolved organic compounds and improve detection sensitivity for trace-level contaminants. Extracted samples will then be reconstituted in an appropriate solvent compatible with liquid

chromatography–mass spectrometry (LC-HRMS) analysis.

2.2 High-Resolution Mass Spectrometry Data Acquisition

All samples will be analyzed using liquid chromatography coupled with high-resolution mass spectrometry (LC-HRMS). An iterative data-dependent acquisition (DDA) approach will be employed, in which each sample is injected and analyzed multiple times. In each successive run, previously fragmented precursor ions will be excluded using dynamic exclusion lists, allowing for increased fragmentation coverage of lower-abundance compounds. This iterative DDA strategy improves the detection of unique spectral features and enhances MS/MS coverage across complex environmental samples. Mass spectra will be collected in both MS1 (full scan) and MS/MS modes to obtain accurate mass measurements and fragmentation patterns necessary for compound identification.

2.3 Data Processing and Feature Detection

Raw LC-HRMS data will be processed using software such as MS-DIAL. Data processing steps will include peak detection, deconvolution, alignment across samples, and normalization. Detected features will be characterized by their mass-to-charge ratio (m/z), retention time, and MS/MS fragmentation spectra. Feature tables will be generated to represent the full set of detected compounds across all samples. Quality control steps, including blank subtraction and filtering of low-confidence features, will be applied to remove background contaminants and noise.

2.4 Suspect Screening for Known Contaminants

To identify known environmental contaminants, suspect screening will be performed using curated chemical databases, such as the Agilent environmental contaminant database. Detected features will be matched against database entries using accurate mass measurements and MS/MS spectral similarity. Candidate identifications will be assigned based on established confidence levels (e.g., Schymanski levels), incorporating mass accuracy, isotopic patterns, and fragmentation matching. This approach will enable the annotation of known compounds such as pharmaceuticals, pesticides, and industrial chemicals, providing a baseline characterization of anthropogenic contamination in the samples.

2.5 Molecular Networking and Non-Targeted Analysis

To characterize unknown compounds not identified through suspect screening, molecular networking will be performed using MS/MS spectral similarity. Spectral data will be uploaded to platforms such as GNPS (Global Natural Products Social Molecular Networking), where pairwise spectral comparisons will be used to construct molecular networks. In these networks, nodes represent individual compounds, and edges represent spectral similarity relationships. Clusters of related compounds will be analyzed to identify structural relationships between known and unknown molecules. Unknown compounds that cluster with annotated compounds can be tentatively characterized based on shared fragmentation patterns and mass differences. Additional annotation tools (e.g., MolNetEnhancer) may be used to integrate chemical class information and improve structural interpretation.

2.6 Structural Interpretation and Data Analysis

Mass differences between related compounds in molecular networks will be examined to infer potential chemical transformations (e.g., oxidation, methylation, or degradation products). Fragmentation patterns will be interpreted to propose tentative structures for unknown compounds. Comparative analysis across sampling locations will be conducted to identify trends in contaminant distribution and to assess the influence of long-range atmospheric transport. Differences in compound profiles may reveal region-specific contamination patterns or shared global sources.

References

- [1] Francesco Asnicar, Andrew Maltez Thomas, Andrea Passerini, Levi Waldron, and Nicola Segata. Machine learning for microbiologists. *Nature Reviews Microbiology*, 22(4):191–205, Apr 2024.
- [2] Jules M. Blais, Michael R. Rosen, and John P. Smol, editors. *Environmental Contaminants: Using Natural Archives to Track Sources and Long-Term Trends of Pollution*. Springer, Dordrecht, 2015.
- [3] Jordi. Catalan, editor. *Tracking Long-Range Atmospheric Transport of Trace Metals, Polycyclic Aromatic Hydrocarbons, and Organohalogen Compounds Using Lake Sediments of Mountain Regions*. Springer, Dordrecht, 2015.
- [4] Alex Chao, Jeffrey M. Minucci, Troy M. Ferland, E. Tyler Carr, Greg Janesch, Safia Rizwan, Heather D. Whitehead, Tommy Cathey, Shirley Pu, Laura D. Brunelle, Angela L. Batt, Jon R. Sobus, and Antony J. Williams. Prioritizing chemical candidates from non-targeted analysis using metadata, spectral similarity, and hazard scoring within INTERPRET NTA. *Analytical Chemistry*, 97(29):15904–15912, 2025.
- [5] Gillian L. Daly and Frank Wania. Organic contaminants in mountains. *Environmental Science & Technology*, 39(2):385–398, 2005.
- [6] Madeleine Ernst, Kyo Bin Kang, Andrés Mauricio Caraballo-Rodríguez, Louis-Felix Nothias, Joe Wandy, Christopher Chen, Mingxun Wang, et al. MolNetEnhancer: Enhanced molecular networks by integrating metabolome mining and annotation tools. *Metabolites*, 9(7):144, 2019.
- [7] Wenkai Fan. Suspect screening and semi-quantification of organic contaminants in snow and water samples on the qinghai–tibet plateau and high mountain regions. Undergraduate honors thesis, The College of Wooster, Wooster, OH, 2025. Unpublished thesis.
- [8] Peter K.T. Newton S.R. et al. Fisher, C.M. Approaches for assessing performance of high-resolution mass spectrometry–based non-targeted analysis methods. *Anal Bioanal Chem*, 414:6455–6471, 2022.
- [9] Rebekah Gray, Eve Painter, Jameson W. Sprankle, Alex D. Crawford, Jillian J. Morrison, Marian Frazier, and Jennifer A. Faust. Suspect screening for pesticides in rain and snow using liquid chromatography high-resolution mass spectrometry. *Atmospheric Environment*, 291:119389, 2022.
- [10] B. M. J. Herbert, S. Villa, and C. J. Halsall. Chemical interactions with snow: Understanding the behavior and fate of semi-volatile organic compounds in snow. *Ecotoxicology and Environmental Safety*, 63(1):3–16, 2006.

- [11] Henriette Hupatz, Ingrid Rahu, Wen-Chao Wang, et al. Critical review on in silico methods for structural annotation of chemicals detected with lc/hrms non-targeted screening. *Analytical and Bioanalytical Chemistry*, 417(2):473–493, Jan 2025.
- [12] Tobias Kind, Hiroshi Tsugawa, Tomas Cajka, Yan Ma, Zijuan Lai, Sajjan S. Mehta, Gert Wohlgemuth, Dinesh Kumar Barupal, Megan R. Showalter, Masanori Arita, and Oliver Fiehn. Identification of small molecules using accurate mass MS/MS search. *Mass Spectrometry Reviews*, 37(4):513–532, 2018.
- [13] Aliz Kiss and Elke Fries. Occurrence of benzotriazoles in the rivers main, hengstbach, and hegbach (germany). *Environmental Science and Pollution Research*, 16(6):702–710, 2009.
- [14] Zhiyuan Ma, Shunlong Meng, Xin Gao, Hui Wan, Longxiang Fang, Lei Zhu, Zhonghua Li, Yuxiao Gao, Liping Qiu, Bo Rong, and Chao Song. Human activity driven spatial heterogeneity of pollutants in a highland river basin revealed by non-target screening and improved pbt evaluation: A case study from the yalong river on the qinghai-tibet plateau,china. *Journal of Hazardous Materials*, 496:139493, 2025.
- [15] K. E. Manz, A. Feerick, J. M. Braun, Y. L. Feng, A. Hall, J. Koelmel, C. Manzano, S. R. Newton, K. D. Pennell, B. J. Place, K. J. Godri Pollitt, C. Prasse, and J. A. Young. Non-targeted analysis (NTA) and suspect screening analysis (SSA): A review of examining the chemical exposome. *Journal of Exposure Science & Environmental Epidemiology*, 33(4):524–536, 2023.
- [16] Stephanie Nino-Suastegui, Eve Painter, Jameson W. Sprankle, Jillian J. Morrison, Jennifer A. Faust, and Rebekah Gray. Non-targeted analysis and suspect screening of organic contaminants in temperate snowfall using liquid chromatography high-resolution mass spectrometry. *Environmental Research*, 266:120494, 2025.
- [17] Florent Olivon, Gwendal Grelier, Fanny Roussi, Marc Litaudon, and David Touboul. MZmine 2 data preprocessing to enhance molecular networking reliability. *Analytical Chemistry*, 89(15):7836–7840, 2017.
- [18] Robin Schmid, Daniel Petras, Louis-Felix Nothias, Mingxun Wang, Allegra T. Aron, Annika Jagels, Hiroshi Tsugawa, et al. Ion identity molecular networking for mass spectrometry-based metabolomics in the GNPS environment. *Nature Communications*, 12:3832, 2021.
- [19] Bastian Schulze, Youngjoon Jeon, Sarit Kaserzon, Amy L. Heffernan, Pradeep Dewapriya, Jake O’Brien, Maria Jose Gomez Ramos, Sara Ghorbani Gorji, Jochen F. Mueller, Kevin V. Thomas, and Saer Samanipour. An assessment of quality assurance/quality control efforts in high resolution mass spectrometry non-target workflows for analysis of environmental samples. *TrAC Trends in Analytical Chemistry*, 133:116063, 2020.

- [20] Emma L. Schymanski, Junho Jeon, Rebekka Gulde, Kathrin Fenner, Matthias Ruff, Heinz P. Singer, and Juliane Hollender. Identifying small molecules via high resolution mass spectrometry: Communicating confidence. *Environmental Science & Technology*, 48(4):2097–2098, 2014.
- [21] Mingxun Wang, Jeremy J. Carver, Vanessa V. Phelan, Laura M. Sanchez, Neha Garg, et al. Sharing and community curation of mass spectrometry data with global natural products social molecular networking. *Nature Biotechnology*, 34(8):828–837, 2016.
- [22] Yue Xu, Gan Zhang, Jun Li, Paromita Chakraborty, Hua Li, and Xiang Liu. Long-range atmospheric transport of persistent organochlorinated compounds to a remote mountain site in southwestern china. *Journal of Environmental Monitoring*, 13:3119–3127, 2011.
- [23] Weiting Ye, Jingcheng Li, and Xianfa Cai. MfGNN: Multi-scale feature-attentive graph neural networks for molecular property prediction. *Journal of Computational Chemistry*, 46(3):e70011, 2025.
- [24] Weimin Zhu, Yi Zhang, Duancheng Zhao, Jianrong Xu, and Ling Wang. HiGNN: A hierarchical informative graph neural network for molecular property prediction. *Journal of Chemical Information and Modeling*, 63(1):43–55, 2023.