

Computational Identification of Unknown Organic Contaminants and Molecules in Remote Environments via Molecular Networking

Raelyn Brooks

April 29, 2026

1 Goal

This project aims to characterize surface water samples from remote high-altitude environments by identifying both known and unknown organic contaminants using high-resolution mass spectrometry (HRMS). While database-driven approaches with lists of known contaminants enable identification of some contaminants, I hypothesize that a significant fraction of uncharacterized molecules can be identified through non-targeted analysis, particularly molecular networking. Together, these approaches will reveal the identity and patterns of contaminants in these remote regions.




Figure 1: Comparison of non-targeted analysis (NTA) and targeted analysis (TA) approaches for identification of organic contaminants [20].

2 Methods

2.1 High-Resolution Mass Spectrometry Data Acquisition

All samples will be analyzed using liquid chromatography coupled with high-resolution mass spectrometry (LC-HRMS). Separation will be performed using a C18 reverse-phase column, allowing for the retention and separation of semi-polar to nonpolar organic compounds commonly found in environmental samples.

Data will be acquired in both positive and negative electrospray ionization (ESI) modes to maximize compound coverage. For positive ionization mode, the mobile phase will consist of 0.1% formic acid in water (aqueous phase) and methanol with 0.1% formic acid (organic phase). For negative ionization mode, the mobile phase will consist of 10 mM ammonium acetate in water (aqueous phase) and 10 mM ammonium acetate in methanol (organic phase). Gradient elution will be used in both modes to effectively separate compounds across a range of polarities.

The analytical workflow will begin with MS1 full-scan acquisition to detect all ionizable features present in the samples. From these data, exclusion and inclusion lists will be generated. The exclusion list will contain background contaminants and redundant high-abundance ions to prevent repeated fragmentation, while the inclusion list will prioritize features of interest, particularly those matching entries in existing chemical database.

Subsequent analyses will employ MS/MS acquisition using an iterative data-dependent acquisition (DDA) strategy. In DDA, precursor ions are selected in real time based on their intensity for fragmentation. In the iterative approach used here, multiple injections of the same sample are performed, and previously fragmented ions are excluded in successive runs. This allows for deeper sampling of the chemical space by enabling fragmentation of lower-abundance features that would otherwise be missed. As a result, iterative DDA significantly increases MS/MS coverage and improves the likelihood of identifying both known and unknown compounds in complex environmental mixtures.




Figure 2: Example of exclusion and inclusion of features to establish our lists. [14]

2.2 Data Processing and Feature Detection

Raw LC-HRMS data will be processed using software such as MS-DIAL. Data processing will include peak detection, alignment across samples, and feature annotation.

Detected signals will initially be treated as features, defined by their mass-to-charge ratio (m/z), retention time, and associated MS/MS spectra. A feature is only considered a confirmed compound once sufficient evidence supports its identification.

Quality control steps, including blank subtraction and filtering of low-confidence signals, will be applied to remove background contaminants and noise. The resulting feature table will represent the set of detected molecular features across all samples and will serve as the basis for downstream identification and analysis.




Figure 3: This workflow illustrates data independent acquisition (DIA) using MS Dial while different from our data dependent acquisition (DDA) we are seeking the flow will be very similar. [?]

2.3 Suspect Screening for Known Contaminants

To identify contaminants of environmental concern (CECs), suspect screening will be performed using curated chemical databases, such as the Agilent environmental contaminant database.

Features will be matched to database entries using a combination of accurate mass, MS/MS spectral similarity, isotopic pattern, and retention time. Retention time provides an additional orthogonal parameter that improves confidence in matching beyond mass and fragmentation data alone.




Figure 4: This figure illustrates the MS/MS database creation and how LCMS is used to identify compounds based on their fragmentation patterns [12].

Rather than assigning definitive identifications, candidate annotations will be evaluated based on their level of confidence using the Schymanski framework as seen in Figure 3 [21]. These levels reflect how well a feature matches known reference data, ranging from tentative identifications (based on mass and fragmentation patterns) to confirmed identifications requiring comparison to authentic reference standards.

This approach enables identification of known contaminants while also highlighting gaps in existing chemical database, where detected features cannot be confidently matched to known compounds.




Figure 5: Confidence levels example for molecular feature identification based on the Schymanski framework. [21]

2.4 Molecular Networking and Non-Targeted Analysis

To extend beyond database-driven identification, molecular networking will be performed to characterize unknown features not identified through suspect screening. Instead of using existing platforms, a custom computational workflow will be developed to compare MS/MS spectra and identify relationships between structurally similar molecules.

Spectral similarity between features will be quantified using cosine similarity scoring. In this approach, each MS/MS spectrum is represented as a vector, where fragment ions (m/z values) are paired with their corresponding intensities. Before comparison, spectra are typically filtered and aligned so that only matching or near-matching fragment ions are considered. The cosine similarity score between two spectra is calculated as:

$$\text{Cosine Similarity} = \frac{\sum(A_i \cdot B_i)}{\sqrt{\sum(A_i^2)} \cdot \sqrt{\sum(B_i^2)}}$$

where A_i and B_i represent the intensities of corresponding fragment ions in the two spectra. This calculation effectively measures the cosine of the angle between the two spectral vectors in multidimensional space.

A score close to 1 indicates that two spectra have very similar fragmentation patterns, suggesting structural similarity between the corresponding molecules. Lower scores indicate less similarity and weaker structural relationships.

Importantly, cosine similarity accounts for both the presence of shared fragment ions and their relative intensities, making it more robust than simple peak matching. Additional considerations, such as allowing small mass tolerances between fragment ions and weighting higher-intensity peaks, can further improve the accuracy of similarity scoring.

Using this approach, groups of features with high spectral similarity will be identified and interpreted as structurally related compounds. Features that are closely related to annotated compounds can be used to infer the identity or chemical class of unknowns, enabling extension beyond existing chemical databases.




Figure 6: This example of Global Natural Product Social (GNPS) molecular networking illustrates the relationships between different molecular features but this is not fitted for environmental samples which causes gaps [19].

2.5 Structural Interpretation and Data Analysis

Structural interpretation will focus on analyzing mass differences and fragmentation patterns within molecular networks to propose tentative structures for unknown compounds.

Mass shifts between related features will be used to infer potential chemical transformations. For example:

- A mass increase of +16 Da may indicate the addition of an oxygen atom (e.g., hydroxylation).
- A mass shift of +14 Da may suggest methylation.
- A mass change of +16 Da combined with loss of 2 hydrogen atoms may indicate the formation of a carbonyl group.

Fragmentation patterns will be further analyzed to support these structural hypotheses. Through this approach, unknown features can be contextualized relative to known compounds, enabling extension beyond existing chemical libraries.

This combined workflow supports (1) comprehensive MS/MS data acquisition, (2) identification of known contaminants using existing databases, and (3) extension of chemical identification through molecular networking, directly aligning with the three specific aims of this study.




Figure 7: This diagram illustrates how a molecular network can be constructed from MS/MS data, enabling the identification of unknown compounds based on their fragmentation patterns while this is ion identity molecular networking a similar graph will be constructed [19].

References

- [1] Francesco Asnicar, Andrew Maltez Thomas, Andrea Passerini, Levi Waldron, and Nicola Segata. Machine learning for microbiologists. *Nature Reviews Microbiology*, 22(4):191–205, Apr 2024.
- [2] Jules M. Blais, Michael R. Rosen, and John P. Smol, editors. *Environmental Contaminants: Using Natural Archives to Track Sources and Long-Term Trends of Pollution*. Springer, Dordrecht, 2015.
- [3] Jordi. Catalan, editor. *Tracking Long-Range Atmospheric Transport of Trace Metals, Polycyclic Aromatic Hydrocarbons, and Organohalogen Compounds Using Lake Sediments of Mountain Regions*. Springer, Dordrecht, 2015.
- [4] Alex Chao, Jeffrey M. Minucci, Troy M. Ferland, E. Tyler Carr, Greg Janesch, Safia Rizwan, Heather D. Whitehead, Tommy Cathey, Shirley Pu, Laura D. Brunelle, Angela L. Batt, Jon R. Sobus, and Antony J. Williams. Prioritizing chemical candidates from non-targeted analysis using metadata, spectral similarity, and hazard scoring within INTERPRET NTA. *Analytical Chemistry*, 97(29):15904–15912, 2025.
- [5] Gillian L. Daly and Frank Wania. Organic contaminants in mountains. *Environmental Science & Technology*, 39(2):385–398, 2005.
- [6] Madeleine Ernst, Kyo Bin Kang, Andrés Mauricio Caraballo-Rodríguez, Louis-Felix Nothias, Joe Wandy, Christopher Chen, Mingxun Wang, et al. MolNetEnhancer: Enhanced molecular networks by integrating metabolome mining and annotation tools. *Metabolites*, 9(7):144, 2019.
- [7] Wenkai Fan. Suspect screening and semi-quantification of organic contaminants in snow and water samples on the qinghai–tibet plateau and high mountain regions. Undergraduate honors thesis, The College of Wooster, Wooster, OH, 2025. Unpublished thesis.
- [8] Peter K.T. Newton S.R. et al. Fisher, C.M. Approaches for assessing performance of high-resolution mass spectrometry–based non-targeted analysis methods. *Anal Bioanal Chem*, 414:6455–6471, 2022.
- [9] Rebekah Gray, Eve Painter, Jameson W. Sprankle, Alex D. Crawford, Jillian J. Morrison, Marian Frazier, and Jennifer A. Faust. Suspect screening for pesticides in rain and snow using liquid chromatography high-resolution mass spectrometry. *Atmospheric Environment*, 291:119389, 2022.
- [10] B. M. J. Herbert, S. Villa, and C. J. Halsall. Chemical interactions with snow: Understanding the behavior and fate of semi-volatile organic compounds in snow. *Ecotoxicology and Environmental Safety*, 63(1):3–16, 2006.

- [11] Henriette Hupatz, Ingrid Rahu, Wen-Chao Wang, et al. Critical review on in silico methods for structural annotation of chemicals detected with lc/hrms non-targeted screening. *Analytical and Bioanalytical Chemistry*, 417(2):473–493, Jan 2025.
- [12] Tobias Kind, Hiroshi Tsugawa, Tomas Cajka, Yan Ma, Zijuan Lai, Sajjan S. Mehta, Gert Wohlgemuth, Dinesh Kumar Barupal, Megan R. Showalter, Masanori Arita, and Oliver Fiehn. Identification of small molecules using accurate mass MS/MS search. *Mass Spectrometry Reviews*, 37(4):513–532, 2018.
- [13] Aliz Kiss and Elke Fries. Occurrence of benzotriazoles in the rivers main, hengstbach, and hegbach (germany). *Environmental Science and Pollution Research*, 16(6):702–710, 2009.
- [14] Jeremy Koelmel, Nicholas Kroeger, Emily Gill, Candice Ulmer Holland, John Bowden, Rainey Patterson, Richard Yost, and Timothy Garrett. Expanding lipidome coverage using lc-ms/ms data-dependent acquisition with automated exclusion list generation. *Journal of The American Society for Mass Spectrometry*, 28:1–10, 03 2017.
- [15] Zhiyuan Ma, Shunlong Meng, Xin Gao, Hui Wan, Longxiang Fang, Lei Zhu, Zhonghua Li, Yuxiao Gao, Liping Qiu, Bo Rong, and Chao Song. Human activity driven spatial heterogeneity of pollutants in a highland river basin revealed by non-target screening and improved pbt evaluation: A case study from the yalong river on the qinghai-tibet plateau,china. *Journal of Hazardous Materials*, 496:139493, 2025.
- [16] K. E. Manz, A. Feerick, J. M. Braun, Y. L. Feng, A. Hall, J. Koelmel, C. Manzano, S. R. Newton, K. D. Pennell, B. J. Place, K. J. Godri Pollitt, C. Prasse, and J. A. Young. Non-targeted analysis (NTA) and suspect screening analysis (SSA): A review of examining the chemical exposome. *Journal of Exposure Science & Environmental Epidemiology*, 33(4):524–536, 2023.
- [17] Stephanie Nino-Suastegui, Eve Painter, Jameson W. Sprankle, Jillian J. Morrison, Jennifer A. Faust, and Rebekah Gray. Non-targeted analysis and suspect screening of organic contaminants in temperate snowfall using liquid chromatography high-resolution mass spectrometry. *Environmental Research*, 266:120494, 2025.
- [18] Florent Olivon, Gwendal Grelier, Fanny Roussi, Marc Litaudon, and David Touboul. MZmine 2 data preprocessing to enhance molecular networking reliability. *Analytical Chemistry*, 89(15):7836–7840, 2017.
- [19] Robin Schmid, Daniel Petras, Louis-Felix Nothias, Mingxun Wang, Allegra T. Aron, Annika Jagels, Hiroshi Tsugawa, et al. Ion identity molecular networking for mass spectrometry-based metabolomics in the GNPS environment. *Nature Communications*, 12:3832, 2021.
- [20] Bastian Schulze, Youngjoon Jeon, Sarit Kaserzon, Amy L. Heffernan, Pradeep Dewapriya, Jake O’Brien, Maria Jose Gomez Ramos, Sara Ghorbani Gorji, Jochen F. Mueller, Kevin V.

- Thomas, and Saer Samanipour. An assessment of quality assurance/quality control efforts in high resolution mass spectrometry non-target workflows for analysis of environmental samples. *TrAC Trends in Analytical Chemistry*, 133:116063, 2020.
- [21] Emma L. Schymanski, Junho Jeon, Rebekka Gulde, Kathrin Fenner, Matthias Ruff, Heinz P. Singer, and Juliane Hollender. Identifying small molecules via high resolution mass spectrometry: Communicating confidence. *Environmental Science & Technology*, 48(4):2097–2098, 2014.
- [22] Hiroshi Tsugawa, Tomas Cajka, Tobias Kind, Yan Ma, Brendan Higgins, Kazutaka Ikeda, Mitsuhiro Kanazawa, Jean VanderGheynst, and Masanori Arita. Ms-dial: Data-independent ms/ms deconvolution for comprehensive metabolome analysis. *Nature methods*, 12, 05 2015.
- [23] Mingxun Wang, Jeremy J. Carver, Vanessa V. Phelan, Laura M. Sanchez, Neha Garg, et al. Sharing and community curation of mass spectrometry data with global natural products social molecular networking. *Nature Biotechnology*, 34(8):828–837, 2016.
- [24] Yue Xu, Gan Zhang, Jun Li, Paromita Chakraborty, Hua Li, and Xiang Liu. Long-range atmospheric transport of persistent organochlorinated compounds to a remote mountain site in southwestern china. *Journal of Environmental Monitoring*, 13:3119–3127, 2011.
- [25] Weiting Ye, Jingcheng Li, and Xianfa Cai. MfGNN: Multi-scale feature-attentive graph neural networks for molecular property prediction. *Journal of Computational Chemistry*, 46(3):e70011, 2025.
- [26] Daniel Zahn, Hans Peter H. Arp, Kathrin Fenner, Anett Georgi, Jasmin Hafner, Sarah E. Hale, Juliane Hollender, Thomas Letzel, Emma L. Schymanski, Gabriel Sigmund, and Thorsten Reemtsma. Should transformation products change the way we manage chemicals? *Environmental Science & Technology*, 58(18):7710–7718, 2024.
- [27] Weimin Zhu, Yi Zhang, Duancheng Zhao, Jianrong Xu, and Ling Wang. HiGNN: A hierarchical informative graph neural network for molecular property prediction. *Journal of Chemical Information and Modeling*, 63(1):43–55, 2023.