




jtools: Analysis and Presentation of Social Scientific Data

Jacob A. Long ¹

¹ School of Journalism and Mass Communications, University of South Carolina, USA

DOI: [10.xxxxxx/draft](https://doi.org/10.xxxxxx/draft)

Software

- [Review](#) 
- [Repository](#) 
- [Archive](#) 

Editor: 

Submitted: 08 January 2024

Published: unpublished

License

Authors of papers retain copyright¹² and release the work under a Creative Commons Attribution 4.0 International License ([CC BY 4.0](#))¹⁴.

Summary

jtools is an R package designed to ease the exploration and presentation of regression models with a focus on the needs of social scientists. Most notably, it generates results summaries that are meant to provide some of the conveniences of commercial software such as Stata (e.g., calculating robust standard errors and integrating them into a results table). Additionally, jtools includes plotting functions to help users better understand and share the results and predictions from fitted regression models.

Statement of need

Among the users of R are working or trainee scientists who are accustomed to software that provides detailed and customizable summaries of regression results “out of the box” (i.e., integrated into the modules that fit the model). While R’s `summary()` function provides useful information for many regression models — such as the method provided for `lm()` including coefficients, standard errors, test statistics, and *p*-values — there is typically little flexibility for customizing the output. Generating results summaries with additional information, such as robust standard errors, confidence intervals, transformed/centered variables, and variance inflation factors demands considerably more programming knowledge and effort for the user, requiring more time and making mistakes more likely. These same concerns exist for the case of plotting predicted values from regression models.

jtools provides an alternative to `summary()`, `summ()`, that allows users more flexibility in generating results tables for supported regression models. Importantly, these capabilities are accessible almost solely through arguments to `summ()`, thereby requiring little more programming knowledge than what is already required to fit a regression model and print a minimal summary to the console. For plotting, the package includes `effect_plot()` for creating line plots for a focal predictor variable. Users may add confidence intervals (including the ability to use robust standard errors for the plotted intervals), the observed data as plotted points, partial residuals as plotted points, and rug plots. Categorical predictors can be plotted as bar charts of single points with error bars as well. Finally, jtools features the function `plot_coefs()` that allows users to plot regression coefficients with confidence intervals for one or more models. This is particularly useful for comparing nested/related model specifications (Kastellec & Leoni, 2007). Plots are generated with `ggplot2` — allowing knowledgeable users to further customize the appearance — but do not require the user to know how to use `ggplot2`.

To support survey researchers, jtools functions generally support the use of sampling weights and survey design objects created by the survey package (Lumley, 2004). To fill in a gap in survey’s offerings, it adds `svycor()`, which calculates survey-weighted correlation matrices for survey design objects. In addition, `weights_tests()` implements the tests for the ignorability of sampling weights first devised by DuMouchel & Duncan (1983) and Pfeffermann & Sverchkov (1999). In layman’s terms, these tests allow researchers to check whether the use of sampling

weights to make data better resemble a population meaningfully change their statistical results compared to ignoring those weights.

Alternative software

Other R packages exist to achieve some of these ends. In some cases, `jtools` is using third-party packages for computation and simply repackaging the results (e.g., `sandwich` [Zeileis et al. (2020)] for robust standard errors and `pbkrtest` [Halekoh & Højsgaard (2014)] for computation of p -values for multilevel models). `modelsummary` (Arel-Bundock, 2022) provides comparable functionality to `summ()` and has some advantages, such as a greater range of supported models and more support for exporting to external documents. `marginalEffects` (Arel-Bundock, 2023), `sjPlot` (Lüdtke, n.d.), and see (Lüdtke et al., 2021) offer support for plotting predicted values from fitted regression models and again have some of their own advantages, such as more supported model types. There is also overlap in functionality with the `car` package (Fox & Weisberg, 2019), like the computation of variance inflation factors and partial residuals, although `jtools` aims to improve the user interface and support more model types in some cases.

Real-world use

At the time of writing, Google Scholar has tracked 556 references to `jtools`. These are predominantly, but not solely, in the social sciences, such as psychology, communication, and political science. For example, Sutin et al. (2023) uses `summ()` to calculate degrees of freedom for a multilevel model in a study of personality and aging. Urban-Wojcik et al. (2022) also use `summ()`, in this case to summarize regression models with robust standard errors in their study of physical activity and the hippocampus. Kraft et al. (2022) generate plotted regression summaries with `plot_coefs()` in a study of asthma sufferers. Finally, Spälti et al. (2023) use `weights_tests()` to assess the sensitivity of their estimates to the influence of survey sampling weights in their research on scientific misperceptions.

References

- Arel-Bundock, V. (2022). `modelsummary`: Data and model summaries in R. *Journal of Statistical Software*, 103(1), 1–23. <https://doi.org/10.18637/jss.v103.i01>
- Arel-Bundock, V. (2023). *MarginalEffects: Predictions, comparisons, slopes, marginal means, and hypothesis tests*. <https://CRAN.R-project.org/package=marginalEffects>
- DuMouchel, W. H., & Duncan, G. J. (1983). Using sample survey weights in multiple regression analyses of stratified samples. *Journal of the American Statistical Association*, 78, 535–543. <https://doi.org/10.2307/2288115>
- Fox, J., & Weisberg, S. (2019). *An R companion to applied regression* (Third). Sage. <https://socialsciences.mcmaster.ca/jfox/Books/Companion/>
- Halekoh, U., & Højsgaard, S. (2014). A kenward-roger approximation and parametric bootstrap methods for tests in linear mixed models – the R package `pbkrtest`. *Journal of Statistical Software*, 59(9), 1–30. <http://www.jstatsoft.org/v59/i09/>
- Kastellec, J. P., & Leoni, E. L. (2007). Using graphs instead of tables in political science. *Perspectives on Politics*, 5(4). <https://doi.org/10.1017/S1537592707072209>
- Kraft, M., Richardson, M., Hallmark, B., Billheimer, D., Van Den Berge, M., Fabbri, L. M., Van Der Molen, T., Nicolini, G., Papi, A., Rabe, K. F., Singh, D., Brightling, C., Siddiqui, S., Pizzichini, E., Cukier, A., Stelmach, R., Olivenstein, R., Zhang, Q., Badorrek, P., ... Hanania, N. (2022). The role of small airway dysfunction in asthma control and exacerbations: A

- 86 longitudinal, observational analysis using data from the ATLANTIS study. *The Lancet*
87 *Respiratory Medicine*, 10(7), 661–668. [https://doi.org/10.1016/S2213-2600\(21\)00536-1](https://doi.org/10.1016/S2213-2600(21)00536-1)
- 88 Lüdtke, D. (n.d.). *sjPlot: Data visualization for statistics in social science*. <https://CRAN.R-project.org/package=sjPlot>
- 90 Lüdtke, D., Patil, I., Ben-Shachar, M. S., Wiernik, B. M., Waggoner, P., & Makowski, D.
91 (2021). see: An R package for visualizing statistical models. *Journal of Open Source*
92 *Software*, 6(64), 3393. <https://doi.org/10.21105/joss.03393>
- 93 Lumley, T. (2004). Analysis of complex survey samples. *Journal of Statistical Software*, 9(1),
94 1–19.
- 95 Pfeiffermann, D., & Sverchkov, M. (1999). Parametric and semi-parametric estimation of
96 regression models fitted to survey data. *Sankhyā: The Indian Journal of Statistics*, 61,
97 166–186. <http://www.jstor.org/stable/25053074>
- 98 Späti, A. K., Lyons, B., Stoeckel, F., Stöckli, S., Szewach, P., Mérola, V., Stednitz, C., López
99 González, P., & Reifler, J. (2023). Partisanship and anti-elite worldviews as correlates of
100 science and health beliefs in the multi-party system of Spain. *Public Understanding of*
101 *Science*, 32(6), 761–780. <https://doi.org/10.1177/09636625231154131>
- 102 Sutin, A. R., Brown, J., Luchetti, M., Aschwanden, D., Stephan, Y., & Terracciano, A. (2023).
103 Five-factor model personality traits and the trajectory of episodic memory: Individual-
104 participant meta-analysis of 471,821 memory assessments from 120,640 participants. *The*
105 *Journals of Gerontology: Series B*, 78(3), 421–433. <https://doi.org/10.1093/geronb/gbac154>
- 106
- 107 Urban-Wojcik, E. J., Lee, S., Grupe, D. W., Quinlan, L., Gresham, L., Hammond, A., Charles, S.
108 T., Lachman, M. E., Almeida, D. M., Davidson, R. J., & Schaefer, S. M. (2022). Diversity
109 of daily activities is associated with greater hippocampal volume. *Cognitive, Affective, &*
110 *Behavioral Neuroscience*, 22(1), 75–87. <https://doi.org/10.3758/s13415-021-00942-5>
- 111 Zeileis, A., Köll, S., & Graham, N. (2020). Various versatile variances: An object-oriented
112 implementation of clustered covariances in R. *Journal of Statistical Software*, 95(1), 1–36.
113 <https://doi.org/10.18637/jss.v095.i01>