

# GRDzhadzha: A code for evolving relativistic matter on analytic metric backgrounds

Josu C. Aurrekoetxea<sup>1</sup>, Jamie Bamber<sup>1</sup>, Sam E. Brady<sup>2</sup>, Katy Clough<sup>2</sup>, Thomas Helfer<sup>3</sup>, James Marsden<sup>1</sup>, Miren Radia<sup>4</sup>, Dina Traykova<sup>5</sup>, and Zipeng Wang<sup>3</sup>

<sup>1</sup> Astrophysics, University of Oxford, Denys Wilkinson Building, Keble Road, Oxford OX1 3RH, United Kingdom <sup>2</sup> School of Mathematical Sciences, Queen Mary University of London, Mile End Road, London E1 4NS, United Kingdom <sup>3</sup> Henry A. Rowland Department of Physics & Astronomy, Johns Hopkins University, 3701 San Martin Drive, Baltimore, Maryland (MD) 21218, United States <sup>4</sup> Department of Applied Mathematics and Theoretical Physics (DAMTP), University of Cambridge, Centre for Mathematical Sciences, Wilberforce Road, Cambridge CB3 0WA, United Kingdom <sup>5</sup> Max Planck Institute for Gravitational Physics (Albert Einstein Institute), Am Mühlenberg 1, Potsdam-Golm, 14476, Germany

DOI: [10.xxxxxx/draft](https://doi.org/10.xxxxxx/draft)

## Software

- [Review](#)
- [Repository](#)
- [Archive](#)

Editor: [Warrick Ball](#)

## Reviewers:

- [@rashti-alireza](#)
- [@ekwessel](#)

Submitted: 16 August 2023

Published: unpublished

## License

Authors of papers retain copyright and release the work under a Creative Commons Attribution 4.0 International License ([CC BY 4.0](#)).

## Summary

Strong gravity environments such as those around black holes provide us with unique opportunities to study questions in fundamental physics (see e.g. ([Barack & others, 2019](#); [Barausse et al., 2015](#); [Bertone & others, 2020](#); [Macedo et al., 2013](#))), such as the existence and properties of dark matter and dark energy. Characterising the behaviour of new fields and other types of matter in highly relativistic environments usually necessitates numerical simulations unless one imposes significant symmetries. Therefore we need to turn to numerical methods to study the dynamics and evolution of the complex systems of black holes and other compact objects in different environments, using numerical relativity (NR). These methods allow us to split the four-dimensional Einstein equations into three-dimensional spatial hypersurfaces and a time-like direction. Then if a solution is known at the initial spatial hypersurface, it can be numerically evolved in time, where an analytic solution no longer exists. Whilst the tools of NR provide the most complete (i.e., approximation free) method for evolving matter in such environments, in many cases of interest, the density of the matter components is negligible in comparison to the curvature scales of the background spacetime metric ([Clough, 2021](#)). In such cases it is a reasonable approximation to neglect the backreaction of the matter environment onto the metric and treat it as fixed (assuming the background itself is stationary or otherwise has an analytic form).

In such cases, one does not need to evolve all the metric degrees of freedom as in NR, but only the additional matter ones. It is possible to do this using any NR code in a trivial way by setting the evolution of the metric variables to zero, but this is clearly rather inefficient. This code, GRDzhadzha, directly evolves the matter variables on an analytically specified background. This significantly speeds up the computation time and reduces the resources needed (both in terms of CPU hours and storage) to perform a given simulation. The code is based on the publicly available NR code GRChombo ([Andrade & others, 2021](#); [Clough et al., 2015](#)), which itself uses the open source Chombo framework ([Adams & others, 2015](#)) for solving PDEs.

In the following sections we discuss the key features and applications of the code, and give an indication of the efficiencies that can be achieved compared to a standard NR code.

## Key features of GRDzhadzha

GRDzhadzha inherits many of the features of GRChombo and Chombo, but avoids the complications introduced when evolving the metric. The key features are:

- **Background metrics:** The currently available backgrounds in the code are a static Kerr black hole in horizon penetrating Kerr-Schild coordinates and a boosted black hole in isotropic Schwarzschild coordinates. As the code is templated over the background, it can easily be changed or adapted to other coordinate systems for different problems without major code modification.
- **Matter evolution:** The code calculates the evolution for the matter variables on the metric background using an ADM decomposition (Arnowitt et al., 2008; York, 1978) in space and time - currently we have implemented a real and a complex scalar field as examples of matter types. Again the code is templated over the matter class so that the matter types can be exchanged with minimal modification.
- **Accuracy:** The metric values and their derivatives are calculated exactly at each point, whereas the matter fields are evolved with a 4th order Runge-Kutta time integration and their derivatives calculated with the same finite difference stencils used in GRChombo (4th and 6th order are currently available).
- **Boundary Conditions:** GRDzhadzha inherits all the available boundary conditions in GRChombo, namely, extrapolating (extrapolating the field value radially from values within the numerical grid), Sommerfeld (radiative) (Sommerfeld, 1912), reflective and periodic.
- **Initial Conditions:** The current examples provide initial data for real and complex scalar field matter. Since backreaction is ignored, there are no constraint equations to satisfy in the case of a scalar field, and the initial data can be freely specified.
- **Diagnostics:** GRDzhadzha has routines for verifying the conservation of matter energy densities, angular and linear momentum densities, and their fluxes, as discussed in (Clough, 2021; Croft, 2023).
- **C++ class structure:** Following the structure of GRChombo, GRDzhadzha is also written in C++ and uses object oriented programming (OOP) and templating.
- **Parallelism:** GRChombo uses hybrid OpenMP/MPI parallelism with explicit vectorisation of the evolution equations via intrinsics, and is AVX-512 compliant.
- **Adaptive Mesh Refinement:** The code inherits the flexible AMR grid structure of Chombo, which provides Berger-Oliger style (M. J. Berger & Oliger, 1984) AMR with block-structured Berger-Rigoutsos grid generation (M. Berger & Rigoutsos, 1991). Depending on the problem, the user may specify the refinement to be triggered by the matter or the background spacetime (Radia et al., 2022). One nice feature is that one does not need to resolve the horizon of the black hole unless matter is present at that location, so for an incoming wave a lot of storage and processing time can be saved by only resolving the wave, and not the spacetime background.

## Statement of Need

As mentioned in the introduction, any numerical relativity code like GRChombo can undertake these simulations. Examples of these include the Einstein Toolkit (<http://einstein toolkit.org/>), with its related Cactus (<http://cactuscode.org>) (Loffler & others, 2012; Schnetter et al., 2004), and Kranc (<http://kranccode.org>) (Husa et al., 2006) infrastructure used by LEAN (Sperhake, 2007; Zilhao et al., 2010) and Canuda (<https://bitbucket.org/canuda>) (Witek et al., 2019). Other notable but non-public codes include BAM (Bruegmann et al., 2008; Marronetti et al., 2007), AMSS-NCKU (Galaviz et al., 2010), PAMR/AMRD and HAD (East et al., 2012; Neilsen et al., 2007). Codes such as SPeC (Pfeiffer et al., 2003) and bamps (Hilditch et al., 2016) implement the generalised harmonic formulation of the Einstein equations using a pseudospectral method, and discontinuous Galerkin methods are

used in SpECTRE (<https://spectre-code.org>) (Cao et al., 2018; Deppe et al., 2021; Kidder & others, 2017). NRPy (<http://astro.phys.wvu.edu/bhathome>) (Ruchlin et al., 2018) is a code aimed for use on non-HPC systems, which generate C code from Python, and uses adapted coordinate systems to minimise computational costs. CosmoGRaPH (<https://cwru-pat.github.io/cosmograph>) (Mertens et al., 2016) and GRAMSES (Barrera-Hinojosa & Li, 2020) are among several NR codes targeted at cosmological applications (see (Adamek et al., 2020) for a comparison) and which also employ particle methods. Simflowny (<https://bitbucket.org/iac3/simflowny/wiki/Home>) (Palenzuela et al., 2018), like CosmoGRaPH, is based on the SAMRAI infrastructure, and has targeted fluid and MHD applications. GRATHENA++ (Daszuta et al., 2021) makes use of oct-tree AMR to maximise scaling.

Whilst there exist many NR codes (both public and private), which can in principle be used to perform simulations of fundamental fields on a fixed BH background, most do not have the efficiency advantages of GRDzhadza<sup>1</sup>. In particular, the fact that the ADM variables and their derivatives are not evolved or stored on the grid saves both a lot of simulation run time, as well as output file storage space. To get a rough idea of the improvement in storage and CPU hours one can achieve, we performed a short test simulation using GRDzhadza and compared it simulation performed using the full NR capabilities of GRChombo. We find that on average GRDzhadza is 15-20 times faster than GRChombo and requires about 3 times less file storage. An additional advantage of this code versus using a full NR code, for problems with negligible backreaction, is that here the metric variables are calculated analytically at every point on the grid, which significantly decreases the margin for numerical error, and means that resolution can be focussed on the matter location, and not the spacetime curvature.

It is important to note that whilst backreaction is neglected in the metric calculation, this does not mean that the backreaction effects cannot be calculated. Fixed background simulations provide a first order (in the density) estimate of the gravitational effects caused by the matter, taking into account their relativistic behaviour. This is discussed further in (Clough, 2021) and some examples using the approach are (Bamber et al., 2021; Traykova et al., 2021, 2023).


Since the interface and structure of the code is very close to the GRChombo numerical relativity code, it is possible for the results of these fixed background simulations to be used as initial data in full numerical relativity simulations (and vice versa), as was done in (Bamber et al., 2023). Therefore if the backreaction is found to be significant due to some growth mechanism, the simulation can be continued in full NR.

## Key research projects using GRDzhadza

So far the code has been used to study a range of fundamental physics problems:

- Studying the interference patterns in neutrino flavour oscillations around a static black hole (Alexandre & Clough, 2018)
- Growth of scalar hair around a Schwarzschild (Clough et al., 2019) and a Kerr (Bamber et al., 2021) black hole
- Determining the relativistic drag forces on a Schwarzschild black hole moving through a cloud of scalar field dark matter (Traykova et al., 2021, 2023)

<sup>1</sup>As far as we are aware, only NRPy and Canuda offer the same functionality. Some private codes also have such capabilities (see e.g. (Traykova et al., 2018), based on (Braden et al., 2015)). Other codes may have similar features that are not explicitly separated out, so this makes it difficult to identify them.



Some examples of the physics studied with GRDzhadzha. The left image is taken from (Traykova et al., 2021) in which the dynamical friction of light dark matter was studied in the relativistic regime. The right image is from a study of the scalar clouds around black hole binaries in (Bamber et al., 2023), in which the initial conditions were generated with a modified version of GRDzhadzha.

- Studying the dynamical friction effects on a Kerr black hole (Magnus effect) (Wang et al., 2023)
- Superradiance with self-interacting vector field (Clough et al., 2022) and with spatially varying mass (Wang et al., 2022)
- BH mergers in wave dark matter environments (Bamber et al., 2023)

## Acknowledgements

We thank the GRChombo collaboration (<http://www.grchombo.org>) for their support and code development work. JB and JM acknowledge funding from a UK Science and Technology Facilities Council (STFC) studentship. JCA acknowledges funding from the Beecroft Trust and The Queen's College via an extraordinary Junior Research Fellowship (eJRF). KC acknowledges funding from the UKRI Ernest Rutherford Fellowship (grant number ST/V003240/1). SEB is supported by a QMUL Principal studentship. ZW is supported by NSF Grants No.~AST-2006538, PHY-2207502, PHY-090003 and PHY-20043, and by NASA Grants No. 20-LPS20-0011 and 21-ATP21-0010. A few of the projects using this code have received funding from the European Research Council (ERC) under the European Union's Horizon 2020 research and innovation programme (Grant Agreement No 693024).

GRDzhadzha users have benefited from the provision of HPC resources from:

- DiRAC (Distributed Research utilising Advanced Computing) resources under the projects ACSP218, ACSP191, ACTP183, ACTP186 and ACTP316. Systems used include:
  - Cambridge Service for Data Driven Discovery (CSD3), part of which is operated by the University of Cambridge Research Computing on behalf of the STFC DiRAC HPC Facility ([www.dirac.ac.uk](http://www.dirac.ac.uk)). The DiRAC component of CSD3 was funded by BEIS capital funding via STFC capital grants ST/P002307/1 and ST/R002452/1 and STFC operations grant ST/R00689X/1. DiRAC is part of the National e-Infrastructure.
  - DiRAC Data Intensive service at Leicester, operated by the University of Leicester IT Services, which forms part of the STFC DiRAC HPC Facility ([www.dirac.ac.uk](http://www.dirac.ac.uk)). The equipment was funded by BEIS capital funding via STFC capital grants ST/K000373/1 and ST/R002363/1 and STFC DiRAC Operations grant ST/R001014/1. DiRAC is part of the National e-Infrastructure.

- DiRAC at Durham facility managed by the Institute for Computational Cosmology on behalf of the STFC DiRAC HPC Facility ([www.dirac.ac.uk](http://www.dirac.ac.uk)). The equipment was funded by BEIS capital funding via STFC capital grants ST/P002293/1 and ST/R002371/1, Durham University and STFC operations grant ST/R000832/1. DiRAC is part of the National e-Infrastructure.
- DiRAC Complexity system, operated by the University of Leicester IT Services, which forms part of the STFC DiRAC HPC Facility ([www.dirac.ac.uk](http://www.dirac.ac.uk)). This equipment is funded by BIS National E-Infrastructure capital grant ST/K000373/1 and STFC DiRAC Operations grant ST/K0003259/1. DiRAC is part of the National e-Infrastructure.
- Sakura, Cobra and Raven clusters at the Max Planck Computing and Data Facility (MPCDF) in Garching, Germany,
- PRACE (Partnership for Advanced Computing in Europe) resources under grant numbers 2018194669, 2020225359. Systems used include:
  - SuperMUCNG, Leibniz Supercomputing Center (LRZ), Germany
  - JUWELS, Juelich Supercomputing Centre (JSC), Germany
- the Argonne Leadership Computing Facility, including the Joint Laboratory for System Evaluation (JLSE), which is a U.S. Department of Energy (DOE) Office of Science User Facility supported under Contract DE-AC02-06CH11357.
- the Texas Advanced Computing Center (TACC) at the University of Austin HPC and visualization resources URL: <http://www.tacc.utexas.edu>, and the San Diego Supercomputing Center (SDSC) URL: <https://www.sdsc.edu>, under project PHY-20043 and XSEDE Grant No. NSF-PHY-090003
- the Glamdring cluster, Astrophysics, Oxford, UK.

## References

- Adamek, J., Barrera-Hinojosa, C., Bruni, M., Li, B., Macpherson, H. J., & Mertens, J. B. (2020). Numerical solutions to Einstein's equations in a shearing-dust Universe: a code comparison. *Class. Quant. Grav.*, 37(15), 154001. <https://doi.org/10.1088/1361-6382/ab939b>
- Adams, M., & others. (2015). *Chombo software package for AMR applications - design document*.
- Alexandre, J., & Clough, K. (2018). Black hole interference patterns in flavor oscillations. *Phys. Rev. D*, 98(4), 043004. <https://doi.org/10.1103/PhysRevD.98.043004>
- Andrade, T., & others. (2021). GRChombo: An adaptable numerical relativity code for fundamental physics. *J. Open Source Softw.*, 6(68), 3703. <https://doi.org/10.21105/joss.03703>
- Arnowitt, R. L., Deser, S., & Misner, C. W. (2008). The Dynamics of general relativity. *Gen. Rel. Grav.*, 40, 1997–2027. <https://doi.org/10.1007/s10714-008-0661-1>
- Bamber, J., Aurrekoetxea, J. C., Clough, K., & Ferreira, P. G. (2023). Black hole merger simulations in wave dark matter environments. *Phys. Rev. D*, 107(2), 024035. <https://doi.org/10.1103/PhysRevD.107.024035>
- Bamber, J., Clough, K., Ferreira, P. G., Hui, L., & Lagos, M. (2021). Growth of accretion driven scalar hair around Kerr black holes. *Phys. Rev. D*, 103(4), 044059. <https://doi.org/10.1103/PhysRevD.103.044059>



- Barack, L., & others. (2019). Black holes, gravitational waves and fundamental physics: a roadmap. *Class. Quant. Grav.*, 36(14), 143001. <https://doi.org/10.1088/1361-6382/ab0587>
- Barausse, E., Cardoso, V., & Pani, P. (2015). Environmental Effects for Gravitational-wave Astrophysics. *J. Phys. Conf. Ser.*, 610(1), 012044. <https://doi.org/10.1088/1742-6596/610/1/012044>
- Barrera-Hinojosa, C., & Li, B. (2020). GRAMES: a new route to general relativistic  $N$ -body simulations in cosmology. Part I. Methodology and code description. *JCAP*, 01, 007. <https://doi.org/10.1088/1475-7516/2020/01/007>
- Berger, M. J., & Oliger, J. (1984). Adaptive Mesh Refinement for Hyperbolic Partial Differential Equations. *J. Comput. Phys.*, 53, 484. [https://doi.org/10.1016/0021-9991\(84\)90073-1](https://doi.org/10.1016/0021-9991(84)90073-1)
- Berger, M., & Rigoutsos, I. (1991). An algorithm for point clustering and grid generation. *IEEE Transactions on Systems, Man, and Cybernetics*, 21(5), 1278–1286. <https://doi.org/10.1109/21.120081>
- Bertone, G., & others. (2020). Gravitational wave probes of dark matter: challenges and opportunities. *SciPost Phys. Core*, 3, 007. <https://doi.org/10.21468/SciPostPhysCore.3.2.007>
- Braden, J., Bond, J. R., & Mersini-Houghton, L. (2015). Cosmic bubble and domain wall instabilities I: parametric amplification of linear fluctuations. *JCAP*, 03, 007. <https://doi.org/10.1088/1475-7516/2015/03/007>
- Bruegmann, B., Gonzalez, J. A., Hannam, M., Husa, S., Sperhake, U., & Tichy, W. (2008). Calibration of Moving Puncture Simulations. *Phys. Rev. D*, 77, 024027. <https://doi.org/10.1103/PhysRevD.77.024027>
- Cao, Z., Fu, P., Ji, L.-W., & Xia, Y. (2018). Application of local discontinuous Galerkin method to Einstein equations. *Int. J. Mod. Phys. D*, 28(01), 1950014. <https://doi.org/10.1142/S0218271819500147>
- Clough, K. (2021). Continuity equations for general matter: applications in numerical relativity. *Class. Quant. Grav.*, 38(16), 167001. <https://doi.org/10.1088/1361-6382/ac10ee>
- Clough, K., Ferreira, P. G., & Lagos, M. (2019). Growth of massive scalar hair around a Schwarzschild black hole. *Phys. Rev. D*, 100(6), 063014. <https://doi.org/10.1103/PhysRevD.100.063014>
- Clough, K., Figueras, P., Finkel, H., Kunesch, M., Lim, E. A., & Tunyasuvunakool, S. (2015). GRChombo : Numerical Relativity with Adaptive Mesh Refinement. *Class. Quant. Grav.*, 32(24), 245011. <https://doi.org/10.1088/0264-9381/32/24/245011>
- Clough, K., Helfer, T., Witek, H., & Berti, E. (2022). Ghost Instabilities in Self-Interacting Vector Fields: The Problem with Proca Fields. *Phys. Rev. Lett.*, 129(15), 151102. <https://doi.org/10.1103/PhysRevLett.129.151102>
- Croft, R. (2023). Local continuity of angular momentum and noether charge for matter in general relativity. *Class. Quant. Grav.*, 40(10), 105007. <https://doi.org/10.1088/1361-6382/accc6a>
- Daszuta, B., Zappa, F., Cook, W., Radice, D., Bernuzzi, S., & Morozova, V. (2021). GR-Athena++: Puncture Evolutions on Vertex-centered Oct-tree Adaptive Mesh Refinement. *Astrophys. J. Supp.*, 257(2), 25. <https://doi.org/10.3847/1538-4365/ac157b>
- Deppe, N., Throwe, W., Kidder, L. E., Fischer, N. L., Armaza, C., Bonilla, G. S., Hébert, F., Kumar, P., Lovelace, G., Moxon, J., O'Shea, E., Pfeiffer, H. P., Scheel, M. A., Teukolsky, S. A., Anantpurkar, I., Boyle, M., Foucart, F., Giesler, M., Izzo, D. A. B., ... Włodarczyk,

- 258 T. (2021). *SpECTRE* (Version 2021.05.03). Zenodo. [https://doi.org/10.5281/zenodo.](https://doi.org/10.5281/zenodo.4734670)  
259 [4734670](https://doi.org/10.5281/zenodo.4734670)
- 260 East, W. E., Pretorius, F., & Stephens, B. C. (2012). Hydrodynamics in full general relativity  
261 with conservative AMR. *Phys. Rev. D*, 85, 124010. [https://doi.org/10.1103/PhysRevD.](https://doi.org/10.1103/PhysRevD.85.124010)  
262 [85.124010](https://doi.org/10.1103/PhysRevD.85.124010)
- 263 Galaviz, P., Bruegmann, B., & Cao, Z. (2010). Numerical evolution of multiple black holes  
264 with accurate initial data. *Phys. Rev. D*, 82, 024005. [https://doi.org/10.1103/PhysRevD.](https://doi.org/10.1103/PhysRevD.82.024005)  
265 [82.024005](https://doi.org/10.1103/PhysRevD.82.024005)
- 266 Hilditch, D., Weyhausen, A., & Brügmann, B. (2016). Pseudospectral method for gravitational  
267 wave collapse. *Phys. Rev. D*, 93(6), 063006. [https://doi.org/10.1103/PhysRevD.93.](https://doi.org/10.1103/PhysRevD.93.063006)  
268 [063006](https://doi.org/10.1103/PhysRevD.93.063006)
- 269 Husa, S., Hinder, I., & Lechner, C. (2006). Kranc: A Mathematica application to generate  
270 numerical codes for tensorial evolution equations. *Comput. Phys. Commun.*, 174, 983–1004.  
271 <https://doi.org/10.1016/j.cpc.2006.02.002>
- 272 Kidder, L. E., & others. (2017). SpECTRE: A Task-based Discontinuous Galerkin Code for  
273 Relativistic Astrophysics. *J. Comput. Phys.*, 335, 84–114. [https://doi.org/10.1016/j.jcp.](https://doi.org/10.1016/j.jcp.2016.12.059)  
274 [2016.12.059](https://doi.org/10.1016/j.jcp.2016.12.059)
- 275 Löffler, F., & others. (2012). The Einstein Toolkit: A Community Computational Infrastructure  
276 for Relativistic Astrophysics. *Class. Quant. Grav.*, 29, 115001. [https://doi.org/10.1088/](https://doi.org/10.1088/0264-9381/29/11/115001)  
277 [0264-9381/29/11/115001](https://doi.org/10.1088/0264-9381/29/11/115001)
- 278 Macedo, C. F. B., Pani, P., Cardoso, V., & Crispino, L. C. B. (2013). Into the lair: gravitational-  
279 wave signatures of dark matter. *Astrophys. J.*, 774, 48. [https://doi.org/10.1088/](https://doi.org/10.1088/0004-637X/774/1/48)  
280 [0004-637X/774/1/48](https://doi.org/10.1088/0004-637X/774/1/48)
- 281 Marronetti, P., Tichy, W., Bruegmann, B., Gonzalez, J., Hannam, M., Husa, S., & Sperhake,  
282 U. (2007). Binary black holes on a budget: Simulations using workstations. *Class. Quant.*  
283 *Grav.*, 24, S43–S58. <https://doi.org/10.1088/0264-9381/24/12/S05>
- 284 Mertens, J. B., Giblin, J. T., & Starkman, G. D. (2016). Integration of inhomogeneous  
285 cosmological spacetimes in the BSSN formalism. *Phys. Rev. D*, 93(12), 124059. <https://doi.org/10.1103/PhysRevD.93.124059>  
286 <https://doi.org/10.1103/PhysRevD.93.124059>
- 287 Neilsen, D., Hirschmann, E. W., Anderson, M., & Liebling, S. L. (2007). Adaptive Mesh  
288 Refinement and Relativistic MHD. *11th Marcel Grossmann Meeting on General Relativity*,  
289 1579–1581. [https://doi.org/10.1142/9789812834300\\_0200](https://doi.org/10.1142/9789812834300_0200)
- 290 Palenzuela, C., Miñano, B., Viganò, D., Arbona, A., Bona-Casas, C., Rigo, A., Bezares,  
291 M., Bona, C., & Massó, J. (2018). A Simflowny-based finite-difference code for high-  
292 performance computing in numerical relativity. *Class. Quant. Grav.*, 35(18), 185007.  
293 <https://doi.org/10.1088/1361-6382/aad7f6>
- 294 Pfeiffer, H. P., Kidder, L. E., Scheel, M. A., & Teukolsky, S. A. (2003). A Multidomain  
295 spectral method for solving elliptic equations. *Comput. Phys. Commun.*, 152, 253–273.  
296 [https://doi.org/10.1016/S0010-4655\(02\)00847-0](https://doi.org/10.1016/S0010-4655(02)00847-0)
- 297 Radia, M., Sperhake, U., Drew, A., Clough, K., Figueras, P., Lim, E. A., Ripley, J. L.,  
298 Aurrekoetxea, J. C., França, T., & Helfer, T. (2022). Lessons for adaptive mesh refinement  
299 in numerical relativity. *Class. Quant. Grav.*, 39(13), 135006. [https://doi.org/10.1088/](https://doi.org/10.1088/1361-6382/ac6fa9)  
300 [1361-6382/ac6fa9](https://doi.org/10.1088/1361-6382/ac6fa9)
- 301 Ruchlin, I., Etienne, Z. B., & Baumgarte, T. W. (2018). SENR/NRP+: Numerical Relativity  
302 in Singular Curvilinear Coordinate Systems. *Phys. Rev. D*, 97(6), 064036. <https://doi.org/10.1103/PhysRevD.97.064036>  
303 <https://doi.org/10.1103/PhysRevD.97.064036>

- 304 Schnetter, E., Hawley, S. H., & Hawke, I. (2004). Evolutions in 3-D numerical relativity using  
 305 fixed mesh refinement. *Class. Quant. Grav.*, 21, 1465–1488. [https://doi.org/10.1088/](https://doi.org/10.1088/0264-9381/21/6/014)  
 306 [0264-9381/21/6/014](https://doi.org/10.1088/0264-9381/21/6/014)
- 307 Sommerfeld, A. (1912). Die greensche funktion der schwingungsgleichung. *Jahresbericht Der*  
 308 *Deutschen Mathematiker-Vereinigung*, 21, 309–352. <http://eudml.org/doc/145344>
- 309 Sperhake, U. (2007). Binary black-hole evolutions of excision and puncture data. *Phys. Rev.*  
 310 *D*, 76, 104015. <https://doi.org/10.1103/PhysRevD.76.104015>
- 311 Traykova, D., Braden, J., & Peiris, H. V. (2018). Accretion of a Symmetry Breaking Scalar  
 312 Field by a Schwarzschild Black Hole. *Phil. Trans. Roy. Soc. Lond. A*, 376(2114),  
 313 20170122. <https://doi.org/10.1098/rsta.2017.0122>
- 314 Traykova, D., Clough, K., Helfer, T., Berti, E., Ferreira, P. G., & Hui, L. (2021). Dynamical  
 315 friction from scalar dark matter in the relativistic regime. *Phys. Rev. D*, 104(10), 103014.  
 316 <https://doi.org/10.1103/PhysRevD.104.103014>
- 317 Traykova, D., Vicente, R., Clough, K., Helfer, T., Berti, E., Ferreira, P. G., & Hui, L.  
 318 (2023). *Relativistic drag forces on black holes from scalar dark matter clouds of all sizes.*  
 319 <https://arxiv.org/abs/2305.10492>
- 320 Wang, Z., Helfer, T., Clough, K., & Berti, E. (2022). Superradiance in massive vector fields  
 321 with spatially varying mass. *Phys. Rev. D*, 105(10), 104055. [https://doi.org/10.1103/](https://doi.org/10.1103/PhysRevD.105.104055)  
 322 [PhysRevD.105.104055](https://doi.org/10.1103/PhysRevD.105.104055)
- 323 Wang, Z., Helfer, T., Traykova, D., Clough, K., & Berti, E. (2023). The Magnus Effect on  
 324 black holes in scalar dark matter environments. *In Preparation.*
- 325 Witek, H., Gualtieri, L., Pani, P., & Sotiriou, T. P. (2019). Black holes and binary mergers  
 326 in scalar Gauss-Bonnet gravity: scalar field dynamics. *Phys. Rev. D*, 99(6), 064035.  
 327 <https://doi.org/10.1103/PhysRevD.99.064035>
- 328 York, J. W., Jr. (1978). Kinematics and Dynamics of General Relativity. *Workshop on Sources*  
 329 *of Gravitational Radiation*, 83–126.
- 330 Zilhao, M., Witek, H., Sperhake, U., Cardoso, V., Gualtieri, L., Herdeiro, C., & Nerozzi, A.  
 331 (2010). Numerical relativity for D dimensional axially symmetric space-times: formalism  
 332 and code tests. *Phys. Rev. D*, 81, 084052. <https://doi.org/10.1103/PhysRevD.81.084052>