

**Genome wide association study using phenotypes estimated from a kinetic-pharmacodynamic model of chemotherapy-induced peripheral neuropathy (CIPN)**

Bhongsatiern, J.,<sup>1</sup> Carbonetto, P.,<sup>2</sup> Chua, K.,<sup>3</sup> Mehrotra, S.,<sup>4</sup> Kroetz, D.,<sup>3</sup> Gobburu, J.V.,<sup>4</sup> Cox, N.J.,<sup>5</sup> Sharma, M.R.,<sup>1</sup>

<sup>1</sup>Department of Medicine, University of Chicago, Chicago, IL

<sup>2</sup>Research Computing Center, University of Chicago, Chicago, IL

<sup>3</sup>Department of Engineering and Therapeutic Sciences, University of California San Francisco, San Francisco, CA

<sup>4</sup>Center for Translational Medicine, University of Maryland, Baltimore, MD

<sup>5</sup>Division of Genetic Medicine, Vanderbilt University, Nashville, TN

Chemotherapy-induced peripheral neuropathy (CIPN) is a common toxicity caused by exposure to anticancer agents that impairs quality of life. Recently, a kinetic-pharmacodynamic model has been developed to quantitate the dose-CIPN relationship using data on patient-reported CIPN in a randomized phase III trial of paclitaxel, nab-paclitaxel and ixabepilone in metastatic breast cancer. Five parameters (BASE, IRG, KDE, KIN, SLOPE) were estimated in this model. Parameter estimates had relatively large between-subject variability, and genetic variation is hypothesized to describe some of this variability. A genome-wide association study was done using the model-estimated parameters as phenotypes (n=524). KDE was significantly associated with rs11952896 in patients of European ancestry ( $p < 5 \times 10^{-8}$ ), but was not significant with log-transformed KDE. Future directions are to explore whether groups of common variant SNPs and/or gene expression are associated with parameter estimates.

(989 including spaces)