

# Quantum spin defects in industrially important semiconductors for scalable solid-state quantum technology




H. Seo, A. L. Falk, P. V. Klimov, K. C. Miao, W. F. Koehl, M. Govoni, H. Ma, D. D. Awschalom and G. Galli  
 Institute for Molecular Engineering, University of Chicago, Chicago, IL 60615

## Defect qubits in solids: NV in diamond vs. Divacancy in SiC

### Nitrogen-vacancy (NV) center in diamond: optically addressable single spin<sup>1</sup>

#### Electronic structure


- Robust S=1 spin-triplet ground state
- Zero-field splitting (2.9 GHz)




- The NV center is being developed for quantum information processing, quantum sensors in physical and biological systems, and hybrid quantum systems<sup>2</sup>.

#### Optically detected magnetic resonance (ODMR)


- Intra-defect excitation and intersystem crossing
- Microwave control of spin transitions
- Spin-dependent photoluminescence



### Divacancy in SiC: NV-analogs in industrial materials<sup>3-9</sup>



- Excellent materials systems for optoelectronic devices, micromechanical systems and power electronics industry.
- Advanced material technologies are available.
- Wafer-scale growth
- Doping and device fabrication



- Common features**
- C<sub>3v</sub>, 6 e<sup>-</sup>, S=1<sup>4</sup>
- Optically addressable<sup>6</sup>
- Single-spin isolation<sup>9</sup>
- Room-T operation<sup>6</sup>

- Recent developments of SiC divacancy includes: dynamic nuclear polarization and electrical control, etc<sup>7</sup>.
- Related systems in SiC include silicon vacancy<sup>8</sup>.

**The divacancy coherence times have been measured much longer than that of the NV center in diamond:** 1.2 ms for divacancy in SiC<sup>9</sup> (T = 20 K) and 0.63 ms for NV in diamond<sup>10</sup> (Room T.).


## First-principles design of new spin qubits in piezoelectric AlN: Motivation and methods

### Industrial materials + coherent spin qubits → scalable solid-state quantum devices<sup>3,7</sup>

#### Material

Material	Band gap (eV)
Diamond	5.5
4H-SiC	3.3
w-AlN	6.2
GaN	3.4
ZnO	3.4
Si	1.1

We consider w-AlN as a new potential host material for defect qubits<sup>15,16</sup>.




**Question:** Can we identify a NV-analog in piezoelectric w-AlN?  
**Method:** Materials design from first-principles!

We use density functional theory to predict and design new defect spin qubits.

#### Qubit Requirements

- Triplet spin, Localization, Intra-defect excitation, ...

#### First-principles Computational Modeling



#### Spectroscopy

- EPR, Photoluminescence, ...

Optical transition energies, Reasonable parameter range, ...

**WEST!**  
<http://www.west-code.org/>

#### Materials Issues

$$v_{xc}(r, r') = \alpha v_{xc}^{ex}(r, r') + (1 - \alpha)v_x(r) + v_c(r)$$

Which  $\alpha$  for solids? Relation ship  $\alpha = \epsilon_{\infty}^{-1}$  is inferred from:

- Empirical observations<sup>18</sup>
- Many-body perturbation theory<sup>17</sup> (static COHSEX approximation).

→  $\alpha$  is self-consistently determined<sup>17</sup>. For AlN  $\alpha = \epsilon_{\infty}^{-1} = 0.25$

#### GW many-body perturbation theory<sup>19</sup>


$$E_n^{QP} = \epsilon_n^{KS} + \langle (\Sigma E_n^{QP})_n \rangle - \langle V_{xc} \rangle$$

$$\Sigma^{GW} = iGW$$


## First-principles design of new spin qubits in piezoelectric AlN: Results

### Negatively charged N vacancy under strain is a strong candidate for qubits in AlN.


#### Defect formation energy<sup>20</sup> of V<sub>N</sub> and O<sub>N</sub>




Uniaxial strain is applied to the host lattice.



$$\Delta E = E(S=1) - E(S=0)$$



#### Defect level diagram (G<sub>0</sub>W<sub>0</sub>@PBE)



## Conclusions, Reference, and Acknowledgement

### Quantum decoherence dynamics of divacancy spins in 4H-SiC<sup>11</sup>


- We showed theoretically and experimentally that the  $T_2$  time of the divacancy reaches 1.3 ms.
- We found that the robust coherence in SiC stems from the combination of the two effects:
  - The <sup>29</sup>Si and <sup>13</sup>C spin baths are decoupled.
  - The active homonuclear spins are much further apart than those in diamond.
- Our results point to polyatomic crystals as promising hosts for coherent qubits in solids.

### Reference

- F. Jelezko, T. Gaebel, I. Popa, A. Gruber, J. Wrachtrup, *Phys. Rev. Lett.* **92**, 076401 (2004).
- F. J. Heremans, C. G. Vale, and D. D. Awschalom, *Proc. IEEE*, early access (2016).
- J.R. Weber, W.F. Koehl, J.B. Varley, A. Janotti, B.B. Buckley, C.G. Van de Walle, and D.D. Awschalom, *PNAS* **107**, 8513 (2010).
- A. Galli, *Phys. Stat. Solidi B* **248**, 1337 (2011).
- P.G. Baranov, Y.I. Il'in, E.N. Mokhov, M.V. Muzafarov, S.B. Orlinskii, J. Schmidt, *IETP Lett.* **82**, 441 (2005).
- W.F. Koehl, B.B. Buckley, F.J. Heremans, G. Calusine, and D.D. Awschalom, *Nature* **84**, 479 (2011).
- W.F. Koehl, H. Seo, G. Galli, and D.D. Awschalom, *MRS Bulletin* **40**, 1146 (2015).
- M. Widmann et al., *Nat. Mater.* (2015); Yang et al., *PRB* (2014); Soltamov et al., *PRL* (2012); Simin et al., *PRX* **6**, 031014 (2016).
- D.J. Christie, A.L. Falk, P. Andrich, P.V. Klimov, J.U. Hassan, N.T. Son, E. Janzen, T. Ohshima, and D.D. Awschalom, *Nat. Mater.* **14**, 160 (2014).
- P.L. Stanwix, L.M. Pham, J.R. Maze, D. Le Sage, T.K. Yeung, P. Cappellaro, P.R. Hemmer, A. Yacoby, M. D. Lukin, and R.L. Walsworth, *PRB* **82**, 201201(R) (2010).
- H. Seo, A.L. Falk, P.V. Klimov, K.C. Miao, G. Galli, and D.D. Awschalom, *Nat. Comm.*, to be published (2016).
- W. Yang and R. B. Liu, *PRB* **78**, 085315 (2008); W. M. Witczak and S. Das Sarma, *PRB* **74**, 035322 (2006).
- J.R. Maze et al., *PRB* (2008); N. Zhao et al., *PRB* (2012); L.T. Hall et al., *PRB* (2014).
- N. Mizouchi et al., *Phys. Rev. B* **80**, 041201 (2009).
- H. Seo, M. Govoni, and G. Galli, *Sci. Rep.* **6**, 20803 (2016).
- J. B. Varley, A. Janotti, and C.G. Van de Walle, *Phys. Rev. B* **93**, 161201(R) (2016).
- J. H. Skone, M. Govoni, and G. Galli, *Phys. Rev. B* **93**, 195112 (2016).
- A. Alkauskas, P. Broqvist, and A. Pasquarolo, *Phys. Stat. Sol.* **B248**, 775 (2011).
- M. Govoni and G. Galli, *J. Chem. Theory Comp.* **11**, 2680 (2015).
- C. Freyoldt, J. Neugebauer, and C.G. Van de Walle, *Phys. Rev. Lett.* **102**, 016402 (2009).

### Design of new defect qubits in aluminum nitride<sup>15</sup>

- We found that negatively charged N-vacancy ( $V_N^-$ ) have <sup>3</sup>A<sup>"</sup> and <sup>3</sup>A<sub>2</sub> spin-triplet ground states under uniaxial and biaxial strain, respectively.
- The ground states of Al-related defects may not be suitable as spin qubits as they undergo strong resonance with the valence states of AlN.



### Acknowledgements

- The computing resources were provided by:
- NATIONAL SCIENCE FOUNDATION MRSEC
  - U.S. DEPARTMENT OF ENERGY
  - Office of Science
  - AFOSR AIR FORCE OFFICE OF SCIENTIFIC RESEARCH
  - SCIENCE CHIEF OF THE UNITED STATES AIR FORCE
  - NERSC National Energy Research Scientific Computing Center
  - Argonne NATIONAL LABORATORY