

Predator-Prey Dynamics

For this lab, we will be analyzing different models of predator prey dynamics to determine the effects of different assumptions on long-term behavior. Complete the following problems and address any included questions. For your submission, include any requested figures, answers to all questions, and your MATLAB code.

Problem 1: Simulation of Lotka-Volterra Model

The simplest possible model of predator-prey interactions is the Lotka-Volterra model, in which the abundance of predator, P , and prey, H , are given by the ordinary differential equations:

$$\begin{aligned} 1. \quad \frac{dH}{dt} &= aH - bHP \\ 2. \quad \frac{dP}{dt} &= cHP - mP \end{aligned}$$

Where a is the prey reproduction time, b is the prey death due to predation, c is the predator reproduction based on consumption of prey, and m is predator death. Note that here predator survival is dependent on the abundance of prey. For the following problems, use $a = 0.5, b = 3, c = 3, m = 0.1$. For all parameters and species time is in units of days^{-1} and species abundances are in units of cells/mL .

- Write a function in MATLAB to be passed into an ODE solver that given time t , a vector of abundances $y = [H, P]$, and a vector of parameters $p = [a, b, c, d]$ outputs the rates of change $dydt = [\frac{dH}{dt}, \frac{dP}{dt}]$ based on the Lotka-Volterra model. Calculate the rates of change given $t = 0$, $y = [0.4, 0.4]$, and the parameters specified above.
- Using the function that you wrote in part a) and MATLAB's ode solver ode15s (or ode45), simulate the model given initial conditions $[H_0, P_0] = [0.4, 0.4], [0.3, 0.3], [0.2, 0.2]$ over a time period of 500 days. Plot H and P v. t for your first initial conditions of $[H_0, P_0] = [0.4, 0.4]$ on a semiology scale and also P v. H for all your initial conditions on a single plot on a log-log scale. On your second plot, include the nullclines for the differential equations ($\frac{dH}{dt} = 0, \frac{dP}{dt} = 0$) using MATLAB's fimplicit function. You should see that your system never intersects the steady state (intersection of the nullclines). What does this mean?
- How do the initial conditions affect the long term behavior of predator and prey abundances? Is this realistic? Explain. Hint: Realistically there should be other limiting factors to species abundances than the presence of predators or prey, i.e., a carrying capacity.

Problem 2: Introducing a carrying capacity on prey growth.

We will now more realistically model growth of prey abundance with logistic growth, i.e. add a carrying capacity to this term. This system is now modeled by the ordinary differential equations:

$$\begin{aligned} 1. \quad \frac{dH}{dt} &= aH \left(1 - \frac{H}{K}\right) - bHP \\ 2. \quad \frac{dP}{dt} &= cHP - mP \end{aligned}$$

Where K is the carrying capacity of the prey population. For this problem, use $a = 0.5, b = 3, c = 3, m = 0.1, K = 0.4$.

- Write a function in MATLAB to be passed into an ODE solver that given time t , a vector of abundances $y = [H, P]$, and a vector of parameters $p = [a, b, c, d, K]$ outputs the rates of change

$dydt = [\frac{dH}{dt}, \frac{dP}{dt}]$ based on the modified model. Calculate the rates of change given $t = 0$, $y = [0.4, 0.4]$, and the parameters specified above.

- b) Using the function that you wrote in part a) and MATLAB's ode solver ode15s (or ode45), simulate the model given initial conditions $[H_0, P_0] =$ over a time period of 500 days. Plot H and P v. t for your first initial conditions of $[H_0, P_0] = [0.4, 0.4]$, $[0.3, 0.3]$, $[0.2, 0.2]$ on a semiology scale and also P v. H for all your initial conditions on a single plot on a log-log scale. On your second plot, include the nullclines for the differential equations ($\frac{dH}{dt} = 0, \frac{dP}{dt} = 0$) using MATLAB's fimplicit function. How do these simulations compare to those in Problem 1? Do they seem more realistic? In what ways?

Problem 3: Simulating a Macarthur-Rozenzweig model.

Now we will make one more modification to our model to make it more realistic. Here, predator consumption of prey is modified to include an efficiency ϵ and a timescale τ . Our differential equations now take the form:

$$\begin{aligned} 1. \quad \frac{dH}{dt} &= aH \left(1 - \frac{H}{K}\right) - \frac{bH}{1+b\tau H} P \\ 2. \quad \frac{dP}{dt} &= \frac{\epsilon bH}{1+b\tau H} P - mP \end{aligned}$$

For this problem, use $a = 0.5, b = 3, \epsilon = 1, \tau = 3, m = 0.1, K = 0.4$

- a) Write a function in MATLAB to be passed into an ODE solver that given time t , a vector of abundances $y = [H, P]$, and a vector of parameters $p = [a, b, \epsilon, \tau, d, K]$ outputs the rates of change $dydt = [\frac{dH}{dt}, \frac{dP}{dt}]$ based on the modified model. Calculate the rates of change given $t = 0$, $y = [0.4, 0.4]$, and the parameters specified above.
- b) Using the function that you wrote in part a) and MATLAB's ode solver ode15s (or ode45), simulate the model given initial conditions $[H_0, P_0] = [0.4, 0.4], [0.3, 0.3], [0.2, 0.2]$ over a time period of 500 days Plot H and P v. t for your first initial conditions of $[H_0, P_0] = [0.4, 0.4]$ on a semiology scale and also P v. H for all your initial conditions on a single plot on a log-log scale.. On your second plot, include the nullclines for the differential equations ($\frac{dH}{dt} = 0, \frac{dP}{dt} = 0$) using MATLAB's fimplicit function. Compare these simulations to those from Problems 1 and 2. Do these seem more realistic?