

Math 320: Chapter 1

rctcwyvrn

September 2020

1 Common sets

- Natural numbers \mathbb{N} (Rudin uses \mathbb{J})
 - Set notation $\{1, 2, 3, 4, \dots\}$
 - Closed under addition and multiplication
 - Not closed under subtraction (might get a negative number)
- Integers \mathbb{Z}
 - Set notation $\dots - 3, -2, -1, 0, 1, 2, 3, 4, \dots$
 - Closed under subtraction and addition
 - But not under division
- Rationals \mathbb{Q}
 - Set notation: $\{\frac{m}{n} : m \in \mathbb{Z}, n \in \mathbb{N}\}$
 - * Where $\frac{m_1}{n_1} = \frac{m_2}{n_2}$ iff $m_1 * n_2 = m_2 * n_1$
 - * Our intuitive idea of division
 - Alternative we could define \mathbb{Q} in a more straightforward way
 - * Define \mathbb{Q} as the set of ordered pairs $\{(m, n) : m \in \mathbb{Z}, n \in \mathbb{N}\}$
 - * We also need to make sure the equivalent fractions are accounted for
 - * where (m_1, n_1) is equivalent to (m_2, n_2) (ie $(m_1, n_1) \sim (m_2, n_2)$) if $m_1 n_2 = m_2 n_1$
 - Closed under addition, subtraction, multiplication, and division (if divisor is non-zero)
 - Question: Are the rationals sufficient for everything we want to do in real analysis (in calculus)?
 - * Can we do things in calculus?
 - * Can we take limits? (and have them work properly?)
 - Nope! (note: the name of this course is **real** analysis, not **rational** analysis)

- The problem is that the rationals have holes, they're not "filled in all the way" like the reals are

Example: Holes in the rationals

(Rudin 1.1a) We want to show that there's a number we can't reach in the rationals ($\sqrt{2}$). $\nexists p \in \mathbb{Q} : p^2 = 2$

Proof:

By contradiction: Suppose that $\exists p \in \mathbb{Q} : p^2 = 2$. Then since p is rational, we can write $p = \frac{m}{n}$, $m \in \mathbb{Z}$, $n \in \mathbb{N}$.

- WLOG we may suppose that m and n are not both even (because if not then we could just divide both by 2 and have a new m and n and repeat until this statement is true)

- Q: How can we be sure that this dividing by 2 will eventually end? (More explicitly, how many times do we need to do it?)

- We have $2 = p^2 = \frac{m^2}{n^2}$

$$2 = \frac{m^2}{n^2} \tag{1}$$

$$m^2 = 2n^2 \tag{2}$$

- So m must be even
- Ie $m = 2k$ for some odd integer k

$$2n^2 = m^2 \tag{3}$$

$$2n^2 = (2k)^2 \tag{4}$$

$$n^2 = 2k^2 \tag{5}$$

- So n must be even, but we said that one of them had to be odd.
- Contradiction!

Example:

(Rudin 1.1b)

- Let $A = \{p \in \mathbb{Q} : p > 0, p^2 < 2\}$
- Let $B = \{p \in \mathbb{Q} : p > 0, p^2 > 2\}$
- Consider the area where the two sets meet. We know that they meet at $\sqrt{2}$, which is not a rational.
- So we know that the set of rationals has little holes in it, like $\sqrt{2}$ which make us unable to use limits, which are important for real analysis

Then:

- $\forall p \in A, \exists q \in A : p < q$ (A does not have a largest element)
- Similarly B does not have a smallest element ($\forall p \in B, \exists q \in B : q < p$)

Proof:

First one

- Consider arbitrary $p \in A$. Try $q = \frac{2p+2}{2+p}$
- Check $q \in \mathbb{Q}$, the denominator is non-zero and rational, so the result is rational.
- Check $q > 0$, both numerator and denominator are positive
- Check $q^2 < 2$.

$$q^2 = \frac{(2p+2)^2}{(2+p)^2} \quad (6)$$

$$= \frac{2(p^2 - 2)}{(p+2)^2} + 2 \quad (7)$$

- The fraction is less than zero, because the numerator is negative ($p^2 < 2$ because $p \in A$), so $q^2 < 2$
- Check that $p < q$. Well $q = p + \frac{2-p^2}{2+p}$, which is positive, so $q > p$

Second one

- Exercise (Try the exact same choice of q)

Q: Where did that q come from? (Think calculus)

Q: Why do we care so much that there is $q^2 = 2$, but don't care that there isn't a $q^2 = -1$?