

Retiring the third law of black hole thermodynamics

Ryan Unger

Princeton University & University of Cambridge

Oxbridge PDE conference
March 2023

joint work with Christoph Kehle (ETH Zürich)

Classical thermodynamics

Law	Thermodynamics

Classical thermodynamics

Law	Thermodynamics
Zeroth	T constant throughout body in thermal equilibrium

Classical thermodynamics

Law	Thermodynamics
Zeroth	T constant throughout body in thermal equilibrium
First	$dE = TdS + \text{work terms } (PdV)$

Classical thermodynamics

Law	Thermodynamics
Zeroth	T constant throughout body in thermal equilibrium
First	$dE = TdS + \text{work terms } (PdV)$
Second	$dS \geq 0$ in any process

Classical thermodynamics

Law	Thermodynamics
Zeroth	T constant throughout body in thermal equilibrium
First	$dE = TdS + \text{work terms } (PdV)$
Second	$dS \geq 0$ in any process
Third	Impossible to achieve $T = 0$ in finite time

Black hole thermodynamics

Commun. math. Phys. 31, 161–170 (1973)
© by Springer-Verlag 1973

The Four Laws of Black Hole Mechanics

J. M. Bardeen*

Department of Physics, Yale University, New Haven, Connecticut, USA

B. Carter and S. W. Hawking

Institute of Astronomy, University of Cambridge, England

Received January 24, 1973

Black hole thermodynamics is a proposed close mathematical **analogy** between black hole dynamics and classical thermodynamics.

Black hole thermodynamics

Commun. math. Phys. 31, 161–170 (1973)
© by Springer-Verlag 1973

The Four Laws of Black Hole Mechanics

J. M. Bardeen*

Department of Physics, Yale University, New Haven, Connecticut, USA

B. Carter and S. W. Hawking

Institute of Astronomy, University of Cambridge, England

Received January 24, 1973

Black hole thermodynamics is a proposed close mathematical **analogy** between black hole dynamics and classical thermodynamics.

- ▶ What is a black hole?
- ▶ What is the analogue of temperature T ?
- ▶ What is the analogue of entropy S ?

Spacetimes and Einstein's equations

The setting is Einstein's theory of general relativity. A **spacetime** consists of a 4-manifold \mathcal{M}^{3+1} and a Lorentzian metric g satisfying the **Einstein field equations**

$$\operatorname{Ric}(g) - \frac{1}{2}R(g)g = 2T. \quad (\text{EFE})$$

Spacetimes and Einstein's equations

The setting is Einstein's theory of general relativity. A **spacetime** consists of a 4-manifold \mathcal{M}^{3+1} and a Lorentzian metric g satisfying the **Einstein field equations**

$$\text{Ric}(g) - \frac{1}{2}R(g)g = 2T. \quad (\text{EFE})$$

Example: Minkowski space. $\mathcal{M}^{3+1} = \mathbb{R}_{t,x,y,z}^{3+1}$ and

$$g = -dt^2 + dx^2 + dy^2 + dz^2$$

This metric describes the geometry of special relativity with speed of light $c = 1$.

Spacetimes and Einstein's equations

The setting is Einstein's theory of general relativity. A **spacetime** consists of a 4-manifold \mathcal{M}^{3+1} and a Lorentzian metric g satisfying the **Einstein field equations**

$$\text{Ric}(g) - \frac{1}{2}R(g)g = 2T. \quad (\text{EFE})$$

Example: Minkowski space. $\mathcal{M}^{3+1} = \mathbb{R}_{t,x,y,z}^{3+1}$ and

$$g = -dt^2 + dx^2 + dy^2 + dz^2$$

This metric describes the geometry of special relativity with speed of light $c = 1$.

Given a vector $v \in T_p\mathcal{M}$:

- ▶ $g(v, v) < 0$, v is *timelike*
- ▶ $g(v, v) = 0$, v is *null*
- ▶ $g(v, v) > 0$, v is *spacelike*

Curves with null or timelike tangent vector let us define **causality** on a spacetime.

The Reissner–Nordström black hole spacetime

- Mass $M > 0$ and charge $0 \leq |e| \leq M$

$$g_{M,e} = - \left(1 - \frac{2M}{r} + \frac{e^2}{r^2} \right) dt^2 + \left(1 - \frac{2M}{r} + \frac{e^2}{r^2} \right)^{-1} dr^2 + r^2 (d\vartheta^2 + \sin^2 \vartheta d\varphi^2)$$


- Solves the **Einstein-Maxwell** equations
- Describes a static, charged, nonrotating black hole

The Reissner–Nordström black hole spacetime

- Mass $M > 0$ and charge $0 \leq |e| \leq M$

$$g_{M,e} = - \left(1 - \frac{2M}{r} + \frac{e^2}{r^2} \right) dt^2 + \left(1 - \frac{2M}{r} + \frac{e^2}{r^2} \right)^{-1} dr^2 + r^2 (d\vartheta^2 + \sin^2 \vartheta d\varphi^2)$$

- Solves the **Einstein-Maxwell** equations
- Describes a static, charged, nonrotating black hole




Subextremal $|e| < M$

The Reissner–Nordström black hole spacetime


- Mass $M > 0$ and charge $0 \leq |e| \leq M$

$$g_{M,e} = - \left(1 - \frac{2M}{r} + \frac{e^2}{r^2} \right) dt^2 + \left(1 - \frac{2M}{r} + \frac{e^2}{r^2} \right)^{-1} dr^2 + r^2 (d\vartheta^2 + \sin^2 \vartheta d\varphi^2)$$

- Solves the **Einstein-Maxwell** equations
- Describes a static, charged, nonrotating black hole



Subextremal $|e| < M$




Extremal $|e| = M$

The Reissner–Nordström black hole spacetime


- Mass $M > 0$ and charge $0 \leq |e| \leq M$

$$g_{M,e} = - \left(1 - \frac{2M}{r} + \frac{e^2}{r^2} \right) dt^2 + \left(1 - \frac{2M}{r} + \frac{e^2}{r^2} \right)^{-1} dr^2 + r^2 (d\vartheta^2 + \sin^2 \vartheta d\varphi^2)$$

- Solves the **Einstein-Maxwell** equations
- Describes a static, charged, nonrotating black hole



Subextremal $|e| < M$



Extremal $|e| = M$

Superextremal $|e| > M$ does not contain a black hole!

Surface gravity and extremality

- Associated to the horizon is a function $\kappa \geq 0$, called the **surface gravity**

Surface gravity and extremality

- ▶ Associated to the horizon is a function $\kappa \geq 0$, called the **surface gravity**
- ▶ Physically, κ measures the force required to hold a test mass “at the horizon” “from infinity”

Surface gravity and extremality

- ▶ Associated to the horizon is a function $\kappa \geq 0$, called the **surface gravity**
- ▶ Physically, κ measures the force required to hold a test mass “at the horizon” “from infinity”
- ▶ It is also related to the celebrated **(local) redshift effect**

Surface gravity and extremality

- ▶ Associated to the horizon is a function $\kappa \geq 0$, called the **surface gravity**
- ▶ Physically, κ measures the force required to hold a test mass “at the horizon” “from infinity”
- ▶ It is also related to the celebrated **(local) redshift effect**
- ▶ For Reissner–Nordström,

$$\kappa(g_{M,e}) = \frac{\sqrt{M^2 - e^2}}{(M + \sqrt{M^2 - e^2})^2}$$

Surface gravity and extremality

- ▶ Associated to the horizon is a function $\kappa \geq 0$, called the **surface gravity**
- ▶ Physically, κ measures the force required to hold a test mass “at the horizon” “from infinity”
- ▶ It is also related to the celebrated **(local) redshift effect**
- ▶ For Reissner–Nordström,

$$\kappa(g_{M,e}) = \frac{\sqrt{M^2 - e^2}}{(M + \sqrt{M^2 - e^2})^2}$$

- ▶ **Subextremal horizon:** $\kappa > 0$
- ▶ **Extremal horizon:** $\kappa \equiv 0$

Black hole thermodynamics

Law	Classical thermodynamics	Black holes

Black hole thermodynamics

Law	Classical thermodynamics	Black holes
Zeroth	T constant in equilibrium	

Black hole thermodynamics

Law	Classical thermodynamics	Black holes
Zeroth	T constant in equilibrium	surface gravity κ constant on stationary horizon

Black hole thermodynamics

Law	Classical thermodynamics	Black holes
Zeroth	T constant in equilibrium	surface gravity κ constant on stationary horizon
First	$dE = TdS + \dots$	

Black hole thermodynamics

Law	Classical thermodynamics	Black holes
Zeroth	T constant in equilibrium	surface gravity κ constant on stationary horizon
First	$dE = TdS + \dots$	$dM = \kappa dA + \dots$

Black hole thermodynamics

Law	Classical thermodynamics	Black holes
Zeroth	T constant in equilibrium	surface gravity κ constant on stationary horizon
First	$dE = TdS + \dots$	$dM = \kappa dA + \dots$
Second	$dS \geq 0$	

Black hole thermodynamics

Law	Classical thermodynamics	Black holes
Zeroth	T constant in equilibrium	surface gravity κ constant on stationary horizon
First	$dE = TdS + \dots$	$dM = \kappa dA + \dots$
Second	$dS \geq 0$	$dA \geq 0$

Black hole thermodynamics

Law	Classical thermodynamics	Black holes
Zeroth	T constant in equilibrium	surface gravity κ constant on stationary horizon
First	$dE = TdS + \dots$	$dM = \kappa dA + \dots$
Second	$dS \geq 0$	$dA \geq 0$
Third	$T \not\rightarrow 0$ in finite process	

Black hole thermodynamics

Law	Classical thermodynamics	Black holes
Zeroth	T constant in equilibrium	surface gravity κ constant on stationary horizon
First	$dE = TdS + \dots$	$dM = \kappa dA + \dots$
Second	$dS \geq 0$	$dA \geq 0$
Third	$T \not\rightarrow 0$ in finite process	surface gravity $\kappa \not\rightarrow 0$ in finite advanced time

Black hole thermodynamics

Law	Classical thermodynamics	Black holes
Zeroth	T constant in equilibrium	surface gravity κ constant on stationary horizon
First	$dE = TdS + \dots$	$dM = \kappa dA + \dots$
Second	$dS \geq 0$	$dA \geq 0$
Third	$T \not\rightarrow 0$ in finite process	surface gravity $\kappa \not\rightarrow 0$ in finite advanced time

- Proposed by Bardeen–Carter–Hawking '73

Black hole thermodynamics

Law	Classical thermodynamics	Black holes
Zeroth	T constant in equilibrium	surface gravity κ constant on stationary horizon
First	$dE = TdS + \dots$	$dM = \kappa dA + \dots$
Second	$dS \geq 0$	$dA \geq 0$
Third	$T \not\rightarrow 0$ in finite process	surface gravity $\kappa \not\rightarrow 0$ in finite advanced time

- Proposed by Bardeen–Carter–Hawking '73
- Hawking radiation—interpret $\kappa \propto T$

Black hole thermodynamics

Law	Classical thermodynamics	Black holes
Zeroth	T constant in equilibrium	surface gravity κ constant on stationary horizon
First	$dE = TdS + \dots$	$dM = \kappa dA + \dots$
Second	$dS \geq 0$	$dA \geq 0$
Third	$T \not\rightarrow 0$ in finite process	surface gravity $\kappa \not\rightarrow 0$ in finite advanced time

- ▶ Proposed by Bardeen–Carter–Hawking '73
- ▶ Hawking radiation—interpret $\kappa \propto T$
- ▶ Bekenstein entropy—interpret $A \propto S$

Black hole thermodynamics

Law	Classical thermodynamics	Black holes
Zeroth	T constant in equilibrium	surface gravity κ constant on stationary horizon
First	$dE = TdS + \dots$	$dM = \kappa dA + \dots$
Second	$dS \geq 0$	$dA \geq 0$
Third	$T \not\rightarrow 0$ in finite process	surface gravity $\kappa \not\rightarrow 0$ in finite advanced time

- ▶ Proposed by Bardeen–Carter–Hawking '73
- ▶ Hawking radiation—interpret $\kappa \propto T$
- ▶ Bekenstein entropy—interpret $A \propto S$
- ▶ These identifications are further “justified” once quantum effects are considered, but are **formal analogies** in classical GR.

Black hole thermodynamics

Law	Classical thermodynamics	Black holes
Zeroth	T constant in equilibrium	surface gravity κ constant on stationary horizon
First	$dE = TdS + \dots$	$dM = \kappa dA + \dots$
Second	$dS \geq 0$	$dA \geq 0$
Third	$T \not\rightarrow 0$ in finite process	surface gravity $\kappa \not\rightarrow 0$ in finite advanced time

- ▶ Proposed by Bardeen–Carter–Hawking '73
- ▶ Hawking radiation—interpret $\kappa \propto T$
- ▶ Bekenstein entropy—interpret $A \propto S$
- ▶ These identifications are further “justified” once quantum effects are considered, but are **formal analogies** in classical GR.
- ▶ Laws 0, 1, and 2 proved by Hawking, Carter, Bardeen–Carter–Hawking, Wald, ...

Retiring the third law

Original formulation of Bardeen–Carter–Hawking:

Extending the analogy even further one would postulate:

The Third Law

It is impossible by any procedure, no matter how idealized, to reduce κ to zero by a finite sequence of operations.

Retiring the third law

Original formulation of Bardeen–Carter–Hawking:

Extending the analogy even further one would postulate:

The Third Law

It is impossible by any procedure, no matter how idealized, to reduce κ to zero by a finite sequence of operations.

Subextremal: $\kappa > 0 \iff T > 0$

Extremal: $\kappa = 0 \iff T = 0$

Retiring the third law

Conjecture (The third law, BCH '73, Israel '86).

*A subextremal black hole cannot become extremal in finite time by any continuous process, no matter how idealized, in which the spacetime and matter fields remain *regular* and obey the *weak energy condition*.*

Retiring the third law

Conjecture (The third law, BCH '73, Israel '86).


A subextremal black hole cannot become extremal in finite time by any continuous process, no matter how idealized, in which the spacetime and matter fields remain regular and obey the weak energy condition.

Theorem (Kehle–U. '22).

Subextremal black holes can become extremal in finite time, evolving from regular initial data. In particular, the “third law of black hole thermodynamics” is false.


Counterexample to the third law: detailed statement of our theorem

There exist regular solutions of the Einstein–Maxwell–charged scalar field system with the following behavior:



Counterexample to the third law: detailed statement of our theorem


There exist regular solutions of the Einstein–Maxwell–charged scalar field system with the following behavior:



- Spherically symmetric Cauchy data for the Einstein–Maxwell–charged scalar field system which undergo gravitational collapse.

Counterexample to the third law: detailed statement of our theorem


There exist regular solutions of the Einstein–Maxwell–charged scalar field system with the following behavior:



- ▶ Spherically symmetric Cauchy data for the Einstein–Maxwell–charged scalar field system which undergo gravitational collapse.
- ▶ Forms an exactly Schwarzschild “apparent horizon.”

Counterexample to the third law: detailed statement of our theorem


There exist regular solutions of the Einstein–Maxwell–charged scalar field system with the following behavior:



- Spherically symmetric Cauchy data for the Einstein–Maxwell–charged scalar field system which undergo gravitational collapse.
- Forms an exactly Schwarzschild “apparent horizon.”
- Forms an exactly extremal Reissner–Nordström event horizon at a later advanced time.

Counterexample to the third law: detailed statement of our theorem


There exist regular solutions of the Einstein–Maxwell–charged scalar field system with the following behavior:



- ▶ Spherically symmetric Cauchy data for the Einstein–Maxwell–charged scalar field system which undergo gravitational collapse.
- ▶ Forms an exactly Schwarzschild “apparent horizon.”
- ▶ Forms an exactly extremal Reissner–Nordström event horizon at a later advanced time.
- ▶ These solutions are arbitrarily regular: for any $k \in \mathbb{N}$, there exists a C^k example.

Counterexample to the third law: detailed statement of our theorem

There exist regular solutions of the Einstein–Maxwell–charged scalar field system with the following behavior:



- ▶ Spherically symmetric Cauchy data for the Einstein–Maxwell–charged scalar field system which undergo gravitational collapse.
- ▶ Forms an exactly Schwarzschild “apparent horizon.”
- ▶ Forms an exactly extremal Reissner–Nordström event horizon at a later advanced time.
- ▶ These solutions are arbitrarily regular: for any $k \in \mathbb{N}$, there exists a C^k example.
- ▶ Model satisfies the dominant energy condition.

Einstein–Maxwell-charged scalar field system

- ▶ Lorentzian manifold (\mathcal{M}^{3+1}, g)
- ▶ 2-form $F = dA$ (electromagnetism)
- ▶ Charged (complex) scalar field ϕ

Einstein–Maxwell-charged scalar field system

- Lorentzian manifold (\mathcal{M}^{3+1}, g)
- 2-form $F = dA$ (electromagnetism)
- Charged (complex) scalar field ϕ
- Equations:

$$R_{\mu\nu}(g) - \frac{1}{2}R(g)g_{\mu\nu} = 2 \left(T_{\mu\nu}^{\text{EM}} + T_{\mu\nu}^{\text{CSF}} \right) \quad (1)$$

$$\nabla^\mu F_{\mu\nu} = 2\epsilon \operatorname{Im}(\phi \overline{D_\nu \phi}) \quad (2)$$

$$g^{\mu\nu} D_\mu D_\nu \phi = 0 \quad (3)$$

$$T_{\mu\nu}^{\text{EM}} = g^{\alpha\beta} F_{\alpha\nu} F_{\beta\mu} - \frac{1}{4} F^{\alpha\beta} F_{\alpha\beta} g_{\mu\nu} \quad (4)$$

$$T_{\mu\nu}^{\text{CSF}} = \operatorname{Re}(D_\mu \phi \overline{D_\nu \phi}) - \frac{1}{2} g_{\mu\nu} g^{\alpha\beta} D_\alpha \phi \overline{D_\beta \phi} \quad (5)$$

Einstein–Maxwell-charged scalar field system

- Lorentzian manifold (\mathcal{M}^{3+1}, g)
- 2-form $F = dA$ (electromagnetism)
- Charged (complex) scalar field ϕ
- Equations:

$$R_{\mu\nu}(g) - \frac{1}{2}R(g)g_{\mu\nu} = 2 \left(T_{\mu\nu}^{\text{EM}} + T_{\mu\nu}^{\text{CSF}} \right) \quad (1)$$

$$\nabla^\mu F_{\mu\nu} = 2\epsilon \operatorname{Im}(\phi \overline{D_\nu \phi}) \quad (2)$$

$$g^{\mu\nu} D_\mu D_\nu \phi = 0 \quad (3)$$

$$T_{\mu\nu}^{\text{EM}} = g^{\alpha\beta} F_{\alpha\nu} F_{\beta\mu} - \frac{1}{4} F^{\alpha\beta} F_{\alpha\beta} g_{\mu\nu} \quad (4)$$

$$T_{\mu\nu}^{\text{CSF}} = \operatorname{Re}(D_\mu \phi \overline{D_\nu \phi}) - \frac{1}{2} g_{\mu\nu} g^{\alpha\beta} D_\alpha \phi \overline{D_\beta \phi} \quad (5)$$

- In an appropriate gauge, this is a nonlinear hyperbolic system for (g, F, ϕ) .

Einstein–Maxwell-charged scalar field system

- Lorentzian manifold (\mathcal{M}^{3+1}, g)
- 2-form $F = dA$ (electromagnetism)
- Charged (complex) scalar field ϕ
- Equations:

$$R_{\mu\nu}(g) - \frac{1}{2}R(g)g_{\mu\nu} = 2 \left(T_{\mu\nu}^{\text{EM}} + T_{\mu\nu}^{\text{CSF}} \right) \quad (1)$$

$$\nabla^\mu F_{\mu\nu} = 2\epsilon \operatorname{Im}(\phi \overline{D_\nu \phi}) \quad (2)$$

$$g^{\mu\nu} D_\mu D_\nu \phi = 0 \quad (3)$$

$$T_{\mu\nu}^{\text{EM}} = g^{\alpha\beta} F_{\alpha\nu} F_{\beta\mu} - \frac{1}{4} F^{\alpha\beta} F_{\alpha\beta} g_{\mu\nu} \quad (4)$$

$$T_{\mu\nu}^{\text{CSF}} = \operatorname{Re}(D_\mu \phi \overline{D_\nu \phi}) - \frac{1}{2} g_{\mu\nu} g^{\alpha\beta} D_\alpha \phi \overline{D_\beta \phi} \quad (5)$$

- In an appropriate gauge, this is a nonlinear hyperbolic system for (g, F, ϕ) .
- Can be thought of as an **initial value problem**.

Einstein–Maxwell-charged scalar field system

- Lorentzian manifold (\mathcal{M}^{3+1}, g)
- 2-form $F = dA$ (electromagnetism)
- Charged (complex) scalar field ϕ
- Equations:

$$R_{\mu\nu}(g) - \frac{1}{2}R(g)g_{\mu\nu} = 2 \left(T_{\mu\nu}^{\text{EM}} + T_{\mu\nu}^{\text{CSF}} \right) \quad (1)$$

$$\nabla^\mu F_{\mu\nu} = 2\epsilon \operatorname{Im}(\phi \overline{D_\nu \phi}) \quad (2)$$

$$g^{\mu\nu} D_\mu D_\nu \phi = 0 \quad (3)$$

$$T_{\mu\nu}^{\text{EM}} = g^{\alpha\beta} F_{\alpha\nu} F_{\beta\mu} - \frac{1}{4} F^{\alpha\beta} F_{\alpha\beta} g_{\mu\nu} \quad (4)$$

$$T_{\mu\nu}^{\text{CSF}} = \operatorname{Re}(D_\mu \phi \overline{D_\nu \phi}) - \frac{1}{2} g_{\mu\nu} g^{\alpha\beta} D_\alpha \phi \overline{D_\beta \phi} \quad (5)$$

- In an appropriate gauge, this is a nonlinear hyperbolic system for (g, F, ϕ) .
- Can be thought of as an **initial value problem**.
- Spherical symmetry!

Einstein–Maxwell-charged scalar field system

- Lorentzian manifold (\mathcal{M}^{3+1}, g)
- 2-form $F = dA$ (electromagnetism)
- Charged (complex) scalar field ϕ
- Equations:

$$R_{\mu\nu}(g) - \frac{1}{2}R(g)g_{\mu\nu} = 2 \left(T_{\mu\nu}^{\text{EM}} + T_{\mu\nu}^{\text{CSF}} \right) \quad (1)$$

$$\nabla^\mu F_{\mu\nu} = 2\epsilon \operatorname{Im}(\phi \overline{D_\nu \phi}) \quad (2)$$


$$g^{\mu\nu} D_\mu D_\nu \phi = 0 \quad (3)$$

$$T_{\mu\nu}^{\text{EM}} = g^{\alpha\beta} F_{\alpha\nu} F_{\beta\mu} - \frac{1}{4} F^{\alpha\beta} F_{\alpha\beta} g_{\mu\nu} \quad (4)$$


$$T_{\mu\nu}^{\text{CSF}} = \operatorname{Re}(D_\mu \phi \overline{D_\nu \phi}) - \frac{1}{2} g_{\mu\nu} g^{\alpha\beta} D_\alpha \phi \overline{D_\beta \phi} \quad (5)$$

- In an appropriate gauge, this is a nonlinear hyperbolic system for (g, F, ϕ) .
- Can be thought of as an **initial value problem**.
- Spherical symmetry!
- Scalar field ϕ degree of freedom required to break the rigidity of pure Einstein–Maxwell in spherical symmetry.


Prototype: Minkowski to Schwarzschild gluing



Prototype: Minkowski to Schwarzschild gluing




Prototype: Minkowski to Schwarzschild gluing



Characteristic gluing for the vacuum Einstein equations recently initiated in fundamental works of Aretakis–Czimek–Rodnianski and refined by Czimek–Rodnianski, but in a **perturbative** regime around Minkowski space which is inapplicable here.

Prototype: Minkowski to Schwarzschild gluing



Characteristic gluing for the vacuum Einstein equations recently initiated in fundamental works of Aretakis–Czimek–Rodnianski and refined by Czimek–Rodnianski, but in a **perturbative** regime around Minkowski space which is inapplicable here.

The present problem is fundamentally **nonlinear** and **nonperturbative**, but is tractable because of the symmetry assumption.

Toy model: gluing for the linear wave equation

We now fix a spacetime (\mathcal{M}^{3+1}, g) and consider the **linear wave equation**

$$\square_g \phi = 0 \tag{*}$$

Toy model: gluing for the linear wave equation

We now fix a spacetime (\mathcal{M}^{3+1}, g) and consider the **linear wave equation**

$$\square_g \phi = 0 \tag{*}$$

Question.

Given two domains $\mathfrak{R}_1, \mathfrak{R}_2 \subset \mathcal{M}$ and functions

$$\phi_1 : \mathfrak{R}_1 \rightarrow \mathbb{R}$$

$$\phi_2 : \mathfrak{R}_2 \rightarrow \mathbb{R}$$

solving (*)

Toy model: gluing for the linear wave equation

We now fix a spacetime (\mathcal{M}^{3+1}, g) and consider the **linear wave equation**

$$\square_g \phi = 0 \tag{*}$$

Question.

Given two domains $\mathfrak{R}_1, \mathfrak{R}_2 \subset \mathcal{M}$ and functions

$$\phi_1 : \mathfrak{R}_1 \rightarrow \mathbb{R}$$

$$\phi_2 : \mathfrak{R}_2 \rightarrow \mathbb{R}$$

solving $(*)$, when does there exist a function


$$\phi : \mathcal{M} \rightarrow \mathbb{R}$$

solving $(*)$, such that


$$\phi|_{\mathfrak{R}_1} = \phi_1$$

$$\phi|_{\mathfrak{R}_2} = \phi_2 ?$$

Characteristic/null gluing for the linear wave equation




Characteristic/null gluing for the linear wave equation



Analogy:

- ▶ \mathfrak{R}_1 is Minkowski
- ▶ \mathfrak{R}_2 is Schwarzschild
- ▶ C will be the event horizon \mathcal{H}^+


Characteristic/null gluing for the linear wave equation



Analogy:

- ▶ \mathfrak{R}_1 is Minkowski
- ▶ \mathfrak{R}_2 is Schwarzschild
- ▶ C will be the event horizon \mathcal{H}^+

Characteristic/null gluing for the linear wave equation



Analogy:

- ▶ \mathfrak{R}_1 is Minkowski
- ▶ \mathfrak{R}_2 is Schwarzschild
- ▶ C will be the event horizon \mathcal{H}^+

The characteristic initial value problem


Bifurcate null hypersurface: $C \cup \underline{C}$ where C and \underline{C} meet transversally, $C \cap \underline{C} \approx S^2$

Proposition (Characteristic IVP for the wave equation).

Given a bifurcate null hypersurface $C \cup \underline{C}$ and $\phi_0 : C \cup \underline{C} \rightarrow \mathbb{R}$, there exists a unique function ϕ defined to the future of $C \cup \underline{C}$ such that

$$\square_g \phi = 0 \quad (*)$$

and $\phi|_{C \cup \underline{C}} = \phi_0$.



The characteristic initial value problem


Bifurcate null hypersurface: $C \cup \underline{C}$ where C and \underline{C} meet transversally, $C \cap \underline{C} \approx S^2$

Proposition (Characteristic IVP for the wave equation).

Given a bifurcate null hypersurface $C \cup \underline{C}$ and $\phi_0 : C \cup \underline{C} \rightarrow \mathbb{R}$, there exists a unique function ϕ defined to the future of $C \cup \underline{C}$ such that

$$\square_g \phi = 0 \tag{*}$$

and $\phi|_{C \cup \underline{C}} = \phi_0$.



The characteristic initial value problem


Bifurcate null hypersurface: $C \cup \underline{C}$ where C and \underline{C} meet transversally, $C \cap \underline{C} \approx S^2$

Proposition (Characteristic IVP for the wave equation).

Given a bifurcate null hypersurface $C \cup \underline{C}$ and $\phi_0 : C \cup \underline{C} \rightarrow \mathbb{R}$, there exists a unique function ϕ defined to the future of $C \cup \underline{C}$ such that

$$\square_g \phi = 0 \tag{*}$$

and $\phi|_{C \cup \underline{C}} = \phi_0$.



Characteristic data for the wave equation does not involve derivatives.

The null “constraints”

In any well posed initial value problem, there needs to be a mechanism to compute the full solution on the initial data hypersurface.

The null “constraints”

In any well posed initial value problem, there needs to be a mechanism to compute the full solution on the initial data hypersurface. Toy problem for the toy problem:

- ▶ Wave equation on Minkowski \mathbb{R}^{3+1}
- ▶ $u = t - r$, $v = t + r$ double null coordinates

The null “constraints”

In any well posed initial value problem, there needs to be a mechanism to compute the full solution on the initial data hypersurface. Toy problem for the toy problem:

- ▶ Wave equation on Minkowski \mathbb{R}^{3+1}
- ▶ $u = t - r$, $v = t + r$ double null coordinates
- ▶ $\phi = \phi(t, r, \vartheta, \varphi) = \phi(u, v, \vartheta, \varphi)$ satisfies

$$\square\phi = -\partial_t^2\phi + r^{-2}\partial_r(r^2\partial_r\phi) + \Delta\phi = 0$$

$$\partial_u\partial_v(r\phi) = \Delta(r\phi) \quad (*)$$

The null “constraints”

In any well posed initial value problem, there needs to be a mechanism to compute the full solution on the initial data hypersurface. Toy problem for the toy problem:

- ▶ Wave equation on Minkowski \mathbb{R}^{3+1}
- ▶ $u = t - r$, $v = t + r$ double null coordinates
- ▶ $\phi = \phi(t, r, \vartheta, \varphi) = \phi(u, v, \vartheta, \varphi)$ satisfies

$$\square\phi = -\partial_t^2\phi + r^{-2}\partial_r(r^2\partial_r\phi) + \Delta\phi = 0$$

$$\partial_u\partial_v(r\phi) = \Delta(r\phi) \quad (*)$$

- ▶ Integrate $(*)$ on C :

$$\partial_u(r\phi)|_{v_2} - \partial_u(r\phi)|_{v_1} = \int_{v_1}^{v_2} \Delta(r\phi) dv'$$

The null “constraints”

In any well posed initial value problem, there needs to be a mechanism to compute the full solution on the initial data hypersurface. Toy problem for the toy problem:

- ▶ Wave equation on Minkowski \mathbb{R}^{3+1}
- ▶ $u = t - r$, $v = t + r$ double null coordinates
- ▶ $\phi = \phi(t, r, \vartheta, \varphi) = \phi(u, v, \vartheta, \varphi)$ satisfies

$$\square\phi = -\partial_t^2\phi + r^{-2}\partial_r(r^2\partial_r\phi) + \Delta\phi = 0$$

$$\partial_u\partial_v(r\phi) = \Delta(r\phi) \quad (*)$$

- ▶ Integrate $(*)$ on C :

$$\partial_u(r\phi)|_{v_2} - \partial_u(r\phi)|_{v_1} = \int_{v_1}^{v_2} \Delta(r\phi) dv'$$

- ▶ $\implies \partial_u\phi|_C$ can be calculated from (1) $\partial_u\phi$ at the bifurcation sphere and (2) $\phi|_C$

The null “constraints”

In any well posed initial value problem, there needs to be a mechanism to compute the full solution on the initial data hypersurface. Toy problem for the toy problem:

- ▶ Wave equation on Minkowski \mathbb{R}^{3+1}
- ▶ $u = t - r$, $v = t + r$ double null coordinates
- ▶ $\phi = \phi(t, r, \vartheta, \varphi) = \phi(u, v, \vartheta, \varphi)$ satisfies

$$\square\phi = -\partial_t^2\phi + r^{-2}\partial_r(r^2\partial_r\phi) + \Delta\phi = 0$$

$$\partial_u\partial_v(r\phi) = \Delta(r\phi) \quad (*)$$

- ▶ Integrate $(*)$ on C :

$$\partial_u(r\phi)|_{v_2} - \partial_u(r\phi)|_{v_1} = \int_{v_1}^{v_2} \Delta(r\phi) dv'$$

- ▶ $\implies \partial_u\phi|_C$ can be calculated from (1) $\partial_u\phi$ at the bifurcation sphere and (2) $\phi|_C$
- ▶ $(*)$ lets us compute $\partial_u^i\phi$ on C to all orders

The null “constraints”

In general, the wave equation can be interpreted as a transport equation for the transverse derivative of ϕ , sourced by ϕ and tangential derivatives (which are a part of the characteristic data)

The null “constraints”

In general, the wave equation can be interpreted as a transport equation for the transverse derivative of ϕ , sourced by ϕ and tangential derivatives (which are a part of the characteristic data)

On C , the wave equation is equivalent to

$$(\partial_v + \text{known function})\partial_u\phi = \text{known function},$$

where “known function” depends on underlying geometry and data.

The null “constraints”

In general, the wave equation can be interpreted as a transport equation for the transverse derivative of ϕ , sourced by ϕ and tangential derivatives (which are a part of the characteristic data)


On C , the wave equation is equivalent to

$$(\partial_v + \text{known function})\partial_u\phi = \text{known function},$$

where “known function” depends on underlying geometry and data.


Commuting the wave equation with transverse derivatives gives an inductive system of transport equations for all orders!

Back to characteristic gluing



The C^k characteristic gluing problem reduces to:


Back to characteristic gluing



The C^k characteristic gluing problem reduces to:

- ▶ Given two spheres S_1 and S_2 along an outgoing null cone C
- ▶ Given k ingoing and outgoing derivatives of ϕ at S_1 and S_2


Back to characteristic gluing



The C^k characteristic gluing problem reduces to:

- ▶ Given two spheres S_1 and S_2 along an outgoing null cone C
- ▶ Given k ingoing and outgoing derivatives of ϕ at S_1 and S_2
- ▶ Prescribe ϕ along the part of C between S_1 and S_2 so that the outgoing derivatives agree with the given ones and the solutions of the transport equations for the ingoing derivatives have the specified initial and final values.


Back to characteristic gluing




The C^k characteristic gluing problem reduces to:

- ▶ Given two spheres S_1 and S_2 along an outgoing null cone C
- ▶ Given k ingoing and outgoing derivatives of ϕ at S_1 and S_2
- ▶ Prescribe ϕ along the part of C between S_1 and S_2 so that the outgoing derivatives agree with the given ones and the solutions of the transport equations for the ingoing derivatives have the specified initial and final values.
- ▶ Initial/boundary value problem for a system of coupled ODEs where you are allowed to (almost) freely prescribe the potential


Minkowski to Schwarzschild gluing



Minkowski to Schwarzschild gluing



Minkowski to Schwarzschild gluing



Einstein-scalar field in spherical symmetry

► $\mathcal{M}^{3+1} = \mathcal{Q}^{1+1} \times S^2$

$$g = -\Omega^2 du dv + r^2 g_{S^2}$$

► $\Omega(u, v) > 0$ lapse, $r(u, v) > 0$ area-radius

Einstein-scalar field in spherical symmetry

► $\mathcal{M}^{3+1} = \mathcal{Q}^{1+1} \times S^2$

$$g = -\Omega^2 du dv + r^2 g_{S^2}$$

► $\Omega(u, v) > 0$ lapse, $r(u, v) > 0$ area-radius

► Wave equations

$$\partial_u \partial_v \phi = -\frac{\partial_u \phi \partial_v r}{r} - \frac{\partial_u r \partial_v \phi}{r} \quad (6)$$

$$\partial_u \partial_v r = -\frac{\Omega^2}{4r} - \frac{\partial_u r \partial_v r}{r} \quad (7)$$

$$\partial_u \partial_v \log(\Omega^2) = \frac{\Omega^2}{2r^2} + 2 \frac{\partial_u r \partial_v r}{r^2} \quad (8)$$

Einstein-scalar field in spherical symmetry

► $\mathcal{M}^{3+1} = \mathcal{Q}^{1+1} \times S^2$

$$g = -\Omega^2 du dv + r^2 g_{S^2}$$

► $\Omega(u, v) > 0$ lapse, $r(u, v) > 0$ area-radius

► Wave equations

$$\partial_u \partial_v \phi = -\frac{\partial_u \phi \partial_v r}{r} - \frac{\partial_u r \partial_v \phi}{r} \quad (6)$$

$$\partial_u \partial_v r = -\frac{\Omega^2}{4r} - \frac{\partial_u r \partial_v r}{r} \quad (7)$$

$$\partial_u \partial_v \log(\Omega^2) = \frac{\Omega^2}{2r^2} + 2 \frac{\partial_u r \partial_v r}{r^2} \quad (8)$$

► Raychaudhuri's equations (constraints)


$$\partial_u \left(\frac{\partial_u r}{\Omega^2} \right) = -\frac{r}{\Omega^2} (\partial_u \phi)^2 \quad (9)$$

$$\partial_v \left(\frac{\partial_v r}{\Omega^2} \right) = -\frac{r}{\Omega^2} (\partial_v \phi)^2 \quad (10)$$

Minkowski to Schwarzschild gluing

Theorem (Kehle–U. '22).

For any $k \in \mathbb{N}$ and $0 < R_i < 2M_f$, the Minkowski sphere of radius R_i can be characteristically glued to the Schwarzschild event horizon sphere with mass M_f to order C^k within the Einstein-scalar field model in spherical symmetry.



Idea of the proof

- ▶ Null constraint system becomes a coupled nonlinear system of ODEs sourced by $\phi(v)$

Idea of the proof

- ▶ Null constraint system becomes a coupled nonlinear system of ODEs sourced by $\phi(v)$
- ▶ Scalar field ansatz

$$\phi_\alpha(v) = \sum_{1 \leq j \leq k+1} \alpha_j \chi_j(v), \quad \alpha \in \mathbb{R}^{k+1}$$

Idea of the proof

- ▶ Null constraint system becomes a coupled nonlinear system of ODEs sourced by $\phi(v)$
- ▶ Scalar field ansatz

$$\phi_\alpha(v) = \sum_{1 \leq j \leq k+1} \alpha_j \chi_j(v), \quad \alpha \in \mathbb{R}^{k+1}$$

- ▶ Initialize $\partial_u \phi_\alpha(0) = \dots = \partial_u^k \phi_\alpha(0) = 0$

Idea of the proof

- ▶ Null constraint system becomes a coupled nonlinear system of ODEs sourced by $\phi(v)$
- ▶ Scalar field ansatz

$$\phi_\alpha(v) = \sum_{1 \leq j \leq k+1} \alpha_j \chi_j(v), \quad \alpha \in \mathbb{R}^{k+1}$$

- ▶ Initialize $\partial_u \phi_\alpha(0) = \dots = \partial_u^k \phi_\alpha(0) = 0$
- ▶ Initialize $r(v)$ teleologically at $v = 1$ ($\partial_v r(0)$ is a gauge choice, but $\partial_v r(1) = 0!$)

Idea of the proof

- ▶ Null constraint system becomes a coupled nonlinear system of ODEs sourced by $\phi(v)$
- ▶ Scalar field ansatz

$$\phi_\alpha(v) = \sum_{1 \leq j \leq k+1} \alpha_j \chi_j(v), \quad \alpha \in \mathbb{R}^{k+1}$$

- ▶ Initialize $\partial_u \phi_\alpha(0) = \dots = \partial_u^k \phi_\alpha(0) = 0$
- ▶ Initialize $r(v)$ teleologically at $v = 1$ ($\partial_v r(0)$ is a gauge choice, but $\partial_v r(1) = 0!$)
- ▶ For each α , generate $r(v)$, $\partial_u r(v)$, $\partial_u \phi_\alpha(v)$, etc.

Idea of the proof

- ▶ Null constraint system becomes a coupled nonlinear system of ODEs sourced by $\phi(v)$
- ▶ Scalar field ansatz

$$\phi_\alpha(v) = \sum_{1 \leq j \leq k+1} \alpha_j \chi_j(v), \quad \alpha \in \mathbb{R}^{k+1}$$

- ▶ Initialize $\partial_u \phi_\alpha(0) = \dots = \partial_u^k \phi_\alpha(0) = 0$
- ▶ Initialize $r(v)$ teleologically at $v = 1$ ($\partial_v r(0)$ is a gauge choice, but $\partial_v r(1) = 0!$)
- ▶ For each α , generate $r(v)$, $\partial_u r(v)$, $\partial_u \phi_\alpha(v)$, etc.
- ▶ The antipodal map $\alpha \mapsto -\alpha$ leaves geometric quantities fixed (r , Ω^2 , and derivatives)

Idea of the proof

- ▶ Null constraint system becomes a coupled nonlinear system of ODEs sourced by $\phi(v)$
- ▶ Scalar field ansatz

$$\phi_\alpha(v) = \sum_{1 \leq j \leq k+1} \alpha_j \chi_j(v), \quad \alpha \in \mathbb{R}^{k+1}$$

- ▶ Initialize $\partial_u \phi_\alpha(0) = \dots = \partial_u^k \phi_\alpha(0) = 0$
- ▶ Initialize $r(v)$ teleologically at $v = 1$ ($\partial_v r(0)$ is a gauge choice, but $\partial_v r(1) = 0!$)
- ▶ For each α , generate $r(v)$, $\partial_u r(v)$, $\partial_u \phi_\alpha(v)$, etc.
- ▶ The antipodal map $\alpha \mapsto -\alpha$ leaves geometric quantities fixed (r , Ω^2 , and derivatives)
- ▶ The antipodal map $\alpha \mapsto -\alpha$ changes the sign of the scalar field and its derivatives

Idea of the proof

- ▶ Null constraint system becomes a coupled nonlinear system of ODEs sourced by $\phi(v)$
- ▶ Scalar field ansatz

$$\phi_\alpha(v) = \sum_{1 \leq j \leq k+1} \alpha_j \chi_j(v), \quad \alpha \in \mathbb{R}^{k+1}$$

- ▶ Initialize $\partial_u \phi_\alpha(0) = \dots = \partial_u^k \phi_\alpha(0) = 0$
- ▶ Initialize $r(v)$ teleologically at $v = 1$ ($\partial_v r(0)$ is a gauge choice, but $\partial_v r(1) = 0!$)
- ▶ For each α , generate $r(v)$, $\partial_u r(v)$, $\partial_u \phi_\alpha(v)$, etc.
- ▶ The antipodal map $\alpha \mapsto -\alpha$ leaves geometric quantities fixed (r , Ω^2 , and derivatives)
- ▶ The antipodal map $\alpha \mapsto -\alpha$ changes the sign of the scalar field and its derivatives
- ▶ In particular, the map

$$\alpha \mapsto \left(\partial_u \phi_\alpha(1), \dots, \partial_u^k \phi_\alpha(1) \right)$$

Idea of the proof

- ▶ Null constraint system becomes a coupled nonlinear system of ODEs sourced by $\phi(v)$
- ▶ Scalar field ansatz

$$\phi_\alpha(v) = \sum_{1 \leq j \leq k+1} \alpha_j \chi_j(v), \quad \alpha \in \mathbb{R}^{k+1}$$

- ▶ Initialize $\partial_u \phi_\alpha(0) = \dots = \partial_u^k \phi_\alpha(0) = 0$
- ▶ Initialize $r(v)$ teleologically at $v = 1$ ($\partial_v r(0)$ is a gauge choice, but $\partial_v r(1) = 0!$)
- ▶ For each α , generate $r(v)$, $\partial_u r(v)$, $\partial_u \phi_\alpha(v)$, etc.
- ▶ The antipodal map $\alpha \mapsto -\alpha$ leaves geometric quantities fixed (r , Ω^2 , and derivatives)
- ▶ The antipodal map $\alpha \mapsto -\alpha$ changes the sign of the scalar field and its derivatives
- ▶ In particular, the map


$$\alpha \mapsto \left(\partial_u \phi_\alpha(1), \dots, \partial_u^k \phi_\alpha(1) \right)$$

is odd

- ▶ **Borsuk–Ulam theorem:** there exists α_* such that

$$\left(\partial_u \phi_{\alpha_*}(1), \dots, \partial_u^k \phi_{\alpha_*}(1) \right) = 0.$$

Disproof of the third law



The vacuum case

We believe that the charged scalar field is not necessary to provide a counterexample to the third law:

The vacuum case

We believe that the charged scalar field is not necessary to provide a counterexample to the third law:

Conjecture.

There exist Cauchy data for the Einstein vacuum equations

$$R_{\mu\nu} = 0$$

which undergo gravitational collapse and form an exactly Schwarzschild apparent horizon, only for the spacetime to form an exactly extremal Kerr event horizon at a later advanced time. In particular, already in vacuum, the “third law of black hole thermodynamics” is false.

The vacuum case

We believe that the charged scalar field is not necessary to provide a counterexample to the third law:

Conjecture.

There exist Cauchy data for the Einstein vacuum equations

$$R_{\mu\nu} = 0$$

which undergo gravitational collapse and form an exactly Schwarzschild apparent horizon, only for the spacetime to form an exactly extremal Kerr event horizon at a later advanced time. In particular, already in vacuum, the “third law of black hole thermodynamics” is false.

This would follow from characteristic gluing, in vacuum, of a Minkowski sphere to a Kerr exterior sphere.

Conclusions

Conclusions

- ▶ Subextremal black holes can become extremal in finite time, so the third law of black hole thermodynamics is false.

Conclusions


- ▶ Subextremal black holes can become extremal in finite time, so the third law of black hole thermodynamics is false.
- ▶ Our nonperturbative characteristic gluing scheme opens up a new avenue for constructing solutions of the Einstein equation with explicit behavior.

Conclusions

- ▶ Subextremal black holes can become extremal in finite time, so the third law of black hole thermodynamics is false.
- ▶ Our nonperturbative characteristic gluing scheme opens up a new avenue for constructing solutions of the Einstein equation with explicit behavior.
- ▶ Understanding dynamics *near* these new spacetimes is a rich source of problems going forward.

Conclusions

- ▶ Subextremal black holes can become extremal in finite time, so the third law of black hole thermodynamics is false.
- ▶ Our nonperturbative characteristic gluing scheme opens up a new avenue for constructing solutions of the Einstein equation with explicit behavior.
- ▶ Understanding dynamics *near* these new spacetimes is a rich source of problems going forward.



Thank you!