

# Retiring the third law of black hole thermodynamics

Ryan Unger

Department of Mathematics, Princeton University

Vanderbilt Analysis Seminar  
February 2023

joint work with Christoph Kehle (ETH Zürich)  
[arXiv:2211.15742](https://arxiv.org/abs/2211.15742)

# Classical thermodynamics

Law	Thermodynamics

# Classical thermodynamics

Law	Thermodynamics
Zeroth	$T$ constant throughout body in thermal equilibrium

# Classical thermodynamics

Law	Thermodynamics
Zeroth	$T$ constant throughout body in thermal equilibrium
First	$dE = TdS + \text{work terms } (PdV)$

# Classical thermodynamics

Law	Thermodynamics
Zeroth	$T$ constant throughout body in thermal equilibrium
First	$dE = TdS + \text{work terms } (PdV)$
Second	$dS \geq 0$ in any process

# Classical thermodynamics

Law	Thermodynamics
Zeroth	$T$ constant throughout body in thermal equilibrium
First	$dE = TdS + \text{work terms } (PdV)$
Second	$dS \geq 0$ in any process
Third	Impossible to achieve $T = 0$ in finite time

# Black hole thermodynamics

Commun. math. Phys. 31, 161–170 (1973)  
© by Springer-Verlag 1973

## The Four Laws of Black Hole Mechanics

J. M. Bardeen\*

Department of Physics, Yale University, New Haven, Connecticut, USA

B. Carter and S. W. Hawking

Institute of Astronomy, University of Cambridge, England

Received January 24, 1973

**Black hole thermodynamics** is a proposed close mathematical **analogy** between black hole dynamics and classical thermodynamics.

# Black hole thermodynamics

Commun. math. Phys. 31, 161–170 (1973)  
© by Springer-Verlag 1973

## The Four Laws of Black Hole Mechanics

J. M. Bardeen\*

Department of Physics, Yale University, New Haven, Connecticut, USA

B. Carter and S. W. Hawking

Institute of Astronomy, University of Cambridge, England

Received January 24, 1973

**Black hole thermodynamics** is a proposed close mathematical **analogy** between black hole dynamics and classical thermodynamics.

- ▶ What is a black hole?
- ▶ What is the analogue of temperature  $T$ ?
- ▶ What is the analogue of entropy  $S$ ?



# Crash course on black holes: Spacetimes and Einstein's equations

The setting is Einstein's theory of general relativity. A **spacetime** consists of a 4-manifold  $\mathcal{M}^{3+1}$  and a Lorentzian metric  $g$  satisfying the **Einstein field equations**

$$\text{Ric}(g) - \frac{1}{2}R(g)g = 2T. \quad (\text{EFE})$$

# Crash course on black holes: Spacetimes and Einstein's equations

The setting is Einstein's theory of general relativity. A **spacetime** consists of a 4-manifold  $\mathcal{M}^{3+1}$  and a Lorentzian metric  $g$  satisfying the **Einstein field equations**

$$\text{Ric}(g) - \frac{1}{2}R(g)g = 2T. \quad (\text{EFE})$$

**Example: Minkowski space.**  $\mathcal{M}^{3+1} = \mathbb{R}_{t,x,y,z}^{3+1}$  and

$$g = -dt^2 + dx^2 + dy^2 + dz^2$$

This metric describes the geometry of special relativity with speed of light  $c = 1$ .

# Crash course on black holes: Spacetimes and Einstein's equations

The setting is Einstein's theory of general relativity. A **spacetime** consists of a 4-manifold  $\mathcal{M}^{3+1}$  and a Lorentzian metric  $g$  satisfying the **Einstein field equations**

$$\text{Ric}(g) - \frac{1}{2}R(g)g = 2T. \quad (\text{EFE})$$

**Example: Minkowski space.**  $\mathcal{M}^{3+1} = \mathbb{R}_{t,x,y,z}^{3+1}$  and

$$g = -dt^2 + dx^2 + dy^2 + dz^2$$


This metric describes the geometry of special relativity with speed of light  $c = 1$ .

Given a vector  $v \in T_p\mathcal{M}$ :

- ▶  $g(v, v) < 0$ ,  $v$  is *timelike*
- ▶  $g(v, v) = 0$ ,  $v$  is *null*
- ▶  $g(v, v) > 0$ ,  $v$  is *spacelike*

Curves with null or timelike tangent vector let us define **causality** on a spacetime.

# Crash course on black holes: Geometry of Minkowski space



## Crash course on black holes: What is a black hole?

**Asymptotic flatness:**  $g$  approaches the Minkowski metric “at infinity.”

## Crash course on black holes: What is a black hole?

**Asymptotic flatness:**  $g$  approaches the Minkowski metric “at infinity.”

An asymptotically flat spacetime is said to contain a **black hole region**

$$\mathcal{BH} \subset \mathcal{M}$$

if observers and signals originating in  $\mathcal{BH}$  cannot reach infinity.

## Crash course on black holes: What is a black hole?

**Asymptotic flatness:**  $g$  approaches the Minkowski metric “at infinity.”

An asymptotically flat spacetime is said to contain a **black hole region**

$$\mathcal{BH} \subset \mathcal{M}$$

if observers and signals originating in  $\mathcal{BH}$  cannot reach infinity.

The **event horizon** of the black hole is its topological boundary

$$\mathcal{H}^+ := \partial(\mathcal{BH}).$$

# Crash course on black holes: What is a black hole?

**Asymptotic flatness:**  $g$  approaches the Minkowski metric “at infinity.”

An asymptotically flat spacetime is said to contain a **black hole region**

$$\mathcal{BH} \subset \mathcal{M}$$

if observers and signals originating in  $\mathcal{BH}$  cannot reach infinity.

The **event horizon** of the black hole is its topological boundary

$$\mathcal{H}^+ := \partial(\mathcal{BH}).$$

The event horizon  $\mathcal{H}^+$  is a **null hypersurface**: it is ruled by geodesics with null tangent vectors.



# Crash course on black holes: What is a black hole?

**Asymptotic flatness:**  $g$  approaches the Minkowski metric “at infinity.”

An asymptotically flat spacetime is said to contain a **black hole region**

$$\mathcal{BH} \subset \mathcal{M}$$

if observers and signals originating in  $\mathcal{BH}$  cannot reach infinity.

The **event horizon** of the black hole is its topological boundary

$$\mathcal{H}^+ := \partial(\mathcal{BH}).$$

The event horizon  $\mathcal{H}^+$  is a **null hypersurface**: it is ruled by geodesics with null tangent vectors.

Examples to come shortly!

## Crash course on black holes: Teleology



The black hole region  $\mathcal{BH}$  and event horizon  $\mathcal{H}^+$  of a spacetime are **teleological** notions—they can't be known to exist, much less be located, *without knowing the entire future of the spacetime*.


## Crash course on black holes: Teleology



The black hole region  $\mathcal{BH}$  and event horizon  $\mathcal{H}^+$  of a spacetime are **teleological** notions—they can't be known to exist, much less be located, *without knowing the entire future of the spacetime*.

The event horizon  $\mathcal{H}^+$  is not a singular object.

## Crash course on black holes: Teleology



The black hole region  $\mathcal{BH}$  and event horizon  $\mathcal{H}^+$  of a spacetime are **teleological** notions—they can't be known to exist, much less be located, *without knowing the entire future of the spacetime*.

The event horizon  $\mathcal{H}^+$  is not a singular object.

The formation, dynamics, and stability of black holes can be rigorously studied using the tools of Lorentzian geometry and hyperbolic PDE.


## Crash course on black holes: Penrose diagrams

**Penrose diagrams:** A  $(1 + 1)$ -dimensional diagrammatic representation of spacetimes. This is particularly useful for visualizing causality.

# Crash course on black holes: Penrose diagrams


**Penrose diagrams:** A  $(1 + 1)$ -dimensional diagrammatic representation of spacetimes. This is particularly useful for visualizing causality.

- ▶ 45 degree lines correspond to null hypersurfaces in  $(3 + 1)$  dimensions
- ▶ steeper than 45: timelike
- ▶ shallower than 45: spacelike




Penrose diagram of Minkowski space

## Crash course on black holes: Minkowski space




## Crash course on black holes: Minkowski space






## Crash course on black holes: Minkowski space



## Crash course on black holes: Minkowski space



## Crash course on black holes: The Schwarzschild solution

- Mass  $M > 0$

$$g_M = - \left(1 - \frac{2M}{r}\right) dt^2 + \left(1 - \frac{2M}{r}\right)^{-1} dr^2 + r^2(d\theta^2 + \sin^2 \theta d\phi^2)$$


- Solves the **Einstein vacuum** equations
- Describes a static, nonrotating black hole

## Crash course on black holes: The Schwarzschild solution

- Mass  $M > 0$

$$g_M = - \left(1 - \frac{2M}{r}\right) dt^2 + \left(1 - \frac{2M}{r}\right)^{-1} dr^2 + r^2(d\theta^2 + \sin^2 \theta d\phi^2)$$

- Solves the **Einstein vacuum** equations
- Describes a static, nonrotating black hole




## Crash course on black holes: The Schwarzschild solution

- Mass  $M > 0$

$$g_M = - \left(1 - \frac{2M}{r}\right) dt^2 + \left(1 - \frac{2M}{r}\right)^{-1} dr^2 + r^2(d\theta^2 + \sin^2 \theta d\phi^2)$$

- Solves the **Einstein vacuum** equations
- Describes a static, nonrotating black hole




As  $M \rightarrow 0$ , recover the Minkowski metric.

## Crash course on black holes: The Schwarzschild solution

- Mass  $M > 0$

$$g_M = - \left(1 - \frac{2M}{r}\right) dt^2 + \left(1 - \frac{2M}{r}\right)^{-1} dr^2 + r^2(d\theta^2 + \sin^2 \theta d\phi^2)$$

- Solves the **Einstein vacuum** equations
- Describes a static, nonrotating black hole



As  $M \rightarrow 0$ , recover the Minkowski metric.

## Crash course on black holes: Reissner–Nordström

- Mass  $M > 0$  and charge  $0 \leq |e| \leq M$

$$g_{M,e} = - \left( 1 - \frac{2M}{r} + \frac{e^2}{r^2} \right) dt^2 + \left( 1 - \frac{2M}{r} + \frac{e^2}{r^2} \right)^{-1} dr^2 + r^2 (d\theta^2 + \sin^2 \theta d\phi^2)$$


- Solves the **Einstein-Maxwell** equations
- Describes a static, charged, nonrotating black hole

## Crash course on black holes: Reissner–Nordström

- Mass  $M > 0$  and charge  $0 \leq |e| \leq M$

$$g_{M,e} = - \left( 1 - \frac{2M}{r} + \frac{e^2}{r^2} \right) dt^2 + \left( 1 - \frac{2M}{r} + \frac{e^2}{r^2} \right)^{-1} dr^2 + r^2 (d\theta^2 + \sin^2 \theta d\phi^2)$$

- Solves the **Einstein-Maxwell** equations
- Describes a static, charged, nonrotating black hole



**Subextremal**  $|e| < M$




# Crash course on black holes: Reissner–Nordström


- Mass  $M > 0$  and charge  $0 \leq |e| \leq M$

$$g_{M,e} = - \left( 1 - \frac{2M}{r} + \frac{e^2}{r^2} \right) dt^2 + \left( 1 - \frac{2M}{r} + \frac{e^2}{r^2} \right)^{-1} dr^2 + r^2 (d\theta^2 + \sin^2 \theta d\phi^2)$$

- Solves the **Einstein-Maxwell** equations
- Describes a static, charged, nonrotating black hole



**Subextremal**  $|e| < M$




**Extremal**  $|e| = M$

# Crash course on black holes: Reissner–Nordström


- Mass  $M > 0$  and charge  $0 \leq |e| \leq M$

$$g_{M,e} = - \left( 1 - \frac{2M}{r} + \frac{e^2}{r^2} \right) dt^2 + \left( 1 - \frac{2M}{r} + \frac{e^2}{r^2} \right)^{-1} dr^2 + r^2 (d\theta^2 + \sin^2 \theta d\phi^2)$$

- Solves the **Einstein-Maxwell** equations
- Describes a static, charged, nonrotating black hole



**Subextremal**  $|e| < M$



**Extremal**  $|e| = M$

**Superextremal**  $|e| > M$  does not contain a black hole!

## Crash course on black holes: Surface gravity and extremality

- The event horizons in these exact solutions are **Killing horizons**: The spacetime admits a Killing field which is null and tangent to the horizon

## Crash course on black holes: Surface gravity and extremality

- ▶ The event horizons in these exact solutions are **Killing horizons**: The spacetime admits a Killing field which is null and tangent to the horizon
- ▶ Associated to a Killing horizon is a function  $\kappa \geq 0$ , called the **surface gravity**

## Crash course on black holes: Surface gravity and extremality

- ▶ The event horizons in these exact solutions are **Killing horizons**: The spacetime admits a Killing field which is null and tangent to the horizon
- ▶ Associated to a Killing horizon is a function  $\kappa \geq 0$ , called the **surface gravity**
- ▶ Physically,  $\kappa$  measures the force required to hold a test mass “at the horizon” “from infinity”

## Crash course on black holes: Surface gravity and extremality

- ▶ The event horizons in these exact solutions are **Killing horizons**: The spacetime admits a Killing field which is null and tangent to the horizon
- ▶ Associated to a Killing horizon is a function  $\kappa \geq 0$ , called the **surface gravity**
- ▶ Physically,  $\kappa$  measures the force required to hold a test mass “at the horizon” “from infinity”
- ▶ It is also related to the celebrated **(local) redshift effect**

## Crash course on black holes: Surface gravity and extremality

- ▶ The event horizons in these exact solutions are **Killing horizons**: The spacetime admits a Killing field which is null and tangent to the horizon
- ▶ Associated to a Killing horizon is a function  $\kappa \geq 0$ , called the **surface gravity**
- ▶ Physically,  $\kappa$  measures the force required to hold a test mass “at the horizon” “from infinity”
- ▶ It is also related to the celebrated **(local) redshift effect**
- ▶ For Reissner–Nordström,

$$\kappa(g_{M,e}) = \frac{\sqrt{M^2 - e^2}}{(M + \sqrt{M^2 - e^2})^2}$$

## Crash course on black holes: Surface gravity and extremality

- ▶ The event horizons in these exact solutions are **Killing horizons**: The spacetime admits a Killing field which is null and tangent to the horizon
- ▶ Associated to a Killing horizon is a function  $\kappa \geq 0$ , called the **surface gravity**
- ▶ Physically,  $\kappa$  measures the force required to hold a test mass “at the horizon” “from infinity”
- ▶ It is also related to the celebrated **(local) redshift effect**
- ▶ For Reissner–Nordström,

$$\kappa(g_{M,e}) = \frac{\sqrt{M^2 - e^2}}{(M + \sqrt{M^2 - e^2})^2}$$

- ▶ **Subextremal Killing horizon**:  $\kappa > 0$
- ▶ **Extremal Killing horizon**:  $\kappa \equiv 0$



# Black hole thermodynamics

Law	Classical thermodynamics	Black holes

# Black hole thermodynamics

Law	Classical thermodynamics	Black holes
Zeroth	$T$ constant in equilibrium	

# Black hole thermodynamics

Law	Classical thermodynamics	Black holes
Zeroth	$T$ constant in equilibrium	surface gravity $\kappa$ constant on stationary horizon

# Black hole thermodynamics

Law	Classical thermodynamics	Black holes
Zeroth	$T$ constant in equilibrium	surface gravity $\kappa$ constant on stationary horizon
First	$dE = TdS + \dots$	

# Black hole thermodynamics

Law	Classical thermodynamics	Black holes
Zeroth	$T$ constant in equilibrium	surface gravity $\kappa$ constant on stationary horizon
First	$dE = TdS + \dots$	$dM = \kappa dA + \dots$

# Black hole thermodynamics

Law	Classical thermodynamics	Black holes
Zeroth	$T$ constant in equilibrium	surface gravity $\kappa$ constant on stationary horizon
First	$dE = TdS + \dots$	$dM = \kappa dA + \dots$
Second	$dS \geq 0$	

# Black hole thermodynamics

Law	Classical thermodynamics	Black holes
Zeroth	$T$ constant in equilibrium	surface gravity $\kappa$ constant on stationary horizon
First	$dE = TdS + \dots$	$dM = \kappa dA + \dots$
Second	$dS \geq 0$	$dA \geq 0$

# Black hole thermodynamics

Law	Classical thermodynamics	Black holes
Zeroth	$T$ constant in equilibrium	surface gravity $\kappa$ constant on stationary horizon
First	$dE = TdS + \dots$	$dM = \kappa dA + \dots$
Second	$dS \geq 0$	$dA \geq 0$
Third	$T \not\rightarrow 0$ in finite process	



# Black hole thermodynamics

Law	Classical thermodynamics	Black holes
Zeroth	$T$ constant in equilibrium	surface gravity $\kappa$ constant on stationary horizon
First	$dE = TdS + \dots$	$dM = \kappa dA + \dots$
Second	$dS \geq 0$	$dA \geq 0$
Third	$T \not\rightarrow 0$ in finite process	surface gravity $\kappa \not\rightarrow 0$ in finite advanced time

# Black hole thermodynamics

Law	Classical thermodynamics	Black holes
Zeroth	$T$ constant in equilibrium	surface gravity $\kappa$ constant on stationary horizon
First	$dE = TdS + \dots$	$dM = \kappa dA + \dots$
Second	$dS \geq 0$	$dA \geq 0$
Third	$T \not\rightarrow 0$ in finite process	surface gravity $\kappa \not\rightarrow 0$ in finite advanced time

- Proposed by Bardeen–Carter–Hawking '73

# Black hole thermodynamics

Law	Classical thermodynamics	Black holes
Zeroth	$T$ constant in equilibrium	surface gravity $\kappa$ constant on stationary horizon
First	$dE = TdS + \dots$	$dM = \kappa dA + \dots$
Second	$dS \geq 0$	$dA \geq 0$
Third	$T \not\rightarrow 0$ in finite process	surface gravity $\kappa \not\rightarrow 0$ in finite advanced time

- Proposed by Bardeen–Carter–Hawking '73
- Hawking radiation—interpret  $\kappa \propto T$

# Black hole thermodynamics

Law	Classical thermodynamics	Black holes
Zeroth	$T$ constant in equilibrium	surface gravity $\kappa$ constant on stationary horizon
First	$dE = TdS + \dots$	$dM = \kappa dA + \dots$
Second	$dS \geq 0$	$dA \geq 0$
Third	$T \not\rightarrow 0$ in finite process	surface gravity $\kappa \not\rightarrow 0$ in finite advanced time

- ▶ Proposed by Bardeen–Carter–Hawking '73
- ▶ Hawking radiation—interpret  $\kappa \propto T$
- ▶ Bekenstein entropy—interpret  $A \propto S$

# Black hole thermodynamics

Law	Classical thermodynamics	Black holes
Zeroth	$T$ constant in equilibrium	surface gravity $\kappa$ constant on stationary horizon
First	$dE = TdS + \dots$	$dM = \kappa dA + \dots$
Second	$dS \geq 0$	$dA \geq 0$
Third	$T \not\rightarrow 0$ in finite process	surface gravity $\kappa \not\rightarrow 0$ in finite advanced time

- ▶ Proposed by Bardeen–Carter–Hawking '73
- ▶ Hawking radiation—interpret  $\kappa \propto T$
- ▶ Bekenstein entropy—interpret  $A \propto S$
- ▶ These identifications are further justified once quantum effects are considered, but are **formal analogies** in classical GR.

# Black hole thermodynamics

Law	Classical thermodynamics	Black holes
Zeroth	$T$ constant in equilibrium	surface gravity $\kappa$ constant on stationary horizon
First	$dE = TdS + \dots$	$dM = \kappa dA + \dots$
Second	$dS \geq 0$	$dA \geq 0$
Third	$T \not\rightarrow 0$ in finite process	surface gravity $\kappa \not\rightarrow 0$ in finite advanced time

- ▶ Proposed by Bardeen–Carter–Hawking '73
- ▶ Hawking radiation—interpret  $\kappa \propto T$
- ▶ Bekenstein entropy—interpret  $A \propto S$
- ▶ These identifications are further justified once quantum effects are considered, but are **formal analogies** in classical GR.
- ▶ Laws 0, 1, and 2 proved by Hawking, Carter, Bardeen–Carter–Hawking, Wald, ...

# Retiring the third law

Original formulation of Bardeen–Carter–Hawking:

Extending the analogy even further one would postulate:

## *The Third Law*

It is impossible by any procedure, no matter how idealized, to reduce  $\kappa$  to zero by a finite sequence of operations.

## Retiring the third law

### Conjecture (The third law, BCH '73, Israel '86).

*A subextremal black hole cannot become extremal in finite time by any continuous process, no matter how idealized, in which the spacetime and matter fields remain regular and obey the weak energy condition.*



# Retiring the third law

## Conjecture (The third law, BCH '73, Israel '86).

*A subextremal black hole cannot become extremal in finite time by any continuous process, no matter how idealized, in which the spacetime and matter fields remain regular and obey the weak energy condition.*

## Theorem (Kehle–U. '22).

*Subextremal black holes can become extremal in finite time, evolving from regular initial data. In particular, the “third law of black hole thermodynamics” is false.*

## Retiring the third law

### Conjecture (The third law, BCH '73, Israel '86).

*A subextremal black hole cannot become extremal in finite time by any continuous process, no matter how idealized, in which the spacetime and matter fields remain regular and obey the weak energy condition.*


### Theorem (Kehle–U. '22).

*Subextremal black holes can become extremal in finite time, evolving from regular initial data. In particular, the “third law of black hole thermodynamics” is false.*

More precisely, there exist regular solutions of the Einstein–Maxwell-charged scalar field system with the following behavior:


## Counterexample to the third law: detailed statement of our theorem

There exist regular solutions of the Einstein–Maxwell–charged scalar field system with the following behavior:



## Counterexample to the third law: detailed statement of our theorem


There exist regular solutions of the Einstein–Maxwell–charged scalar field system with the following behavior:



- Spherically symmetric Cauchy data for the Einstein–Maxwell–charged scalar field system which undergo gravitational collapse.

## Counterexample to the third law: detailed statement of our theorem


There exist regular solutions of the Einstein–Maxwell–charged scalar field system with the following behavior:



- Spherically symmetric Cauchy data for the Einstein–Maxwell–charged scalar field system which undergo gravitational collapse.
- Forms an exactly Schwarzschild “apparent horizon.”

## Counterexample to the third law: detailed statement of our theorem


There exist regular solutions of the Einstein–Maxwell–charged scalar field system with the following behavior:



- Spherically symmetric Cauchy data for the Einstein–Maxwell–charged scalar field system which undergo gravitational collapse.
- Forms an exactly Schwarzschild “apparent horizon.”
- Forms an exactly extremal Reissner–Nordström event horizon at a later advanced time.

## Counterexample to the third law: detailed statement of our theorem


There exist regular solutions of the Einstein–Maxwell–charged scalar field system with the following behavior:



- ▶ Spherically symmetric Cauchy data for the Einstein–Maxwell–charged scalar field system which undergo gravitational collapse.
- ▶ Forms an exactly Schwarzschild “apparent horizon.”
- ▶ Forms an exactly extremal Reissner–Nordström event horizon at a later advanced time.
- ▶ These solutions are arbitrarily regular: for any  $k \in \mathbb{N}$ , there exists a  $C^k$  example.

## Counterexample to the third law: detailed statement of our theorem

There exist regular solutions of the Einstein–Maxwell–charged scalar field system with the following behavior:



- ▶ Spherically symmetric Cauchy data for the Einstein–Maxwell–charged scalar field system which undergo gravitational collapse.
- ▶ Forms an exactly Schwarzschild “apparent horizon.”
- ▶ Forms an exactly extremal Reissner–Nordström event horizon at a later advanced time.
- ▶ These solutions are arbitrarily regular: for any  $k \in \mathbb{N}$ , there exists a  $C^k$  example.
- ▶ Model satisfies the dominant energy condition.



# Einstein–Maxwell-charged scalar field system

- ▶ Lorentzian manifold  $(\mathcal{M}^{3+1}, g)$
- ▶ 2-form  $F = dA$  (electromagnetism)
- ▶ Charged (complex) scalar field  $\phi$

# Einstein–Maxwell-charged scalar field system

- Lorentzian manifold  $(\mathcal{M}^{3+1}, g)$
- 2-form  $F = dA$  (electromagnetism)
- Charged (complex) scalar field  $\phi$
- Equations:

$$R_{\mu\nu}(g) - \frac{1}{2}R(g)g_{\mu\nu} = 2 \left( T_{\mu\nu}^{\text{EM}} + T_{\mu\nu}^{\text{CSF}} \right) \quad (1)$$

$$\nabla^\mu F_{\mu\nu} = 2\mathfrak{e} \operatorname{Im}(\phi \overline{D_\nu \phi}) \quad (2)$$

$$g^{\mu\nu} D_\mu D_\nu \phi = 0 \quad (3)$$

$$T_{\mu\nu}^{\text{EM}} = g^{\alpha\beta} F_{\alpha\nu} F_{\beta\mu} - \frac{1}{4} F^{\alpha\beta} F_{\alpha\beta} g_{\mu\nu} \quad (4)$$

$$T_{\mu\nu}^{\text{CSF}} = \operatorname{Re}(D_\mu \phi \overline{D_\nu \phi}) - \frac{1}{2} g_{\mu\nu} g^{\alpha\beta} D_\alpha \phi \overline{D_\beta \phi} \quad (5)$$

# Einstein–Maxwell-charged scalar field system

- Lorentzian manifold  $(\mathcal{M}^{3+1}, g)$
- 2-form  $F = dA$  (electromagnetism)
- Charged (complex) scalar field  $\phi$
- Equations:

$$R_{\mu\nu}(g) - \frac{1}{2}R(g)g_{\mu\nu} = 2 \left( T_{\mu\nu}^{\text{EM}} + T_{\mu\nu}^{\text{CSF}} \right) \quad (1)$$

$$\nabla^\mu F_{\mu\nu} = 2\mathfrak{e} \operatorname{Im}(\phi \overline{D_\nu \phi}) \quad (2)$$

$$g^{\mu\nu} D_\mu D_\nu \phi = 0 \quad (3)$$

$$T_{\mu\nu}^{\text{EM}} = g^{\alpha\beta} F_{\alpha\nu} F_{\beta\mu} - \frac{1}{4} F^{\alpha\beta} F_{\alpha\beta} g_{\mu\nu} \quad (4)$$

$$T_{\mu\nu}^{\text{CSF}} = \operatorname{Re}(D_\mu \phi \overline{D_\nu \phi}) - \frac{1}{2} g_{\mu\nu} g^{\alpha\beta} D_\alpha \phi \overline{D_\beta \phi} \quad (5)$$

- In an appropriate gauge, this is a nonlinear hyperbolic system for  $(g, F, \phi)$ .

# Einstein–Maxwell-charged scalar field system

- ▶ Lorentzian manifold  $(\mathcal{M}^{3+1}, g)$
- ▶ 2-form  $F = dA$  (electromagnetism)
- ▶ Charged (complex) scalar field  $\phi$
- ▶ Equations:

$$R_{\mu\nu}(g) - \frac{1}{2}R(g)g_{\mu\nu} = 2 \left( T_{\mu\nu}^{\text{EM}} + T_{\mu\nu}^{\text{CSF}} \right) \quad (1)$$

$$\nabla^\mu F_{\mu\nu} = 2\mathfrak{e} \operatorname{Im}(\phi \overline{D_\nu \phi}) \quad (2)$$

$$g^{\mu\nu} D_\mu D_\nu \phi = 0 \quad (3)$$

$$T_{\mu\nu}^{\text{EM}} = g^{\alpha\beta} F_{\alpha\nu} F_{\beta\mu} - \frac{1}{4} F^{\alpha\beta} F_{\alpha\beta} g_{\mu\nu} \quad (4)$$

$$T_{\mu\nu}^{\text{CSF}} = \operatorname{Re}(D_\mu \phi \overline{D_\nu \phi}) - \frac{1}{2} g_{\mu\nu} g^{\alpha\beta} D_\alpha \phi \overline{D_\beta \phi} \quad (5)$$

- ▶ In an appropriate gauge, this is a nonlinear hyperbolic system for  $(g, F, \phi)$ .
- ▶ Can be thought of as an **initial value problem**.

# Einstein–Maxwell-charged scalar field system

- ▶ Lorentzian manifold  $(\mathcal{M}^{3+1}, g)$
- ▶ 2-form  $F = dA$  (electromagnetism)
- ▶ Charged (complex) scalar field  $\phi$
- ▶ Equations:

$$R_{\mu\nu}(g) - \frac{1}{2}R(g)g_{\mu\nu} = 2 \left( T_{\mu\nu}^{\text{EM}} + T_{\mu\nu}^{\text{CSF}} \right) \quad (1)$$

$$\nabla^\mu F_{\mu\nu} = 2\mathfrak{e} \operatorname{Im}(\phi \overline{D_\nu \phi}) \quad (2)$$

$$g^{\mu\nu} D_\mu D_\nu \phi = 0 \quad (3)$$

$$T_{\mu\nu}^{\text{EM}} = g^{\alpha\beta} F_{\alpha\nu} F_{\beta\mu} - \frac{1}{4} F^{\alpha\beta} F_{\alpha\beta} g_{\mu\nu} \quad (4)$$

$$T_{\mu\nu}^{\text{CSF}} = \operatorname{Re}(D_\mu \phi \overline{D_\nu \phi}) - \frac{1}{2} g_{\mu\nu} g^{\alpha\beta} D_\alpha \phi \overline{D_\beta \phi} \quad (5)$$

- ▶ In an appropriate gauge, this is a nonlinear hyperbolic system for  $(g, F, \phi)$ .
- ▶ Can be thought of as an **initial value problem**.
- ▶ Spherical symmetry!

# Einstein–Maxwell-charged scalar field system

- ▶ Lorentzian manifold  $(\mathcal{M}^{3+1}, g)$
- ▶ 2-form  $F = dA$  (electromagnetism)
- ▶ Charged (complex) scalar field  $\phi$
- ▶ Equations:

$$R_{\mu\nu}(g) - \frac{1}{2}R(g)g_{\mu\nu} = 2 \left( T_{\mu\nu}^{\text{EM}} + T_{\mu\nu}^{\text{CSF}} \right) \quad (1)$$

$$\nabla^\mu F_{\mu\nu} = 2\epsilon \operatorname{Im}(\phi \overline{D_\nu \phi}) \quad (2)$$


$$g^{\mu\nu} D_\mu D_\nu \phi = 0 \quad (3)$$

$$T_{\mu\nu}^{\text{EM}} = g^{\alpha\beta} F_{\alpha\nu} F_{\beta\mu} - \frac{1}{4} F^{\alpha\beta} F_{\alpha\beta} g_{\mu\nu} \quad (4)$$


$$T_{\mu\nu}^{\text{CSF}} = \operatorname{Re}(D_\mu \phi \overline{D_\nu \phi}) - \frac{1}{2} g_{\mu\nu} g^{\alpha\beta} D_\alpha \phi \overline{D_\beta \phi} \quad (5)$$

- ▶ In an appropriate gauge, this is a nonlinear hyperbolic system for  $(g, F, \phi)$ .
- ▶ Can be thought of as an **initial value problem**.
- ▶ Spherical symmetry!
- ▶ Scalar field  $\phi$  degree of freedom required to break the rigidity of pure Einstein–Maxwell in spherical symmetry.

## Prototype: Minkowski to Schwarzschild gluing




## Prototype: Minkowski to Schwarzschild gluing






## Prototype: Minkowski to Schwarzschild gluing



**Characteristic gluing** for the vacuum Einstein equations recently initiated in fundamental works of Aretakis–Czimek–Rodnianski and refined by Czimek–Rodnianski, but in a **perturbative** regime around Minkowski space which is inapplicable here.

## Prototype: Minkowski to Schwarzschild gluing



**Characteristic gluing** for the vacuum Einstein equations recently initiated in fundamental works of Aretakis–Czimek–Rodnianski and refined by Czimek–Rodnianski, but in a **perturbative** regime around Minkowski space which is inapplicable here.

The present problem is fundamentally **nonlinear** and **nonperturbative**, but is tractable because of the symmetry assumption.

## Toy model: gluing for the linear wave equation

We now fix a spacetime  $(\mathcal{M}^{3+1}, g)$  and consider the **linear wave equation**

$$\square_g \phi = 0 \tag{*}$$

## Toy model: gluing for the linear wave equation

We now fix a spacetime  $(\mathcal{M}^{3+1}, g)$  and consider the **linear wave equation**

$$\square_g \phi = 0 \tag{*}$$

### Question.

Given two domains  $\mathfrak{R}_1, \mathfrak{R}_2 \subset \mathcal{M}$  and functions

$$\phi_1 : \mathfrak{R}_1 \rightarrow \mathbb{R}$$

$$\phi_2 : \mathfrak{R}_2 \rightarrow \mathbb{R}$$

solving (\*)

## Toy model: gluing for the linear wave equation

We now fix a spacetime  $(\mathcal{M}^{3+1}, g)$  and consider the **linear wave equation**

$$\square_g \phi = 0 \tag{*}$$

### Question.

Given two domains  $\mathfrak{R}_1, \mathfrak{R}_2 \subset \mathcal{M}$  and functions

$$\phi_1 : \mathfrak{R}_1 \rightarrow \mathbb{R}$$

$$\phi_2 : \mathfrak{R}_2 \rightarrow \mathbb{R}$$

solving  $(*)$ , when does there exist a function

$$\phi : \mathcal{M} \rightarrow \mathbb{R}$$


solving  $(*)$ , such that

$$\phi|_{\mathfrak{R}_1} = \phi_1$$

$$\phi|_{\mathfrak{R}_2} = \phi_2 ?$$


# Spacelike gluing for the linear wave equation

Consider first the case of two spacelike separated regions:



# Spacelike gluing for the linear wave equation

Consider first the case of two spacelike separated regions:



In this case the gluing problem for  $(*)$  is trivial:

## **Proposition (Cauchy problem for the wave equation).**


*Given a spacelike hypersurface  $\Sigma \subset (\mathcal{M}, g)$  and functions  $\phi_0, \phi_1 \in C^\infty(\Sigma)$ , there exists a unique function  $\phi : \mathcal{M} \rightarrow \mathbb{R}$  solving*

$$\square_g \phi = 0 \tag{*}$$

*such that  $\phi|_\Sigma = \phi_0$ ,  $n(\phi)|_\Sigma = \phi_1$ .*

# Spacelike gluing for the linear wave equation

Consider first the case of two spacelike separated regions:



In this case the gluing problem for  $(*)$  is trivial:

## **Proposition (Cauchy problem for the wave equation).**

*Given a spacelike hypersurface  $\Sigma \subset (\mathcal{M}, g)$  and functions  $\phi_0, \phi_1 \in C^\infty(\Sigma)$ , there exists a unique function  $\phi : \mathcal{M} \rightarrow \mathbb{R}$  solving*


$$\square_g \phi = 0 \tag{*}$$

*such that  $\phi|_\Sigma = \phi_0$ ,  $n(\phi)|_\Sigma = \phi_1$ .*


Analogue for the Einstein equations is a highly nontrivial problem in elliptic PDE, pioneered by Corvino–Schoen '06.



## Characteristic/null gluing for the linear wave equation




# Characteristic/null gluing for the linear wave equation



Analogy:

- ▶  $\mathfrak{R}_1$  is Minkowski
- ▶  $\mathfrak{R}_2$  is Schwarzschild
- ▶  $C$  will be the event horizon  $\mathcal{H}^+$


# Characteristic/null gluing for the linear wave equation



Analogy:

- ▶  $\mathfrak{R}_1$  is Minkowski
- ▶  $\mathfrak{R}_2$  is Schwarzschild
- ▶  $C$  will be the event horizon  $\mathcal{H}^+$

# Characteristic/null gluing for the linear wave equation



Analogy:

- ▶  $\mathfrak{R}_1$  is Minkowski
- ▶  $\mathfrak{R}_2$  is Schwarzschild
- ▶  $C$  will be the event horizon  $\mathcal{H}^+$

## The characteristic initial value problem

**Bifurcate null hypersurface:**  $C \cup \underline{C}$  where  $C$  and  $\underline{C}$  meet transversally,  $C \cap \underline{C} \approx S^2$

## The characteristic initial value problem

**Bifurcate null hypersurface:**  $C \cup \underline{C}$  where  $C$  and  $\underline{C}$  meet transversally,  $C \cap \underline{C} \approx S^2$

### **Proposition (Characteristic IVP for the wave equation).**

*Given a bifurcate null hypersurface  $C \cup \underline{C}$  and  $\phi_0 : C \cup \underline{C} \rightarrow \mathbb{R}$ , there exists a unique function  $\phi$  defined to the future of  $C \cup \underline{C}$  such that*

$$\square_g \phi = 0 \tag{*}$$

*and  $\phi|_{C \cup \underline{C}} = \phi_0$ .*

## The characteristic initial value problem


**Bifurcate null hypersurface:**  $C \cup \underline{C}$  where  $C$  and  $\underline{C}$  meet transversally,  $C \cap \underline{C} \approx S^2$

### Proposition (Characteristic IVP for the wave equation).

Given a bifurcate null hypersurface  $C \cup \underline{C}$  and  $\phi_0 : C \cup \underline{C} \rightarrow \mathbb{R}$ , there exists a unique function  $\phi$  defined to the future of  $C \cup \underline{C}$  such that

$$\square_g \phi = 0 \quad (*)$$

and  $\phi|_{C \cup \underline{C}} = \phi_0$ .



## The characteristic initial value problem


**Bifurcate null hypersurface:**  $C \cup \underline{C}$  where  $C$  and  $\underline{C}$  meet transversally,  $C \cap \underline{C} \approx S^2$

### Proposition (Characteristic IVP for the wave equation).

Given a bifurcate null hypersurface  $C \cup \underline{C}$  and  $\phi_0 : C \cup \underline{C} \rightarrow \mathbb{R}$ , there exists a unique function  $\phi$  defined to the future of  $C \cup \underline{C}$  such that

$$\square_g \phi = 0 \tag{*}$$

and  $\phi|_{C \cup \underline{C}} = \phi_0$ .





## The characteristic initial value problem


**Bifurcate null hypersurface:**  $C \cup \underline{C}$  where  $C$  and  $\underline{C}$  meet transversally,  $C \cap \underline{C} \approx S^2$

### Proposition (Characteristic IVP for the wave equation).

Given a bifurcate null hypersurface  $C \cup \underline{C}$  and  $\phi_0 : C \cup \underline{C} \rightarrow \mathbb{R}$ , there exists a unique function  $\phi$  defined to the future of  $C \cup \underline{C}$  such that

$$\square_g \phi = 0 \tag{*}$$

and  $\phi|_{C \cup \underline{C}} = \phi_0$ .



Characteristic data for the wave equation does not involve derivatives.

## The null “constraints”

In any well posed initial value problem, there needs to be a mechanism to compute the full solution on the initial data hypersurface.

# The null “constraints”

In any well posed initial value problem, there needs to be a mechanism to compute the full solution on the initial data hypersurface. Toy problem for the toy problem:

- ▶ Wave equation on Minkowski  $\mathbb{R}^{3+1}$
- ▶  $u = t - r$ ,  $v = t + r$  double null coordinates

# The null “constraints”

In any well posed initial value problem, there needs to be a mechanism to compute the full solution on the initial data hypersurface. Toy problem for the toy problem:

- ▶ Wave equation on Minkowski  $\mathbb{R}^{3+1}$
- ▶  $u = t - r$ ,  $v = t + r$  double null coordinates
- ▶  $\phi = \phi(t, r, \vartheta, \varphi) = \phi(u, v, \vartheta, \varphi)$  satisfies

$$\partial_u \partial_v (r\phi) = \Delta(r\phi) \quad (*)$$

# The null “constraints”

In any well posed initial value problem, there needs to be a mechanism to compute the full solution on the initial data hypersurface. Toy problem for the toy problem:

- ▶ Wave equation on Minkowski  $\mathbb{R}^{3+1}$
- ▶  $u = t - r$ ,  $v = t + r$  double null coordinates
- ▶  $\phi = \phi(t, r, \vartheta, \varphi) = \phi(u, v, \vartheta, \varphi)$  satisfies

$$\partial_u \partial_v (r\phi) = \Delta(r\phi) \quad (*)$$

- ▶ Integrate  $(*)$  on  $C$ :

$$\partial_u(r\phi)|_{v_2} - \partial_u(r\phi)|_{v_1} = \int_{v_1}^{v_2} \Delta(r\phi) dv'$$

# The null “constraints”

In any well posed initial value problem, there needs to be a mechanism to compute the full solution on the initial data hypersurface. Toy problem for the toy problem:

- ▶ Wave equation on Minkowski  $\mathbb{R}^{3+1}$
- ▶  $u = t - r$ ,  $v = t + r$  double null coordinates
- ▶  $\phi = \phi(t, r, \vartheta, \varphi) = \phi(u, v, \vartheta, \varphi)$  satisfies

$$\partial_u \partial_v (r\phi) = \Delta(r\phi) \quad (*)$$

- ▶ Integrate  $(*)$  on  $C$ :

$$\partial_u(r\phi)|_{v_2} - \partial_u(r\phi)|_{v_1} = \int_{v_1}^{v_2} \Delta(r\phi) dv'$$

- ▶  $\implies \partial_u \phi|_C$  can be calculated from (1)  $\partial_u \phi$  at the bifurcation sphere and (2)  $\phi|_C$

# The null “constraints”

In any well posed initial value problem, there needs to be a mechanism to compute the full solution on the initial data hypersurface. Toy problem for the toy problem:

- ▶ Wave equation on Minkowski  $\mathbb{R}^{3+1}$
- ▶  $u = t - r$ ,  $v = t + r$  double null coordinates
- ▶  $\phi = \phi(t, r, \vartheta, \varphi) = \phi(u, v, \vartheta, \varphi)$  satisfies

$$\partial_u \partial_v (r\phi) = \Delta(r\phi) \quad (*)$$

- ▶ Integrate  $(*)$  on  $C$ :

$$\partial_u(r\phi)|_{v_2} - \partial_u(r\phi)|_{v_1} = \int_{v_1}^{v_2} \Delta(r\phi) dv'$$

- ▶  $\implies \partial_u \phi|_C$  can be calculated from (1)  $\partial_u \phi$  at the bifurcation sphere and (2)  $\phi|_C$
- ▶  $(*)$  lets us compute  $\partial_u^i \phi$  on  $C$  to all orders

## The null “constraints”

In general, the wave equation can be interpreted as a transport equation for the transverse derivative of  $\phi$ , sourced by  $\phi$  and tangential derivatives (which are a part of the characteristic data)



## The null “constraints”

In general, the wave equation can be interpreted as a transport equation for the transverse derivative of  $\phi$ , sourced by  $\phi$  and tangential derivatives (which are a part of the characteristic data)

On  $C$ , the wave equation is equivalent to

$$(\partial_v + \text{known function})\partial_u\phi = \text{known function},$$

where “known function” depends on underlying geometry and data.

## The null “constraints”

In general, the wave equation can be interpreted as a transport equation for the transverse derivative of  $\phi$ , sourced by  $\phi$  and tangential derivatives (which are a part of the characteristic data)

On  $C$ , the wave equation is equivalent to

$$(\partial_v + \text{known function})\partial_u\phi = \text{known function},$$

where “known function” depends on underlying geometry and data.

Commuting the wave equation with transverse derivatives gives an inductive system of transport equations for all orders!

## The null “constraints”

In general, the wave equation can be interpreted as a transport equation for the transverse derivative of  $\phi$ , sourced by  $\phi$  and tangential derivatives (which are a part of the characteristic data)

On  $C$ , the wave equation is equivalent to


$$(\partial_v + \text{known function})\partial_u\phi = \text{known function},$$

where “known function” depends on underlying geometry and data.

Commuting the wave equation with transverse derivatives gives an inductive system of transport equations for all orders!


These equations have become known as the *null “constraints”* and are very distinct from the “usual” constraint equations for Einstein.

## Back to characteristic gluing



The  $C^k$  characteristic gluing problem reduces to:


## Back to characteristic gluing



The  $C^k$  characteristic gluing problem reduces to:

- ▶ Given two spheres  $S_1$  and  $S_2$  along an outgoing null cone  $C$
- ▶ Given  $k$  ingoing and outgoing derivatives of  $\phi$  at  $S_1$  and  $S_2$


## Back to characteristic gluing



The  $C^k$  characteristic gluing problem reduces to:

- ▶ Given two spheres  $S_1$  and  $S_2$  along an outgoing null cone  $C$
- ▶ Given  $k$  ingoing and outgoing derivatives of  $\phi$  at  $S_1$  and  $S_2$
- ▶ Prescribe  $\phi$  along the part of  $C$  between  $S_1$  and  $S_2$  so that the outgoing derivatives agree with the given ones and the solutions of the transport equations for the ingoing derivatives have the specified initial and final values.

## Back to characteristic gluing



The  $C^k$  characteristic gluing problem reduces to:

- ▶ Given two spheres  $S_1$  and  $S_2$  along an outgoing null cone  $C$
- ▶ Given  $k$  ingoing and outgoing derivatives of  $\phi$  at  $S_1$  and  $S_2$
- ▶ Prescribe  $\phi$  along the part of  $C$  between  $S_1$  and  $S_2$  so that the outgoing derivatives agree with the given ones and the solutions of the transport equations for the ingoing derivatives have the specified initial and final values.
- ▶ Initial/boundary value problem for a system of coupled ODEs where you are allowed to (almost) freely prescribe the potential

## Einstein-scalar field in spherical symmetry

►  $\mathcal{M}^{3+1} = \mathcal{Q}^{1+1} \times S^2$

$$g = -\Omega^2 du dv + r^2 g_{S^2}$$

►  $\Omega(u, v) > 0$  lapse,  $r(u, v) > 0$  area-radius



# Einstein-scalar field in spherical symmetry

►  $\mathcal{M}^{3+1} = \mathcal{Q}^{1+1} \times S^2$

$$g = -\Omega^2 du dv + r^2 g_{S^2}$$

►  $\Omega(u, v) > 0$  lapse,  $r(u, v) > 0$  area-radius

► Wave equations

$$\partial_u \partial_v \phi = -\frac{\partial_u \phi \partial_v r}{r} - \frac{\partial_u r \partial_v \phi}{r} \quad (6)$$

$$\partial_u \partial_v r = -\frac{\Omega^2}{4r} - \frac{\partial_u r \partial_v r}{r} \quad (7)$$

$$\partial_u \partial_v \log(\Omega^2) = \frac{\Omega^2}{2r^2} + 2 \frac{\partial_u r \partial_v r}{r^2} \quad (8)$$

# Einstein-scalar field in spherical symmetry

►  $\mathcal{M}^{3+1} = \mathcal{Q}^{1+1} \times S^2$

$$g = -\Omega^2 du dv + r^2 g_{S^2}$$

►  $\Omega(u, v) > 0$  lapse,  $r(u, v) > 0$  area-radius

► Wave equations

$$\partial_u \partial_v \phi = -\frac{\partial_u \phi \partial_v r}{r} - \frac{\partial_u r \partial_v \phi}{r} \quad (6)$$

$$\partial_u \partial_v r = -\frac{\Omega^2}{4r} - \frac{\partial_u r \partial_v r}{r} \quad (7)$$


$$\partial_u \partial_v \log(\Omega^2) = \frac{\Omega^2}{2r^2} + 2 \frac{\partial_u r \partial_v r}{r^2} \quad (8)$$

► Raychaudhuri's equations (constraints)


$$\partial_u \left( \frac{\partial_u r}{\Omega^2} \right) = -\frac{r}{\Omega^2} (\partial_u \phi)^2 \quad (9)$$

$$\partial_v \left( \frac{\partial_v r}{\Omega^2} \right) = -\frac{r}{\Omega^2} (\partial_v \phi)^2 \quad (10)$$


# Minkowski to Schwarzschild gluing



# Minkowski to Schwarzschild gluing




# Minkowski to Schwarzschild gluing



# Minkowski to Schwarzschild gluing

## Theorem (Kehle–U. '22).

*For any  $k \in \mathbb{N}$  and  $0 < R_i < 2M_f$ , the Minkowski sphere of radius  $R_i$  can be characteristically glued to the Schwarzschild event horizon sphere with mass  $M_f$  to order  $C^k$  within the Einstein-scalar field model in spherical symmetry.*



## Idea of the proof

- ▶ Null constraint system becomes a coupled nonlinear system of ODEs sourced by  $\phi(v)$

## Idea of the proof

- ▶ Null constraint system becomes a coupled nonlinear system of ODEs sourced by  $\phi(v)$
- ▶ Scalar field ansatz

$$\phi_\alpha(v) = \sum_{1 \leq j \leq k+1} \alpha_j \chi_j(v), \quad \alpha \in \mathbb{R}^{k+1}$$



## Idea of the proof

- ▶ Null constraint system becomes a coupled nonlinear system of ODEs sourced by  $\phi(v)$
- ▶ Scalar field ansatz

$$\phi_\alpha(v) = \sum_{1 \leq j \leq k+1} \alpha_j \chi_j(v), \quad \alpha \in \mathbb{R}^{k+1}$$

- ▶ Initialize  $\partial_u \phi_\alpha(0) = \dots = \partial_u^k \phi_\alpha(0) = 0$

## Idea of the proof

- ▶ Null constraint system becomes a coupled nonlinear system of ODEs sourced by  $\phi(v)$
- ▶ Scalar field ansatz

$$\phi_\alpha(v) = \sum_{1 \leq j \leq k+1} \alpha_j \chi_j(v), \quad \alpha \in \mathbb{R}^{k+1}$$

- ▶ Initialize  $\partial_u \phi_\alpha(0) = \dots = \partial_u^k \phi_\alpha(0) = 0$
- ▶ Initialize  $r(v)$  teleologically at  $v = 1$  ( $\partial_v r(0)$  is a gauge choice, but  $\partial_v r(1) = 0!$ )

## Idea of the proof

- ▶ Null constraint system becomes a coupled nonlinear system of ODEs sourced by  $\phi(v)$
- ▶ Scalar field ansatz

$$\phi_\alpha(v) = \sum_{1 \leq j \leq k+1} \alpha_j \chi_j(v), \quad \alpha \in \mathbb{R}^{k+1}$$

- ▶ Initialize  $\partial_u \phi_\alpha(0) = \dots = \partial_u^k \phi_\alpha(0) = 0$
- ▶ Initialize  $r(v)$  teleologically at  $v = 1$  ( $\partial_v r(0)$  is a gauge choice, but  $\partial_v r(1) = 0!$ )
- ▶ For each  $\alpha$ , generate  $r(v)$ ,  $\partial_u r(v)$ ,  $\partial_u \phi_\alpha(v)$ , etc.

## Idea of the proof

- ▶ Null constraint system becomes a coupled nonlinear system of ODEs sourced by  $\phi(v)$
- ▶ Scalar field ansatz

$$\phi_\alpha(v) = \sum_{1 \leq j \leq k+1} \alpha_j \chi_j(v), \quad \alpha \in \mathbb{R}^{k+1}$$

- ▶ Initialize  $\partial_u \phi_\alpha(0) = \dots = \partial_u^k \phi_\alpha(0) = 0$
- ▶ Initialize  $r(v)$  teleologically at  $v = 1$  ( $\partial_v r(0)$  is a gauge choice, but  $\partial_v r(1) = 0!$ )
- ▶ For each  $\alpha$ , generate  $r(v)$ ,  $\partial_u r(v)$ ,  $\partial_u \phi_\alpha(v)$ , etc.
- ▶ The antipodal map  $\alpha \mapsto -\alpha$  leaves geometric quantities fixed ( $r$ ,  $\Omega^2$ , and derivatives)

## Idea of the proof

- ▶ Null constraint system becomes a coupled nonlinear system of ODEs sourced by  $\phi(v)$
- ▶ Scalar field ansatz

$$\phi_\alpha(v) = \sum_{1 \leq j \leq k+1} \alpha_j \chi_j(v), \quad \alpha \in \mathbb{R}^{k+1}$$

- ▶ Initialize  $\partial_u \phi_\alpha(0) = \dots = \partial_u^k \phi_\alpha(0) = 0$
- ▶ Initialize  $r(v)$  teleologically at  $v = 1$  ( $\partial_v r(0)$  is a gauge choice, but  $\partial_v r(1) = 0!$ )
- ▶ For each  $\alpha$ , generate  $r(v)$ ,  $\partial_u r(v)$ ,  $\partial_u \phi_\alpha(v)$ , etc.
- ▶ The antipodal map  $\alpha \mapsto -\alpha$  leaves geometric quantities fixed ( $r$ ,  $\Omega^2$ , and derivatives)
- ▶ The antipodal map  $\alpha \mapsto -\alpha$  changes the sign of the scalar field and its derivatives

## Idea of the proof

- ▶ Null constraint system becomes a coupled nonlinear system of ODEs sourced by  $\phi(v)$
- ▶ Scalar field ansatz

$$\phi_\alpha(v) = \sum_{1 \leq j \leq k+1} \alpha_j \chi_j(v), \quad \alpha \in \mathbb{R}^{k+1}$$

- ▶ Initialize  $\partial_u \phi_\alpha(0) = \dots = \partial_u^k \phi_\alpha(0) = 0$
- ▶ Initialize  $r(v)$  teleologically at  $v = 1$  ( $\partial_v r(0)$  is a gauge choice, but  $\partial_v r(1) = 0!$ )
- ▶ For each  $\alpha$ , generate  $r(v)$ ,  $\partial_u r(v)$ ,  $\partial_u \phi_\alpha(v)$ , etc.
- ▶ The antipodal map  $\alpha \mapsto -\alpha$  leaves geometric quantities fixed ( $r$ ,  $\Omega^2$ , and derivatives)
- ▶ The antipodal map  $\alpha \mapsto -\alpha$  changes the sign of the scalar field and its derivatives
- ▶ In particular, the map

$$\alpha \mapsto \left( \partial_u \phi_\alpha(1), \dots, \partial_u^k \phi_\alpha(1) \right)$$

## Idea of the proof

- ▶ Null constraint system becomes a coupled nonlinear system of ODEs sourced by  $\phi(v)$
- ▶ Scalar field ansatz

$$\phi_\alpha(v) = \sum_{1 \leq j \leq k+1} \alpha_j \chi_j(v), \quad \alpha \in \mathbb{R}^{k+1}$$

- ▶ Initialize  $\partial_u \phi_\alpha(0) = \dots = \partial_u^k \phi_\alpha(0) = 0$
- ▶ Initialize  $r(v)$  teleologically at  $v = 1$  ( $\partial_v r(0)$  is a gauge choice, but  $\partial_v r(1) = 0!$ )
- ▶ For each  $\alpha$ , generate  $r(v)$ ,  $\partial_u r(v)$ ,  $\partial_u \phi_\alpha(v)$ , etc.
- ▶ The antipodal map  $\alpha \mapsto -\alpha$  leaves geometric quantities fixed ( $r$ ,  $\Omega^2$ , and derivatives)
- ▶ The antipodal map  $\alpha \mapsto -\alpha$  changes the sign of the scalar field and its derivatives
- ▶ In particular, the map


$$\alpha \mapsto \left( \partial_u \phi_\alpha(1), \dots, \partial_u^k \phi_\alpha(1) \right)$$

is odd

- ▶ **Borsuk–Ulam theorem:** there exists  $\alpha_*$  such that

$$\left( \partial_u \phi_{\alpha_*}(1), \dots, \partial_u^k \phi_{\alpha_*}(1) \right) = 0.$$

# Disproof of the third law





## The vacuum case

We believe that the charged scalar field is not necessary to provide a counterexample to the third law:

# The vacuum case

We believe that the charged scalar field is not necessary to provide a counterexample to the third law:

## Conjecture.

*There exist Cauchy data for the Einstein vacuum equations*

$$R_{\mu\nu} = 0$$

*which undergo gravitational collapse and form an exactly Schwarzschild apparent horizon, only for the spacetime to form an exactly extremal Kerr event horizon at a later advanced time. In particular, already in vacuum, the “third law of black hole thermodynamics” is false.*

# The vacuum case

We believe that the charged scalar field is not necessary to provide a counterexample to the third law:

## Conjecture.


*There exist Cauchy data for the Einstein vacuum equations*

$$R_{\mu\nu} = 0$$


*which undergo gravitational collapse and form an exactly Schwarzschild apparent horizon, only for the spacetime to form an exactly extremal Kerr event horizon at a later advanced time. In particular, already in vacuum, the “third law of black hole thermodynamics” is false.*

This would follow from characteristic gluing, in vacuum, of a Minkowski sphere to a Kerr exterior sphere.


## Outlook: the “moduli space” of gravitational (non-)collapse




## Outlook: the “moduli space” of gravitational (non-)collapse




# Outlook: the “moduli space” of gravitational (non-)collapse



## Outlook: the “moduli space” of gravitational (non-)collapse



# Outlook: the “moduli space” of gravitational (non-)collapse



Critical behavior: naked singularities expected when a “low energy” black hole forms.



## Critical behavior near extremality

- We expect that a generic perturbation of an extremal black hole will just make the parameters subextremal.

## Critical behavior near extremality

- ▶ We expect that a generic perturbation of an extremal black hole will just make the parameters subextremal.
- ▶ Despite the presence of an instability for extremal black holes, the **Aretakis instability**, a weak form of stability is believed to be true for extremal black holes.

# Critical behavior near extremality

- ▶ We expect that a generic perturbation of an extremal black hole will just make the parameters subextremal.
- ▶ Despite the presence of an instability for extremal black holes, the **Aretakis instability**, a weak form of stability is believed to be true for extremal black holes.

## Conjecture (Dafermos–Holzegel–Rodnianski–Taylor).

*The stable manifold of very slowly rotating extremal Kerr–Newman delimits the boundary between asymptotically subextremal black holes and spacetimes failing to collapse for a while longer.*

## Critical behavior near extremality

- ▶ We expect that a generic perturbation of an extremal black hole will just make the parameters subextremal.
- ▶ Despite the presence of an instability for extremal black holes, the **Aretakis instability**, a weak form of stability is believed to be true for extremal black holes.

### Conjecture (Dafermos–Holzegel–Rodnianski–Taylor).

*The stable manifold of very slowly rotating extremal Kerr–Newman delimits the boundary between asymptotically subextremal black holes and spacetimes failing to collapse for a while longer.*

Critical behavior: extremal black holes can be perturbed so black hole formation is delayed and the event horizon jumps just as the extremal threshold is crossed.

## Critical behavior near extremality

- ▶ We expect that a generic perturbation of an extremal black hole will just make the parameters subextremal.
- ▶ Despite the presence of an instability for extremal black holes, the **Aretakis instability**, a weak form of stability is believed to be true for extremal black holes.

### Conjecture (Dafermos–Holzegel–Rodnianski–Taylor).

*The stable manifold of very slowly rotating extremal Kerr–Newman delimits the boundary between asymptotically subextremal black holes and spacetimes failing to collapse for a while longer.*

Critical behavior: extremal black holes can be perturbed so black hole formation is delayed and the event horizon jumps just as the extremal threshold is crossed.

### Conjecture.

*There exist regular Cauchy data for the Einstein–Maxwell-charged scalar field system, leading to the formation of an extremal Reissner–Nordström black hole, that can be perturbed to disperse, i.e. no black hole forms.*

## Critical behavior near extremality

- ▶ We expect that a generic perturbation of an extremal black hole will just make the parameters subextremal.
- ▶ Despite the presence of an instability for extremal black holes, the **Aretakis instability**, a weak form of stability is believed to be true for extremal black holes.

### Conjecture (Dafermos–Holzegel–Rodnianski–Taylor).


*The stable manifold of very slowly rotating extremal Kerr–Newman delimits the boundary between asymptotically subextremal black holes and spacetimes failing to collapse for a while longer.*

Critical behavior: extremal black holes can be perturbed so black hole formation is delayed and the event horizon jumps just as the extremal threshold is crossed.

### Conjecture.

*There exist regular Cauchy data for the Einstein–Maxwell-charged scalar field system, leading to the formation of an extremal Reissner–Nordström black hole, that can be perturbed to disperse, i.e. no black hole forms.*

Critical behavior: the very black-holeness of certain spacetimes is unstable.



*Thank you!*