

Retiring the third law of black hole thermodynamics

Ryan Unger

Department of Mathematics, Princeton University

Crete Center for Theoretical Physics
May 2023

joint work with Christoph Kehle (ETH Zürich)
[arXiv:2211.15742](https://arxiv.org/abs/2211.15742), [arXiv:2304.08455](https://arxiv.org/abs/2304.08455)

Outline

1. What is the third law?
2. Main result: the third law is false
3. Geometry and physics of third law violating spacetimes
4. Extremal black hole formation as a critical phenomenon

Black hole thermodynamics

Commun. math. Phys. 31, 161–170 (1973)
© by Springer-Verlag 1973

The Four Laws of Black Hole Mechanics

J. M. Bardeen*

Department of Physics, Yale University, New Haven, Connecticut, USA

B. Carter and S. W. Hawking

Institute of Astronomy, University of Cambridge, England

Received January 24, 1973

Black hole thermodynamics

Commun. math. Phys. 31, 161–170 (1973)
© by Springer-Verlag 1973

The Four Laws of Black Hole Mechanics

J. M. Bardeen*

Department of Physics, Yale University, New Haven, Connecticut, USA

B. Carter and S. W. Hawking

Institute of Astronomy, University of Cambridge, England

Received January 24, 1973

Black hole thermodynamics is a proposed close mathematical **analogy** between **classical** black hole dynamics and classical thermodynamics.

Black hole thermodynamics

Commun. math. Phys. 31, 161–170 (1973)
© by Springer-Verlag 1973

The Four Laws of Black Hole Mechanics

J. M. Bardeen*

Department of Physics, Yale University, New Haven, Connecticut, USA

B. Carter and S. W. Hawking

Institute of Astronomy, University of Cambridge, England

Received January 24, 1973

Black hole thermodynamics is a proposed close mathematical analogy between classical black hole dynamics and classical thermodynamics.

Fundamentally, these are statements about Einstein gravity coupled to classical matter fields (and are already interesting in vacuum!).

Black hole thermodynamics

Law	Classical thermodynamics	Black holes

Black hole thermodynamics

Law	Classical thermodynamics	Black holes
Zeroth	T constant in equilibrium	

Black hole thermodynamics

Law	Classical thermodynamics	Black holes
Zeroth	T constant in equilibrium	surface gravity κ constant on stationary horizon

Black hole thermodynamics

Law	Classical thermodynamics	Black holes
Zeroth	T constant in equilibrium	surface gravity κ constant on stationary horizon
First	$dE = TdS + \dots$	

Black hole thermodynamics

Law	Classical thermodynamics	Black holes
Zeroth	T constant in equilibrium	surface gravity κ constant on stationary horizon
First	$dE = TdS + \dots$	$dM = \kappa dA + \dots$

Black hole thermodynamics

Law	Classical thermodynamics	Black holes
Zeroth	T constant in equilibrium	surface gravity κ constant on stationary horizon
First	$dE = TdS + \dots$	$dM = \kappa dA + \dots$
Second	$dS \geq 0$	

Black hole thermodynamics

Law	Classical thermodynamics	Black holes
Zeroth	T constant in equilibrium	surface gravity κ constant on stationary horizon
First	$dE = TdS + \dots$	$dM = \kappa dA + \dots$
Second	$dS \geq 0$	$dA \geq 0$

Black hole thermodynamics

Law	Classical thermodynamics	Black holes
Zeroth	T constant in equilibrium	surface gravity κ constant on stationary horizon
First	$dE = TdS + \dots$	$dM = \kappa dA + \dots$
Second	$dS \geq 0$	$dA \geq 0$
Third	$T \not\rightarrow 0$ in finite process	

Black hole thermodynamics

Law	Classical thermodynamics	Black holes
Zeroth	T constant in equilibrium	surface gravity κ constant on stationary horizon
First	$dE = TdS + \dots$	$dM = \kappa dA + \dots$
Second	$dS \geq 0$	$dA \geq 0$
Third	$T \not\rightarrow 0$ in finite process	surface gravity $\kappa \not\rightarrow 0$ in finite advanced time

Black hole thermodynamics

Law	Classical thermodynamics	Black holes
Zeroth	T constant in equilibrium	surface gravity κ constant on stationary horizon
First	$dE = TdS + \dots$	$dM = \kappa dA + \dots$
Second	$dS \geq 0$	$dA \geq 0$
Third	$T \not\rightarrow 0$ in finite process	surface gravity $\kappa \not\rightarrow 0$ in finite advanced time

- Hawking radiation—interpret $\kappa \propto T$

Black hole thermodynamics

Law	Classical thermodynamics	Black holes
Zeroth	T constant in equilibrium	surface gravity κ constant on stationary horizon
First	$dE = TdS + \dots$	$dM = \kappa dA + \dots$
Second	$dS \geq 0$	$dA \geq 0$
Third	$T \not\rightarrow 0$ in finite process	surface gravity $\kappa \not\rightarrow 0$ in finite advanced time

- ▶ Hawking radiation—interpret $\kappa \propto T$
- ▶ Bekenstein entropy—interpret $A \propto S$

Black hole thermodynamics

Law	Classical thermodynamics	Black holes
Zeroth	T constant in equilibrium	surface gravity κ constant on stationary horizon
First	$dE = TdS + \dots$	$dM = \kappa dA + \dots$
Second	$dS \geq 0$	$dA \geq 0$
Third	$T \not\rightarrow 0$ in finite process	surface gravity $\kappa \not\rightarrow 0$ in finite advanced time


- ▶ Hawking radiation—interpret $\kappa \propto T$
- ▶ Bekenstein entropy—interpret $A \propto S$
- ▶ These identifications are further justified physically using QFT, but are **formal analogies** in classical GR.

Black hole thermodynamics


Law	Classical thermodynamics	Black holes
Zeroth	T constant in equilibrium	surface gravity κ constant on stationary horizon
First	$dE = TdS + \dots$	$dM = \kappa dA + \dots$
Second	$dS \geq 0$	$dA \geq 0$
Third	$T \not\rightarrow 0$ in finite process	surface gravity $\kappa \not\rightarrow 0$ in finite advanced time

- ▶ Hawking radiation—interpret $\kappa \propto T$
- ▶ Bekenstein entropy—interpret $A \propto S$
- ▶ These identifications are further justified physically using QFT, but are **formal analogies** in classical GR.
- ▶ Laws 0, 1, and 2 proved by Hawking, Carter, Bardeen–Carter–Hawking, Wald, ...

Refresher on Schwarzschild




Refresher on Schwarzschild




Maximally extended Schwarzschild is the unique maximal Cauchy development of the data induced on a spacelike hypersurface Σ as depicted here.

Refresher on Schwarzschild



The black hole interior is foliated by **trapped surfaces** (both future null expansions negative).

Refresher on Schwarzschild




The black hole interior is foliated by **trapped surfaces** (both future null expansions negative).

Theorem (Penrose's incompleteness theorem).


If an asymptotically flat spacetime contains a trapped surface and satisfies the null energy condition, it is geodesically incomplete.

Refresher on gravitational collapse




Penrose diagram of gravitational collapse. One-ended Cauchy data!


Refresher on gravitational collapse




Refresher on subextremal Reissner–Nordström: $0 < |e| < M$




Refresher on subextremal Reissner–Nordström: $0 < |e| < M$




Refresher on subextremal Reissner–Nordström: $0 < |e| < M$




Refresher on subextremal Reissner–Nordström: $0 < |e| < M$




Refresher on extremal Reissner–Nordström: $0 < |e| = M$




Refresher on extremal Reissner–Nordström: $0 < |e| = M$




Refresher on extremal Reissner–Nordström: $0 < |e| = M$




Refresher on extremal Reissner–Nordström: $0 < |e| = M$



Refresher on superextremal Reissner–Nordström: $0 < M < |e|$



Refresher on superextremal Reissner–Nordström: $0 < M < |e|$



Surface gravity κ of Reissner–Nordström

- ▶ $\nabla_T T = \kappa T$ on \mathcal{H}^+
- ▶ RN with mass M and charge e , $|e| \leq M$, has

$$\kappa = 2\pi T = \frac{\sqrt{M^2 - e^2}}{(M + \sqrt{M^2 - e^2})^2}$$

- ▶ **Subextremal:** $\kappa > 0$
- ▶ **Extremal:** $\kappa = 0$

The third law

Original formulation of Bardeen–Carter–Hawking:

Extending the analogy even further one would postulate:

The Third Law

It is impossible by any procedure, no matter how idealized, to reduce κ to zero by a finite sequence of operations.

Difficult to interpret κ dynamically!

The third law

Original formulation of Bardeen–Carter–Hawking:

Extending the analogy even further one would postulate:

The Third Law

It is impossible by any procedure, no matter how idealized, to reduce κ to zero by a finite sequence of operations.

Difficult to interpret κ dynamically!

Statement revised by Israel '86:

VOLUME 57, NUMBER 4

PHYSICAL REVIEW LETTERS

28 JULY 1986

Third Law of Black-Hole Dynamics: A Formulation

W. Israel^(a)

Research Institute for Fundamental Physics, Yukawa Hall, Kyoto University, Kyoto 606, Japan

(Received 19 May 1986)

The third law

Original formulation of Bardeen–Carter–Hawking:

Extending the analogy even further one would postulate:

The Third Law

It is impossible by any procedure, no matter how idealized, to reduce κ to zero by a finite sequence of operations.

Difficult to interpret κ dynamically!

Statement revised by Israel '86:

VOLUME 57, NUMBER 4

PHYSICAL REVIEW LETTERS

28 JULY 1986

Third Law of Black-Hole Dynamics: A Formulation and Proof

W. Israel^(a)

Research Institute for Fundamental Physics, Yukawa Hall, Kyoto University, Kyoto 606, Japan

(Received 19 May 1986)

Israel's formulation of the third law

A nonextremal black hole cannot become extremal (i.e., lose its trapped surfaces) at a finite advanced time in any continuous process in which the stress-energy tensor of accreted matter stays bounded and satisfies the weak energy condition in a neighborhood of the outer apparent horizon.

Israel's formulation of the third law

A nonextremal black hole cannot become extremal (i.e., lose its trapped surfaces) at a finite advanced time in any continuous process in which the stress-energy tensor of accreted matter stays bounded and satisfies the weak energy condition in a neighborhood of the outer apparent horizon.

1. “Finite advanced time” replaces “finite sequence of operations.”

Israel's formulation of the third law

A nonextremal black hole cannot become extremal (i.e., lose its trapped surfaces) at a finite advanced time in any continuous process in which the stress-energy tensor of accreted matter stays bounded and satisfies the weak energy condition in a neighborhood of the outer apparent horizon.

1. “Finite advanced time” replaces “finite sequence of operations.”
 - ▶ Brings the third law onto a similar footing as the second law, no quasistationary assumption.

Israel's formulation of the third law

A nonextremal black hole cannot become extremal (i.e., lose its trapped surfaces) at a finite advanced time in any continuous process in which the stress-energy tensor of accreted matter stays bounded and satisfies the weak energy condition in a neighborhood of the outer apparent horizon.

1. "Finite advanced time" replaces "finite sequence of operations."
 - ▶ Brings the third law onto a similar footing as the second law, no quasistationary assumption.
2. "Stress-energy tensor stays bounded" is a regularity condition.

Israel's formulation of the third law

A nonextremal black hole cannot become extremal (i.e., lose its trapped surfaces) at a finite advanced time in any continuous process in which the stress-energy tensor of accreted matter stays bounded and satisfies the weak energy condition in a neighborhood of the outer apparent horizon.

1. “Finite advanced time” replaces “finite sequence of operations.”
 - ▶ Brings the third law onto a similar footing as the second law, no quasistationary assumption.
2. “Stress-energy tensor stays bounded” is a regularity condition.
 - ▶ If singularities allowed, counterexample using massive dust shell. (Farrugia–Hajicek '79)

Israel's formulation of the third law

A nonextremal black hole cannot become extremal (i.e., lose its trapped surfaces) at a finite advanced time in any continuous process in which the stress-energy tensor of accreted matter stays bounded and satisfies the weak energy condition in a neighborhood of the outer apparent horizon.

1. “Finite advanced time” replaces “finite sequence of operations.”
 - ▶ Brings the third law onto a similar footing as the second law, no quasistationary assumption.
2. “Stress-energy tensor stays bounded” is a regularity condition.
 - ▶ If singularities allowed, counterexample using massive dust shell. (Farrugia–Hajicek '79)
3. Weak energy condition must be enforced.

Israel's formulation of the third law

A nonextremal black hole cannot become extremal (i.e., lose its trapped surfaces) at a finite advanced time in any continuous process in which the stress-energy tensor of accreted matter stays bounded and satisfies the weak energy condition in a neighborhood of the outer apparent horizon.

1. “Finite advanced time” replaces “finite sequence of operations.”
 - ▶ Brings the third law onto a similar footing as the second law, no quasistationary assumption.
2. “Stress-energy tensor stays bounded” is a regularity condition.
 - ▶ If singularities allowed, counterexample using massive dust shell. (Farrugia–Hajicek '79)
3. Weak energy condition must be enforced.
 - ▶ If no energy condition, counterexample using charged null dust. (Sullivan–Israel '80)

Israel's formulation of the third law

A nonextremal black hole cannot become extremal (i.e., lose its trapped surfaces) at a finite advanced time in any continuous process in which the stress-energy tensor of accreted matter stays bounded and satisfies the weak energy condition in a neighborhood of the outer apparent horizon.

1. “Finite advanced time” replaces “finite sequence of operations.”
 - ▶ Brings the third law onto a similar footing as the second law, no quasistationary assumption.
2. “Stress-energy tensor stays bounded” is a regularity condition.
 - ▶ If singularities allowed, counterexample using massive dust shell. (Farrugia–Hajicek '79)
3. Weak energy condition must be enforced.
 - ▶ If no energy condition, counterexample using charged null dust. (Sullivan–Israel '80)
4. Becoming extremal associated with “losing trapped surfaces.”

Israel's formulation of the third law

A nonextremal black hole cannot become extremal (i.e., lose its trapped surfaces) at a finite advanced time in any continuous process in which the stress-energy tensor of accreted matter stays bounded and satisfies the weak energy condition in a neighborhood of the outer apparent horizon.

1. “Finite advanced time” replaces “finite sequence of operations.”
 - ▶ Brings the third law onto a similar footing as the second law, no quasistationary assumption.
2. “Stress-energy tensor stays bounded” is a regularity condition.
 - ▶ If singularities allowed, counterexample using massive dust shell. (Farrugia–Hajicek '79)
3. Weak energy condition must be enforced.
 - ▶ If no energy condition, counterexample using charged null dust. (Sullivan–Israel '80)
4. Becoming extremal associated with “losing trapped surfaces.”
 - ▶ This parenthetical remark is related to the “mistake” in Israel’s paper and we will discuss it later.

Israel's formulation of the third law

A nonextremal black hole cannot become extremal (i.e., lose its trapped surfaces) at a finite advanced time in any continuous process in which the stress-energy tensor of accreted matter stays bounded and satisfies the weak energy condition in a neighborhood of the outer apparent horizon.

1. “Finite advanced time” replaces “finite sequence of operations.”
 - ▶ Brings the third law onto a similar footing as the second law, no quasistationary assumption.
2. “Stress-energy tensor stays bounded” is a regularity condition.
 - ▶ If singularities allowed, counterexample using massive dust shell. (Farrugia–Hajicek '79)
3. Weak energy condition must be enforced.
 - ▶ If no energy condition, counterexample using charged null dust. (Sullivan–Israel '80)
4. Becoming extremal associated with “losing trapped surfaces.”
 - ▶ This parenthetical remark is related to the “mistake” in Israel’s paper and we will discuss it later.
5. “In any continuous process” included because we are dealing with fundamentally nongeneric behavior.

Israel's formulation of the third law

A nonextremal black hole cannot become extremal (i.e., lose its trapped surfaces) at a finite advanced time in any continuous process in which the stress-energy tensor of accreted matter stays bounded and satisfies the weak energy condition in a neighborhood of the outer apparent horizon.

1. “Finite advanced time” replaces “finite sequence of operations.”
 - ▶ Brings the third law onto a similar footing as the second law, no quasistationary assumption.
2. “Stress-energy tensor stays bounded” is a regularity condition.
 - ▶ If singularities allowed, counterexample using massive dust shell. (Farrugia–Hajicek '79)
3. Weak energy condition must be enforced.
 - ▶ If no energy condition, counterexample using charged null dust. (Sullivan–Israel '80)
4. Becoming extremal associated with “losing trapped surfaces.”
 - ▶ This parenthetical remark is related to the “mistake” in Israel’s paper and we will discuss it later.
5. “In any continuous process” included because we are dealing with fundamentally nongeneric behavior.
 - ▶ Adding any kind of genericity assumption to the third law renders it a completely uninteresting and utterly trivial statement, which does not even use the Einstein equations!

Retiring the third law

Conjecture (The third law, BCH '73, Israel '86).

A subextremal black hole cannot become extremal in finite time by any continuous process, no matter how idealized, in which the spacetime and matter fields remain regular and obey the weak energy condition.

Retiring the third law

Conjecture (The third law, BCH '73, Israel '86).

A subextremal black hole cannot become extremal in finite time by any continuous process, no matter how idealized, in which the spacetime and matter fields remain regular and obey the weak energy condition.

Theorem (Kehle–U. '22).

Subextremal black holes can become extremal in finite time, evolving from regular initial data. In particular, the “third law of black hole thermodynamics” is false.

Retiring the third law

Conjecture (The third law, BCH '73, Israel '86).

A subextremal black hole cannot become extremal in finite time by any continuous process, no matter how idealized, in which the spacetime and matter fields remain regular and obey the weak energy condition.


Theorem (Kehle–U. '22).

Subextremal black holes can become extremal in finite time, evolving from regular initial data. In particular, the “third law of black hole thermodynamics” is false.

More precisely, there exist regular solutions of the Einstein–Maxwell-charged scalar field system (which satisfies the weak energy condition) with the following behavior:


Counterexample to the third law: detailed statement of our theorem

There exist **regular** solutions of the Einstein–Maxwell-charged scalar field system with the following behavior:



Counterexample to the third law: detailed statement of our theorem


There exist **regular** solutions of the Einstein–Maxwell-charged scalar field system with the following behavior:



- Spherically symmetric Cauchy data for the Einstein–Maxwell-charged scalar field system which undergo gravitational collapse.

Counterexample to the third law: detailed statement of our theorem


There exist **regular** solutions of the Einstein–Maxwell-charged scalar field system with the following behavior:



- ▶ Spherically symmetric Cauchy data for the Einstein–Maxwell-charged scalar field system which undergo gravitational collapse.
- ▶ Forms an exactly Schwarzschild “apparent horizon.”

Counterexample to the third law: detailed statement of our theorem


There exist **regular** solutions of the Einstein–Maxwell-charged scalar field system with the following behavior:



- ▶ Spherically symmetric Cauchy data for the Einstein–Maxwell-charged scalar field system which undergo gravitational collapse.
- ▶ Forms an exactly Schwarzschild “apparent horizon.”
- ▶ Forms an exactly extremal Reissner–Nordström event horizon at a later advanced time.

Counterexample to the third law: detailed statement of our theorem

There exist **regular** solutions of the Einstein–Maxwell-charged scalar field system with the following behavior:



- ▶ Spherically symmetric Cauchy data for the Einstein–Maxwell-charged scalar field system which undergo gravitational collapse.
- ▶ Forms an exactly Schwarzschild “apparent horizon.”
- ▶ Forms an exactly extremal Reissner–Nordström event horizon at a later advanced time.
- ▶ These solutions are arbitrarily **regular**: for any $k \in \mathbb{N}$, there exists a C^k example.

Einstein–Maxwell-charged scalar field system

- ▶ Lorentzian manifold (\mathcal{M}^{3+1}, g)
- ▶ 2-form $F = dA$ (electromagnetism)
- ▶ Charged (complex) massless scalar field ϕ

Einstein–Maxwell-charged scalar field system

- ▶ Lorentzian manifold (\mathcal{M}^{3+1}, g)
- ▶ 2-form $F = dA$ (electromagnetism)
- ▶ Charged (complex) massless scalar field ϕ
- ▶ Equations:

$$R_{\mu\nu}(g) - \frac{1}{2}R(g)g_{\mu\nu} = 2 \left(T_{\mu\nu}^{\text{EM}} + T_{\mu\nu}^{\text{CSF}} \right) \quad (1)$$

$$\nabla^\mu F_{\mu\nu} = 2\epsilon \operatorname{Im}(\phi \overline{D_\nu \phi}) \quad (2)$$

$$g^{\mu\nu} D_\mu D_\nu \phi = 0 \quad (3)$$

$$T_{\mu\nu}^{\text{EM}} = g^{\alpha\beta} F_{\alpha\nu} F_{\beta\mu} - \frac{1}{4} F^{\alpha\beta} F_{\alpha\beta} g_{\mu\nu} \quad (4)$$

$$T_{\mu\nu}^{\text{CSF}} = \operatorname{Re}(D_\mu \phi \overline{D_\nu \phi}) - \frac{1}{2} g_{\mu\nu} g^{\alpha\beta} D_\alpha \phi \overline{D_\beta \phi} \quad (5)$$

Einstein–Maxwell-charged scalar field system

- ▶ Lorentzian manifold (\mathcal{M}^{3+1}, g)
- ▶ 2-form $F = dA$ (electromagnetism)
- ▶ Charged (complex) massless scalar field ϕ
- ▶ Equations:

$$R_{\mu\nu}(g) - \frac{1}{2}R(g)g_{\mu\nu} = 2 \left(T_{\mu\nu}^{\text{EM}} + T_{\mu\nu}^{\text{CSF}} \right) \quad (1)$$

$$\nabla^\mu F_{\mu\nu} = 2\epsilon \operatorname{Im}(\phi \overline{D_\nu \phi}) \quad (2)$$

$$g^{\mu\nu} D_\mu D_\nu \phi = 0 \quad (3)$$

$$T_{\mu\nu}^{\text{EM}} = g^{\alpha\beta} F_{\alpha\nu} F_{\beta\mu} - \frac{1}{4} F^{\alpha\beta} F_{\alpha\beta} g_{\mu\nu} \quad (4)$$

$$T_{\mu\nu}^{\text{CSF}} = \operatorname{Re}(D_\mu \phi \overline{D_\nu \phi}) - \frac{1}{2} g_{\mu\nu} g^{\alpha\beta} D_\alpha \phi \overline{D_\beta \phi} \quad (5)$$

- ▶ Spherical symmetry, ϕ degree of freedom breaks rigidity from Birkhoff.

Einstein–Maxwell-charged scalar field system

- ▶ Lorentzian manifold (\mathcal{M}^{3+1}, g)
- ▶ 2-form $F = dA$ (electromagnetism)
- ▶ Charged (complex) massless scalar field ϕ
- ▶ Equations:

$$R_{\mu\nu}(g) - \frac{1}{2}R(g)g_{\mu\nu} = 2 \left(T_{\mu\nu}^{\text{EM}} + T_{\mu\nu}^{\text{CSF}} \right) \quad (1)$$

$$\nabla^\mu F_{\mu\nu} = 2\epsilon \operatorname{Im}(\phi \overline{D_\nu \phi}) \quad (2)$$

$$g^{\mu\nu} D_\mu D_\nu \phi = 0 \quad (3)$$

$$T_{\mu\nu}^{\text{EM}} = g^{\alpha\beta} F_{\alpha\nu} F_{\beta\mu} - \frac{1}{4} F^{\alpha\beta} F_{\alpha\beta} g_{\mu\nu} \quad (4)$$

$$T_{\mu\nu}^{\text{CSF}} = \operatorname{Re}(D_\mu \phi \overline{D_\nu \phi}) - \frac{1}{2} g_{\mu\nu} g^{\alpha\beta} D_\alpha \phi \overline{D_\beta \phi} \quad (5)$$

- ▶ Spherical symmetry, ϕ degree of freedom breaks rigidity from Birkhoff.
- ▶ Scalar field carries charge in order to charge up the black hole to extremality.

Einstein–Maxwell-charged scalar field system

- ▶ Lorentzian manifold (\mathcal{M}^{3+1}, g)
- ▶ 2-form $F = dA$ (electromagnetism)
- ▶ Charged (complex) massless scalar field ϕ
- ▶ Equations:

$$R_{\mu\nu}(g) - \frac{1}{2}R(g)g_{\mu\nu} = 2 \left(T_{\mu\nu}^{\text{EM}} + T_{\mu\nu}^{\text{CSF}} \right) \quad (1)$$

$$\nabla^\mu F_{\mu\nu} = 2\epsilon \operatorname{Im}(\phi \overline{D_\nu \phi}) \quad (2)$$

$$g^{\mu\nu} D_\mu D_\nu \phi = 0 \quad (3)$$

$$T_{\mu\nu}^{\text{EM}} = g^{\alpha\beta} F_{\alpha\nu} F_{\beta\mu} - \frac{1}{4} F^{\alpha\beta} F_{\alpha\beta} g_{\mu\nu} \quad (4)$$

$$T_{\mu\nu}^{\text{CSF}} = \operatorname{Re}(D_\mu \phi \overline{D_\nu \phi}) - \frac{1}{2} g_{\mu\nu} g^{\alpha\beta} D_\alpha \phi \overline{D_\beta \phi} \quad (5)$$

- ▶ Spherical symmetry, ϕ degree of freedom breaks rigidity from Birkhoff.
- ▶ Scalar field carries charge in order to charge up the black hole to extremality.
- ▶ Model satisfies the **dominant energy condition** (\implies **weak energy condition**).

Einstein–Maxwell-charged scalar field system

- ▶ Lorentzian manifold (\mathcal{M}^{3+1}, g)
- ▶ 2-form $F = dA$ (electromagnetism)
- ▶ Charged (complex) massless scalar field ϕ
- ▶ Equations:

$$R_{\mu\nu}(g) - \frac{1}{2}R(g)g_{\mu\nu} = 2 \left(T_{\mu\nu}^{\text{EM}} + T_{\mu\nu}^{\text{CSF}} \right) \quad (1)$$

$$\nabla^\mu F_{\mu\nu} = 2\epsilon \operatorname{Im}(\phi \overline{D_\nu \phi}) \quad (2)$$

$$g^{\mu\nu} D_\mu D_\nu \phi = 0 \quad (3)$$

$$T_{\mu\nu}^{\text{EM}} = g^{\alpha\beta} F_{\alpha\nu} F_{\beta\mu} - \frac{1}{4} F^{\alpha\beta} F_{\alpha\beta} g_{\mu\nu} \quad (4)$$

$$T_{\mu\nu}^{\text{CSF}} = \operatorname{Re}(D_\mu \phi \overline{D_\nu \phi}) - \frac{1}{2} g_{\mu\nu} g^{\alpha\beta} D_\alpha \phi \overline{D_\beta \phi} \quad (5)$$

- ▶ Spherical symmetry, ϕ degree of freedom breaks rigidity from Birkhoff.
- ▶ Scalar field carries charge in order to charge up the black hole to extremality.
- ▶ Model satisfies the **dominant energy condition** (\implies **weak energy condition**).
- ▶ Our proof works with a massive scalar field as long as $m \ll \epsilon$.

Losing trapped surfaces and the implicit assumption of connectivity

Losing trapped surfaces and the implicit assumption of connectivity

- Weak energy condition \implies trapped surfaces persist in evolution once formed.

Losing trapped surfaces and the implicit assumption of connectivity

- ▶ Weak energy condition \implies trapped surfaces persist in evolution once formed.
- ▶ Follows immediately from Raychaudhuri's equation

$$\frac{d\theta}{d\lambda} = -\frac{1}{2}\theta^2 - \hat{\sigma}^2 + \hat{\omega}^2 - R_{\mu\nu}k^\mu k^\nu.$$

Losing trapped surfaces and the implicit assumption of connectivity

- ▶ Weak energy condition \implies trapped surfaces persist in evolution once formed.
- ▶ Follows immediately from Raychaudhuri's equation

$$\frac{d\theta}{d\lambda} = -\tfrac{1}{2}\theta^2 - \hat{\sigma}^2 + \hat{\omega}^2 - R_{\mu\nu}k^\mu k^\nu.$$

- ▶ Extremal Reissner–Nordström (and Kerr) have no trapped surfaces, but their subextremal versions do.

Losing trapped surfaces and the implicit assumption of connectivity

- ▶ Weak energy condition \implies trapped surfaces persist in evolution once formed.
- ▶ Follows immediately from Raychaudhuri's equation

$$\frac{d\theta}{d\lambda} = -\tfrac{1}{2}\theta^2 - \hat{\sigma}^2 + \hat{\omega}^2 - R_{\mu\nu}k^\mu k^\nu.$$

- ▶ Extremal Reissner–Nordström (and Kerr) have no trapped surfaces, but their subextremal versions do.
- ▶ In his paper, Israel seems to have implicitly assumed that a regular solution will have a connected outermost apparent horizon.

Infractions can result from the absorption of infinitesimally thin, massive shells,⁵ which force the apparent horizon to jump outward discontinuously

Losing trapped surfaces and the implicit assumption of connectivity

- ▶ Weak energy condition \implies trapped surfaces persist in evolution once formed.
- ▶ Follows immediately from Raychaudhuri's equation

$$\frac{d\theta}{d\lambda} = -\tfrac{1}{2}\theta^2 - \hat{\sigma}^2 + \hat{\omega}^2 - R_{\mu\nu}k^\mu k^\nu.$$

- ▶ Extremal Reissner–Nordström (and Kerr) have no trapped surfaces, but their subextremal versions do.
- ▶ In his paper, Israel seems to have implicitly assumed that a regular solution will have a connected outermost apparent horizon.

Infractions can result from the absorption of infinitesimally thin, massive shells,⁵ which force the apparent horizon to jump outward discontinuously

This is not true. The outermost apparent horizon can jump in smooth spacetimes.

Losing trapped surfaces and the implicit assumption of connectivity

- ▶ Weak energy condition \implies trapped surfaces persist in evolution once formed.
- ▶ Follows immediately from Raychaudhuri's equation

$$\frac{d\theta}{d\lambda} = -\frac{1}{2}\theta^2 - \hat{\sigma}^2 + \hat{\omega}^2 - R_{\mu\nu}k^\mu k^\nu.$$


- ▶ Extremal Reissner–Nordström (and Kerr) have no trapped surfaces, but their subextremal versions do.
- ▶ In his paper, Israel seems to have implicitly assumed that a regular solution will have a connected outermost apparent horizon.

Infractions can result from the absorption of infinitesimally thin, massive shells,⁵ which force the apparent horizon to jump outward discontinuously


This is not true. The outermost apparent horizon can jump in smooth spacetimes.

- ▶ Under this implicit assumption, Israel concluded the third law to be true on the basis of Raychaudhuri's equation.

Israel's argument




Israel's paper reinterpreted



Outermost apparent horizon becomes disconnected the instant the black hole becomes extremal!

Israel's paper reinterpreted




Outermost apparent horizon becomes disconnected the instant the black hole becomes extremal!

However, this is a feature, not a glitch!

Gravitational collapse for any charge to mass ratio

Theorem (Kehle–U.).


For any $q \in [-1, 1]$, there exist regular Cauchy data for the Einstein–Maxwell-charged scalar field system whose evolution has the following Penrose diagram:



Gravitational collapse for any charge to mass ratio

Theorem (Kehle–U.).

For any $q \in [-1, 1]$, there exist regular Cauchy data for the Einstein–Maxwell-charged scalar field system whose evolution has the following Penrose diagram:




- From the point of view of our construction, the extremal case is exactly the same as the subextremal case!

Gravitational collapse for any charge to mass ratio

Theorem (Kehle–U.).

For any $q \in [-1, 1]$, there exist regular Cauchy data for the Einstein–Maxwell-charged scalar field system whose evolution has the following Penrose diagram:




- ▶ From the point of view of our construction, the extremal case is exactly the same as the subextremal case!
- ▶ $|q| \rightarrow 1$ represents a regular limit.

Gravitational collapse for any charge to mass ratio


Theorem (Kehle–U.).

For any $q \in [-1, 1]$, there exist regular Cauchy data for the Einstein–Maxwell-charged scalar field system whose evolution has the following Penrose diagram:




- ▶ From the point of view of our construction, the extremal case is exactly the same as the subextremal case!
- ▶ $|q| \rightarrow 1$ represents a regular limit.
- ▶ $T = 0$ is not fundamentally different than $T > 0$ in these examples.

Interior structure of third law violating solutions




Interior structure of third law violating solutions



- The outermost apparent horizon becomes disconnected, yet the spacetime is highly regular.


Interior structure of third law violating solutions



- ▶ The outermost apparent horizon becomes disconnected, yet the spacetime is highly regular.
- ▶ Trapped surfaces persist for all time and are found at any retarded time in the black hole interior.


So where do the trapped surfaces go?

The geometry of a $|q| = 1 - \varepsilon$ example converges to a $|q| = 1$ example as $\varepsilon \rightarrow 0$.




So where do the trapped surfaces go?

The geometry of a $|q| = 1 - \varepsilon$ example converges to a $|q| = 1$ example as $\varepsilon \rightarrow 0$.




So where do the trapped surfaces go?

The geometry of a $|q| = 1 - \varepsilon$ example converges to a $|q| = 1$ example as $\varepsilon \rightarrow 0$.



So where do the trapped surfaces go?

The geometry of a $|q| = 1 - \varepsilon$ example converges to a $|q| = 1$ example as $\varepsilon \rightarrow 0$.



Genericity

The Third Law

It is impossible by any procedure, no matter how idealized, to reduce κ to zero by a finite sequence of operations.

A nonextremal black hole cannot become extremal (i.e., lose its trapped surfaces) at a finite advanced time in any continuous process in which the stress-energy tensor of accreted matter stays bounded and satisfies the weak energy condition in a neighborhood of the outer apparent horizon.

Genericity

The Third Law

It is impossible by any procedure, no matter how idealized, to reduce κ to zero by a finite sequence of operations.

A nonextremal black hole cannot become extremal (i.e., lose its trapped surfaces) at a finite advanced time in any continuous process in which the stress-energy tensor of accreted matter stays bounded and satisfies the weak energy condition in a neighborhood of the outer apparent horizon.

- ▶ Keeping a horizon at exactly constant temperature, any temperature, is exceptional.

Genericity

The Third Law

It is impossible by any procedure, no matter how idealized, to reduce κ to zero by a finite sequence of operations.

A nonextremal black hole cannot become extremal (i.e., lose its trapped surfaces) at a finite advanced time in any continuous process in which the stress-energy tensor of accreted matter stays bounded and satisfies the weak energy condition in a neighborhood of the outer apparent horizon.

- ▶ Keeping a horizon at exactly constant temperature, any temperature, is exceptional.
- ▶ For example, the set of solutions which become exactly Schwarzschild after finite time is infinite codimension in the moduli space of solutions.

Genericity

The Third Law

It is impossible by any procedure, no matter how idealized, to reduce κ to zero by a finite sequence of operations.

A nonextremal black hole cannot become extremal (i.e., lose its trapped surfaces) at a finite advanced time in any continuous process in which the stress-energy tensor of accreted matter stays bounded and satisfies the weak energy condition in a neighborhood of the outer apparent horizon.

- ▶ Keeping a horizon at exactly constant temperature, any temperature, is exceptional.
- ▶ For example, the set of solutions which become exactly Schwarzschild after finite time is infinite codimension in the moduli space of solutions.
- ▶ In view of our results, forming $T = 0$ in finite time is not different or more difficult than forming fixed $T > 0$.

Genericity

The Third Law

It is impossible by any procedure, no matter how idealized, to reduce κ to zero by a finite sequence of operations.

A nonextremal black hole cannot become extremal (i.e., lose its trapped surfaces) at a finite advanced time in any continuous process in which the stress-energy tensor of accreted matter stays bounded and satisfies the weak energy condition in a neighborhood of the outer apparent horizon.

- ▶ Keeping a horizon at exactly constant temperature, any temperature, is exceptional.
- ▶ For example, the set of solutions which become exactly Schwarzschild after finite time is infinite codimension in the moduli space of solutions.
- ▶ In view of our results, forming $T = 0$ in finite time is not different or more difficult than forming fixed $T > 0$.
- ▶ One might try to save the third law by demanding a process that is stable under perturbations.

The Third Law

It is impossible by any procedure, no matter how idealized, to reduce κ to zero by a finite sequence of operations.

A nonextremal black hole cannot become extremal (i.e., lose its trapped surfaces) at a finite advanced time in any continuous process in which the stress-energy tensor of accreted matter stays bounded and satisfies the weak energy condition in a neighborhood of the outer apparent horizon.

- ▶ Keeping a horizon at exactly constant temperature, any temperature, is exceptional.
- ▶ For example, the set of solutions which become exactly Schwarzschild after finite time is infinite codimension in the moduli space of solutions.
- ▶ In view of our results, forming $T = 0$ in finite time is not different or more difficult than forming fixed $T > 0$.
- ▶ One might try to save the third law by demanding a process that is stable under perturbations.
- ▶ However, such a reformulation would be utterly trivial (and has nothing to do with $T = 0$).

The third law and supercharging

Bardeen–Carter–Hawking:


Another reason for believing the third law is that if one could reduce κ to zero by a finite sequence of operations, then presumably one could carry the process further, thereby creating a naked singularity.

The third law and supercharging

Bardeen–Carter–Hawking:

Another reason for believing the third law is that if one could reduce κ to zero by a finite sequence of operations, then presumably one could carry the process further, thereby creating a naked singularity.

- ▶ Charging (resp., spinning) up a black hole to create a naked singularity as in superextremal RN (resp., Kerr) is known as **supercharging** (resp., **overspinning**).




The third law and supercharging

Bardeen–Carter–Hawking:

Another reason for believing the third law is that if one could reduce κ to zero by a finite sequence of operations, then presumably one could carry the process further, thereby creating a naked singularity.

- ▶ Charging (resp., spinning) up a black hole to create a naked singularity as in superextremal RN (resp., Kerr) is known as **supercharging** (resp., **overspinning**).




- ▶ This is a doomed endeavor because the naked singularity in extremal RN/Kerr is dynamically inaccessible.

The third law and supercharging

Bardeen–Carter–Hawking:

Another reason for believing the third law is that if one could reduce κ to zero by a finite sequence of operations, then presumably one could carry the process further, thereby creating a naked singularity.

- ▶ Charging (resp., spinning) up a black hole to create a naked singularity as in superextremal RN (resp., Kerr) is known as **supercharging** (resp., **overspinning**).




- ▶ This is a doomed endeavor because the naked singularity in extremal RN/Kerr is dynamically inaccessible.
- ▶ Supercharging was definitively disproved in the EMCSF model in spherical symmetry by Kommeli '13.

The third law and supercharging

Bardeen–Carter–Hawking:

Another reason for believing the third law is that if one could reduce κ to zero by a finite sequence of operations, then presumably one could carry the process further, thereby creating a naked singularity.

- ▶ Charging (resp., spinning) up a black hole to create a naked singularity as in superextremal RN (resp., Kerr) is known as **supercharging** (resp., **overspinning**).




- ▶ This is a doomed endeavor because the naked singularity in extremal RN/Kerr is dynamically inaccessible.
- ▶ Supercharging was definitively disproved in the EMCSF model in spherical symmetry by Kommeli '13.
- ▶ Supercharging is impossible, but not because of a third law obstruction!

The third law and supercharging

Bardeen–Carter–Hawking:

Another reason for believing the third law is that if one could reduce κ to zero by a finite sequence of operations, then presumably one could carry the process further, thereby creating a naked singularity.

- ▶ Charging (resp., spinning) up a black hole to create a naked singularity as in superextremal RN (resp., Kerr) is known as **supercharging** (resp., **overspinning**)



- ▶ This is a doomed endeavor because the naked singularity in extremal RN/Kerr is dynamically inaccessible
- ▶ Supercharging was definitively disproved in the EMCSF model in spherical symmetry by Kommeli '13
- ▶ Supercharging is impossible, but not because of a third law obstruction!

Strategy of the construction

Steps in the proof:

Strategy of the construction

Steps in the proof:

1. Construct spacetime teleologically.

Strategy of the construction

Steps in the proof:

1. Construct spacetime teleologically.
 - ▶ Teleological construction allows us to locate the event horizon.

Strategy of the construction


Steps in the proof:

1. Construct spacetime teleologically.
 - ▶ Teleological construction allows us to locate the event horizon.
2. Extract a Cauchy hypersurface Σ and deduce that the teleologically constructed spacetime arises *dynamically* from initial data.

Strategy of the construction


Steps in the proof:

1. Construct spacetime teleologically.
 - ▶ Teleological construction allows us to locate the event horizon.
2. Extract a Cauchy hypersurface Σ and deduce that the teleologically constructed spacetime arises *dynamically* from initial data.




The black hole region \mathcal{BH} and event horizon \mathcal{H}^+ of a spacetime are **teleological** notions—they can't be located *without knowing the entire future of the spacetime*.


Prototype: Minkowski to Schwarzschild gluing



Prototype: Minkowski to Schwarzschild gluing




Prototype: Minkowski to Schwarzschild gluing



Characteristic gluing for the vacuum Einstein equations recently initiated in fundamental works of Aretakis–Czimek–Rodnianski and refined by Czimek–Rodnianski, but in a **perturbative** regime around Minkowski space which is inapplicable here.


Prototype: Minkowski to Schwarzschild gluing




Characteristic gluing for the vacuum Einstein equations recently initiated in fundamental works of Aretakis–Czimek–Rodnianski and refined by Czimek–Rodnianski, but in a **perturbative** regime around Minkowski space which is inapplicable here.

The present problem is fundamentally **nonlinear** and **nonperturbative**, but is tractable because of the symmetry assumption.

Characteristic/null gluing for the linear wave equation



Characteristic/null gluing for the linear wave equation




Question: Given functions

$$\phi_1 : \mathfrak{R}_1 \rightarrow \mathbb{R}, \quad \phi_2 : \mathfrak{R}_2 \rightarrow \mathbb{R}$$

solving $\square_g \phi_i = 0$,

Characteristic/null gluing for the linear wave equation



Question: Given functions

$$\phi_1 : \mathfrak{R}_1 \rightarrow \mathbb{R}, \quad \phi_2 : \mathfrak{R}_2 \rightarrow \mathbb{R}$$


solving $\square_g \phi_i = 0$, when does there exist a function

$$\phi : \mathcal{M} \rightarrow \mathbb{R}$$

solving $\square_g \phi = 0$, such that

$$\phi|_{\mathfrak{R}_i} = \phi_i ?$$

Characteristic/null gluing for the linear wave equation



Question: Given functions

$$\phi_1 : \mathfrak{R}_1 \rightarrow \mathbb{R}, \quad \phi_2 : \mathfrak{R}_2 \rightarrow \mathbb{R}$$

solving $\square_g \phi_i = 0$, when does there exist a function

$$\phi : \mathcal{M} \rightarrow \mathbb{R}$$


solving $\square_g \phi = 0$, such that

$$\phi|_{\mathfrak{R}_i} = \phi_i ?$$

Analogy:

- ▶ \mathfrak{R}_1 is Minkowski
- ▶ \mathfrak{R}_2 is Schwarzschild
- ▶ C will be the event horizon \mathcal{H}^+

Characteristic/null gluing for the linear wave equation



Question: Given functions

$$\phi_1 : \mathfrak{R}_1 \rightarrow \mathbb{R}, \quad \phi_2 : \mathfrak{R}_2 \rightarrow \mathbb{R}$$

solving $\square_g \phi_i = 0$, when does there exist a function

$$\phi : \mathcal{M} \rightarrow \mathbb{R}$$


solving $\square_g \phi = 0$, such that

$$\phi|_{\mathfrak{R}_i} = \phi_i ?$$

Analogy:

- ▶ \mathfrak{R}_1 is Minkowski
- ▶ \mathfrak{R}_2 is Schwarzschild
- ▶ C will be the event horizon \mathcal{H}^+

Characteristic/null gluing for the linear wave equation



Question: Given functions

$$\phi_1 : \mathfrak{R}_1 \rightarrow \mathbb{R}, \quad \phi_2 : \mathfrak{R}_2 \rightarrow \mathbb{R}$$

solving $\square_g \phi_i = 0$, when does there exist a function

$$\phi : \mathcal{M} \rightarrow \mathbb{R}$$

solving $\square_g \phi = 0$, such that

$$\phi|_{\mathfrak{R}_i} = \phi_i ?$$

Analogy:

- ▶ \mathfrak{R}_1 is Minkowski
- ▶ \mathfrak{R}_2 is Schwarzschild
- ▶ C will be the event horizon \mathcal{H}^+

Minkowski to Schwarzschild gluing


Theorem (Kehle–U. '22).

For any $k \in \mathbb{N}$ and $0 < R_i < 2M_f$, the Minkowski sphere of radius R_i can be characteristically glued to the Schwarzschild event horizon sphere with mass M_f to order C^k within the Einstein-scalar field model in spherical symmetry.

Minkowski to Schwarzschild gluing

Theorem (Kehle-U. '22).


For any $k \in \mathbb{N}$ and $0 < R_i < 2M_f$, the Minkowski sphere of radius R_i can be characteristically glued to the Schwarzschild event horizon sphere with mass M_f to order C^k within the Einstein-scalar field model in spherical symmetry.



Minkowski to Schwarzschild gluing


Theorem (Kehle–U. '22).

For any $k \in \mathbb{N}$ and $0 < R_i < 2M_f$, the Minkowski sphere of radius R_i can be characteristically glued to the Schwarzschild event horizon sphere with mass M_f to order C^k within the Einstein-scalar field model in spherical symmetry.




Pulse amplitudes α_j are chosen using the Borsuk–Ulam theorem, making use of the symmetry $\phi \mapsto -\phi$ of the equations.


Minkowski to Schwarzschild gluing




Minkowski to Schwarzschild gluing



Minkowski to Schwarzschild gluing



Blueprint to disproving the third law



The vacuum case

We believe that the charged scalar field is not necessary to provide a counterexample to the third law:

The vacuum case

We believe that the charged scalar field is not necessary to provide a counterexample to the third law:

Conjecture.

There exist Cauchy data for the Einstein vacuum equations

$$R_{\mu\nu} = 0$$

which undergo gravitational collapse and form an exactly Schwarzschild apparent horizon, only for the spacetime to form an exactly extremal Kerr event horizon at a later advanced time. In particular, already in vacuum, the “third law of black hole thermodynamics” is false.

The very slowly rotating case


Theorem (Kehle–U. '23).

For any mass M and specific angular momentum $0 \leq |a| \ll M$, there exists a solution of the Einstein vacuum equations with the following Penrose diagram:

The very slowly rotating case

Theorem (Kehle–U. '23).


For any mass M and specific angular momentum $0 \leq |a| \ll M$, there exists a solution of the Einstein vacuum equations with the following Penrose diagram:



The very slowly rotating case

Theorem (Kehle–U. '23).

For any mass M and specific angular momentum $0 \leq |a| \ll M$, there exists a solution of the Einstein vacuum equations with the following Penrose diagram:




This provides an alternative, and fundamentally different, proof of Christodoulou's theorem that *black holes can form due to focusing of gravitational waves alone*.

The very slowly rotating case

Theorem (Kehle–U. '23).

For any mass M and specific angular momentum $0 \leq |a| \ll M$, there exists a solution of the Einstein vacuum equations with the following Penrose diagram:



This provides an alternative, and fundamentally different, proof of Christodoulou's theorem that *black holes can form due to focusing of gravitational waves alone*.


Outlook: the “moduli space” of gravitational (non-)collapse




Outlook: the “moduli space” of gravitational (non-)collapse




Outlook: the “moduli space” of gravitational (non-)collapse



Outlook: the “moduli space” of gravitational (non-)collapse



Outlook: the “moduli space” of gravitational (non-)collapse



Critical behavior: naked singularities expected when a “low energy” black hole forms.

Critical behavior near extremality

- We expect that a generic perturbation of an extremal black hole will just make the parameters subextremal.

Critical behavior near extremality

- ▶ We expect that a generic perturbation of an extremal black hole will just make the parameters subextremal.
- ▶ Despite the presence of an instability for extremal black holes, the **Aretakis instability**, a weak form of stability is believed to be true for extremal black holes. (See the work of Angelopoulos–Aretakis–Gajic on nonlinear model equations.)

Critical behavior near extremality

- ▶ We expect that a generic perturbation of an extremal black hole will just make the parameters subextremal.
- ▶ Despite the presence of an instability for extremal black holes, the **Aretakis instability**, a weak form of stability is believed to be true for extremal black holes. (See the work of Angelopoulos–Aretakis–Gajic on nonlinear model equations.)

Conjecture (Dafermos–Holzegel–Rodnianski–Taylor).

The stable manifold of very slowly rotating extremal Kerr–Newman delimits the boundary between asymptotically subextremal black holes and spacetimes failing to collapse for a while longer.

Critical behavior near extremality


- ▶ We expect that a generic perturbation of an extremal black hole will just make the parameters subextremal.
- ▶ Despite the presence of an instability for extremal black holes, the **Aretakis instability**, a weak form of stability is believed to be true for extremal black holes. (See the work of Angelopoulos–Aretakis–Gajic on nonlinear model equations.)

Conjecture (Dafermos–Holzegel–Rodnianski–Taylor).


The stable manifold of very slowly rotating extremal Kerr–Newman delimits the boundary between asymptotically subextremal black holes and spacetimes failing to collapse for a while longer.

Critical behavior: extremal black holes can be perturbed so black hole formation is delayed and the event horizon jumps just as the extremal threshold is crossed.


Event horizon jumping




Event horizon jumping




Event horizon jumping




Event horizon jumping



So what does happen if you “go past extremality”?



So what does happen if you “go past extremality”?



Critical behavior: We expect that there are smooth 1-parameter families of solutions passing through extremality where the event horizon “jumps” as soon as the exterior becomes superextremal. There is no naked singularity.

Instability of black hole formation due to electromagnetic repulsion

Conjecture.

There exist regular Cauchy data for the Einstein–Maxwell-charged scalar field system, leading to the formation of an extremal Reissner–Nordström black hole, that can be perturbed to disperse, i.e. no black hole forms.

Instability of black hole formation due to electromagnetic repulsion

Conjecture.

There exist regular Cauchy data for the Einstein–Maxwell-charged scalar field system, leading to the formation of an extremal Reissner–Nordström black hole, that can be perturbed to disperse, i.e. no black hole forms.

Critical behavior: the very black-holeness of certain spacetimes is unstable.

- ▶ This conjecture seems very difficult because dispersion is confounded by the slow decay of the charged scalar field.

Instability of black hole formation due to electromagnetic repulsion

Conjecture.

There exist regular Cauchy data for the Einstein–Maxwell-charged scalar field system, leading to the formation of an extremal Reissner–Nordström black hole, that can be perturbed to disperse, i.e. no black hole forms.

Critical behavior: the very black-holeness of certain spacetimes is unstable.

- ▶ This conjecture seems very difficult because dispersion is confounded by the slow decay of the charged scalar field. (The scalar field part of the system is already scaling critical.)

Instability of black hole formation due to electromagnetic repulsion

Conjecture.

There exist regular Cauchy data for the Einstein–Maxwell-charged scalar field system, leading to the formation of an extremal Reissner–Nordström black hole, that can be perturbed to disperse, i.e. no black hole forms.

Critical behavior: the very black-holeness of certain spacetimes is unstable.

- ▶ This conjecture seems very difficult because dispersion is confounded by the slow decay of the charged scalar field. (The scalar field part of the system is already scaling critical.)
- ▶ We propose to first study this phenomenon in the Einstein–Maxwell-massless Vlasov model, inspired by classical work of Ori and recent work of Moschidis.

Instability of black hole formation due to electromagnetic repulsion

Conjecture.

There exist regular Cauchy data for the Einstein–Maxwell-charged scalar field system, leading to the formation of an extremal Reissner–Nordström black hole, that can be perturbed to disperse, i.e. no black hole forms.

Critical behavior: the very black-holeness of certain spacetimes is unstable.

- ▶ This conjecture seems very difficult because dispersion is confounded by the slow decay of the charged scalar field. (The scalar field part of the system is already scaling critical.)
- ▶ We propose to first study this phenomenon in the Einstein–Maxwell-massless Vlasov model, inspired by classical work of Ori and recent work of Moschidis.
- ▶ (g, F, f)

$$G_{\mu\nu}[g] = 2 \left(T_{\mu\nu}^{\text{EM}}[F] + T_{\mu\nu}^{\text{Vl}}[f] \right) \quad (6)$$

$$\nabla^\nu F_{\mu\nu} = \epsilon N_\mu[f] \quad (7)$$

$$\left(p^\mu \frac{\partial}{\partial x^\mu} - \Gamma^\mu{}_{\alpha\beta} p^\alpha p^\beta \frac{\partial}{\partial p^\mu} + \epsilon F^\mu{}_\nu p^\nu \frac{\partial}{\partial p^\mu} \right) f = 0 \quad (8)$$

$$\text{spt } f \subset \{p^2 = 0\} \quad (9)$$

Injection of charged massless Vlasov beam into Minkowski space

Conjecture.


There exist smooth one-parameter families of solutions $(g_\lambda, F_\lambda, f_\lambda)_{\lambda \in (-\varepsilon, \varepsilon)}$ to the Einstein–Maxwell-massless Vlasov system with the following behavior:

Injection of charged massless Vlasov beam into Minkowski space

Conjecture.

There exist smooth one-parameter families of solutions $(g_\lambda, F_\lambda, f_\lambda)_{\lambda \in (-\varepsilon, \varepsilon)}$ to the Einstein–Maxwell-massless Vlasov system with the following behavior:

$\lambda > 0$: subextremal BH forms




Injection of charged massless Vlasov beam into Minkowski space


Conjecture.

There exist smooth one-parameter families of solutions $(g_\lambda, F_\lambda, f_\lambda)_{\lambda \in (-\varepsilon, \varepsilon)}$ to the Einstein–Maxwell-massless Vlasov system with the following behavior:

$\lambda > 0$: subextremal BH forms



$\lambda = 0$: extremal BH forms




Injection of a charged massless Vlasov beam into Minkowski space


Conjecture.

There exist smooth one-parameter families of solutions $(g_\lambda, F_\lambda, f_\lambda)_{\lambda \in (-\varepsilon, \varepsilon)}$ to the Einstein–Maxwell-massless Vlasov system with the following behavior:


$\lambda > 0$: subextremal BH forms



$\lambda = 0$: extremal BH forms



$\lambda = 0$: no BH forms



Conclusions

Conclusions

- Subextremal black holes can become extremal in finite time, so the third law of black hole thermodynamics is false.
-

Conclusions

- ▶ Subextremal black holes can become extremal in finite time, so the third law of black hole thermodynamics is false.
- ▶ A $T = 0$ black hole is no more difficult to form than a fixed $T > 0$ black hole.

Conclusions


- ▶ Subextremal black holes can become extremal in finite time, so the third law of black hole thermodynamics is false.
- ▶ A $T = 0$ black hole is no more difficult to form than a fixed $T > 0$ black hole.
- ▶ The limit $T \rightarrow 0$ is continuous. Don't be confused by the naked singularity of ERN—not every part of the maximally extended RN solution is physically relevant in this limit.

Conclusions

- ▶ Subextremal black holes can become extremal in finite time, so the third law of black hole thermodynamics is false.
- ▶ A $T = 0$ black hole is no more difficult to form than a fixed $T > 0$ black hole.
- ▶ The limit $T \rightarrow 0$ is continuous. Don't be confused by the naked singularity of ERN—not every part of the maximally extended RN solution is physically relevant in this limit.
- ▶ However, there is critical behavior lurking near the formation of $T = 0$ black holes. We hope that characteristic gluing and PDE techniques can be used to study this phenomenon.

Conclusions

- Subextremal black holes can become extremal in finite time, so the third law of black hole thermodynamics is false.
- A $T = 0$ black hole is no more difficult to form than a fixed $T > 0$ black hole.
- The limit $T \rightarrow 0$ is continuous. Don't be confused by the naked singularity of ERN—not every part of the maximally extended RN solution is physically relevant in this limit.
- However, there is critical behavior lurking near the formation of $T = 0$ black holes. We hope that characteristic gluing and PDE techniques can be used to study this phenomenon.



Thank you!