

Bioinformatic approaches to regulatory genomics and epigenomics

376-1347-00L | week 07

Pierre-Luc Germain

Today's plan

(update epiwraps, download data)

- Debriefing on the assignment
- DNA accessibility
- ATAC-seq analysis (practical)
- Nucleosome positioning

Debriefing on the assignments

Searching for matches: choose the subject carefully !

Find the percentage of motifs across the genome within a peak

```
``{r}  
# when subject peak centers, motifs are only (!) searched within the peak centers  
moi_peaks <- matchMotifs(motif, subject_peak_centers, genome=genome, out="positions")  
sum(overlapsAny(moi_peaks, peak_centers))/length(moi_peaks)  
``
```

```
[1] 1
```

Debriefing on the assignments

Searching for matches: choose the subject carefully !

Find the percentage of motifs across the genome within a peak

```
```{r}
when subject peak centers, motifs are only (!) searched within the peak centers
moi_peaks <- matchMotifs(motif, subject=peak_centers, genome=genome, out="positions")
sum(overlapsAny(moi_peaks, peak_centers))/length(moi_peaks)
```
```

[1] 1

correct:

Find the fraction of motifs across the genome within a peak

```
```{r}
ah <- AnnotationHub(localHub=FALSE)
genome <- ah[["AH68356"]]
genome_seq <- rtracklayer::import(genome)
genome_coords <- GRanges(seqnames=Rle(c("chr1")),
 ranges = IRanges(1, end=200*1e6)) # here for one chromosome

with findMotifInstances (epiwraps)
moi_genome <- findMotifInstances(genome_seq, motif) #epiwraps
|
With matchMotifs (motifmatchr)
moi_genome2 <- matchMotifs(motif2, subject=genome_coords, genome=BSgenome.Mmusculus.UCSC.mm10, out="positions")
```

```{r, get overlaps}
sum(overlapsAny(moi_genome, peak_centers))/length(moi_genome)
```
```

Of all instances of that motif in the genome, what proportion is bound by the factor (i.e. has a peak)?

```
mmusculus <- import(genome, "2bit", which = as(seqinfo(genome), "GenomicRanges"))  
motif_instances_genome <- findMotifInstances(mmusculus, motif, mc.cores=2)
```

```
## Note: motif [motif] has an empty nsites slot, using 100.
```

```
length(motif_instances_genome)
```

```
## [1] 6544422
```

```
motif_with_peaks = overlapsAny(motif_instances_genome, peaks)  
sum(motif_with_peaks)
```

```
## [1] 16856
```

```
percentage2 <- sum(motif_with_peaks)/length(motif_instances_genome)*100  
percentage2
```

```
## [1] 0.2575629
```

Of the 6544422 motif instances, 16856 (0.2575629%) overlap a peak.

TF:
GATA1

Of the 9675
GATA1 peaks,
7277 (~75%)
contain a
GATA1 motif,
but...



Debriefing on the assignment – wrapping up

- The proportion of binding sites that show the motif depends on the TF (ranging from roughly 20 to 95%)
- The proportion of (genome-wide) motif instances that are bound by the factor is typically very very small
- This means that something else determines whether a genomic region will be bound

DNA accessibility, which is associated to lower nucleosome density, reflects activity




Fig. 1 | **A continuum of accessibility states broadly reflects the distribution of chromatin dynamics across the genome.** In contrast to closed chromatin, permissive chromatin is sufficiently dynamic for transcription factors to initiate sequence-specific accessibility remodelling and establish an open chromatin conformation (illustrated here for an active gene locus). Pol II, RNA polymerase II; TF, transcription factor.

(Klemm, Shipony and Greenleaf, 2019)

Returning to our very brief history of genetics & genomics

...

1900 - Rediscovery of Mendel's work (1860s)

1913 - Chromosomes are linear arrays of genes

1941 - the one-gene-one-enzyme hypothesis

1944 - DNA is the genetic material

1951 - First protein sequenced

1977 - DNA sequencing

1977 - Eukaryotic genes are spliced

1995 - First bacterial genomes sequenced

2000 - Next Generation Sequencing (NGS)

2001 - Draft of the human genome

2003 - RNA-seq


2006 - ChIP-seq

2008 - DNase-seq, MNase-seq


2012 - ATAC-seq

} Accessibility
assays

...





ATAC-seq



ATAC-seq recently became extremely popular due to its information content and low material requirement (i.e. # cells)

Chromatin accessibility assays







ATAC signal tells us that something is happening, it just doesn't tell us what exactly...

(Adapted from Fox et al., Nat Comm 2020)


ATACseq



ATACseq




This means that once we have the data, we can split the fragments according to size in order to obtain specific information about different kinds of chromatin signals





Practical

Estimating TF activity from accessibility and footprints



(Baek, Goldstein and Hager, Cell Reports 2017)

Estimating TF activity from accessibility and footprints




(Baek, Goldstein and Hager, Cell Reports 2017)

“Shifting” ATAC-seq alignments


From a given ATAC-seq insertion site, the exact region that is accessible is a few nucleotides from the start of the read

When doing high-resolution things like footprinting, one therefore typically shifts the cut sites by +4/-5nt, so that it is placed in the middle of where the Tn5 was binding



(For most other purposes, this is too fine-grained to make a difference)




(adapted from
Zhijian et al.,
Genome Biology 2019)



Nucleosome positioning





(Klemm, Shipony and Greenleaf, 2019)

Assignment

In the same dataset of ATAC on chr19, plot the insertion (i.e. 'cuts') profile of, respectively, nucleosome-free and nucleosome-containing fragments, around the high-confidence motifs of two factors.

You can choose your own factors of interest, or for instance use GATA1, REST and/or the glucocorticoid receptor (search "GCR")

Expected form of the answer: 2 figures (one for each factor), each containing the two signals around the motifs

Don't forget to render your markdown and push it as [assignment.html](#) !