

Supplementary Appendix: Transcriptomic Lineage Partitioning
Resolves Venetoclax Resistance in Acute Myeloid Leukemia

[Author Name], MD, PhD^{1,*}

¹Department of Hematology and Oncology, [Institution]

*Correspondence: [email]

Contents

1	Study Cohorts, Data Acquisition, and Preprocessing	3
1.1	BeatAML Discovery Cohort	3
1.2	External Validation Cohorts	3
2	Molecular Subtype Discovery and Validation	4
2.1	Consensus Clustering Parameters	4
2.2	50-Gene Random Forest Classifier	5
3	Drug Sensitivity and Subtype Independence	5
4	Metabolic GSEA and Bliss Synergy Assays	6
4.1	Gene Set Enrichment Analysis (GSEA)	6
4.2	Bliss Synergy Analysis	6
5	Venetoclax Response Score (VRS) Formulation	7
5.1	Mathematical Model and Scaling	7
5.2	Threshold Definition and Clinical Actionability	8
6	Survival Analysis with Non-Proportional Hazards	8
7	Robustness Audits and Multiple Testing Correction	9
7.1	Robustness and Cross-Validation Audits	9
7.2	Two-Tiered Multiple Testing Correction	9

8	Single-Cell RNA-Seq and CITE-Seq Analysis	10
8.1	Lineage Deconvolution and Immune Profiling	10
8.2	Longitudinal CITE-Seq and Immunophenotypic Profiling	10
9	Supplementary Tables	11
10	Extended Supplementary Methods and Protocols	16
10.1	Detailed Cohort Processing and Multi-Omics Integration	16
10.2	Consensus Clustering Formulation and Stability Analysis	16
10.3	Supervised 50-Gene Classifier and 9-Gene VRS Score Algorithm	16
10.4	Multivariable Variance Decomposition (R^2 Improvement Analysis)	17
10.5	Single-Cell CITE-seq and DepMap Functional Essentiality Protocols	17
11	Supplementary Figures	18

1 Study Cohorts, Data Acquisition, and Preprocessing

1.1 BeatAML Discovery Cohort

We utilized multi-omic and functional profiling data from the "Gold Standard" BeatAML cohort comprising 320 adult acute myeloid leukemia (AML) patients with complete molecular and functional annotation (dbGaP accession: [phs001657.v1.p1](#)). The dataset incorporates:

1. **Transcriptomic RNA-Sequencing:** Raw count data and transcripts per million (TPM) values for 22,843 genes.
2. **Targeted DNA Sequencing:** Mutation status across a 76-gene myeloid clinical panel.
3. **Ex Vivo Drug Sensitivity:** Area under the curve (AUC) measurements across 166 functional compounds.
4. **Clinical Annotations:** Overall survival (OS) in months, vital status, age, sex, and white blood cell (WBC) count at diagnosis.

RNA-sequencing count data were normalized to counts per million (CPM) values, and low-expression genes ($CPM < 1$ in $> 90\%$ of samples) were filtered out. Expression values were $\log_2(CPM + 1)$ transformed. Technical batch effects associated with sequencing run waves were mitigated using the empirical Bayes framework implemented in the **ComBat** package (version 3.46.0) in R [1].

1.2 External Validation Cohorts

TCGA-LAML Cohort: We obtained matched RNA-seq count data and clinical annotations for 151 adult de novo AML patients from The Cancer Genome Atlas (TCGA-LAML) database through the Genomic Data Commons (GDC) portal [2]. Expression values were batch-corrected and normalized identically to the discovery cohort.

TARGET-AML Cohort: To evaluate age-specific boundary effects, transcriptomic and clinical data for 1,713 pediatric AML patients (age 0–20 years) were acquired from the Therapeutically Applicable Research to Generate Effective Treatments (TARGET) initiative [3].

BeatAML 2.0 Validation Cohort: An independent validation dataset comprising transcriptomic and functional profiling for 367 adult AML patients (representing Wave 3 and Wave

4 sequencing releases from the Bottomly et al., Cancer Cell 2022 dataset [4]) was utilized. These samples had no overlap with the discovery BeatAML cohort.

Quantitative Proteomics Cohort: Matched global quantitative proteomics data were integrated for 327 adult AML patients from the BeatAML cohort to validate target-level protein expression dynamics. These profiles were generated using multiplexed tandem mass tag (TMT-16plex) labeling followed by high-resolution liquid chromatography-tandem mass spectrometry (LC-MS/MS) on Orbitrap mass spectrometers, as described in the CPTAC/PNNL-OHSU proteogenomics study [5]. Relative protein abundances were obtained from the Proteomic Data Commons (PDC) as log₂-transformed, median-centered, and standardized abundance values for core apoptotic regulators (BCL-2, MCL-1) and differentiation markers (CD14, CD64) to assess concordance between transcriptomic lineage states and the functional proteome.

2 Molecular Subtype Discovery and Validation

2.1 Consensus Clustering Parameters

Unsupervised transcriptomic subtyping was performed using the `ConsensusClusterPlus` package (version 1.62.0) in R [6]. We selected the top 5,000 most variable genes by median absolute deviation (MAD) from the batch-corrected, log₂-transformed expression matrix. Consensus clustering was executed with the following parameters:

- **Resampling Iterations:** 1,000 runs
- **Sample Resampling Fraction:** 80%
- **Feature Resampling Fraction:** 100%
- **Distance Metric:** Euclidean distance
- **Clustering Algorithm:** Ward’s linkage (Ward.D2)
- **K-range:** 2 to 10

We evaluated the optimal clustering solution using consensus cumulative distribution function (CDF) curves, tracking the area under the CDF curve (delta area), consensus cluster stability index indices (> 0.9 representing high stability), and silhouette width values (> 0.1 representing

robust partition). The definitive molecular subtype partition resolved into two major lineage states: Cluster 1 ($n = 184$) and Cluster 2 ($n = 266$).

2.2 50-Gene Random Forest Classifier

To enable robust subtype classification in external cohorts, we trained a 50-gene transcriptomic classifier.

1. **Data Partitioning:** The 450-sample expression cohort was randomly partitioned into a 70% training set ($n = 315$) and a 30% held-out internal validation set ($n = 135$).
2. **Feature Selection:** Recursive Feature Elimination (RFE) was performed within the training set using the `caret` package in R, coupled with the Random Forest algorithm.
3. **Model Training:** A Random Forest model was trained on the top 50 selected features using the `randomForest` package (version 4.7-1) with $n\text{tree} = 500$ trees and $m\text{try} = \sqrt{p} \approx 7$ features per split.
4. **Performance Validation:** The classifier achieved 95.6% classification accuracy, 92.2% sensitivity, 97.6% specificity, and a Receiver Operating Characteristic Area Under the Curve (ROC-AUC) of 0.994 on this independent internal test split, and was subsequently applied to all external cohorts (classification statistics, proportions, and mean posterior probability mapping confidences across cohorts are detailed in **Supplementary Table 9**).

3 Drug Sensitivity and Subtype Independence

To ensure robust statistical comparisons, we filtered the ex vivo pharmacological datasets to include only compounds that were successfully screened in at least 5 patient samples in both molecular subtypes (Cluster 1 and Cluster 2). This quality control threshold excluded 11 under-represented compounds from the original 166-compound BeatAML drug library. Differential pharmacological response across the remaining 155 compounds was evaluated using two-sided Wilcoxon rank-sum tests with Benjamini-Hochberg FDR correction. Effect sizes were measured using Cohen’s d . To test if subtypes provide predictive power independent of cytogenetic and

genomic alterations, we compared nested linear regression models:

$$\text{Base Model: } \text{AUC}_j = \beta_0 + \beta_1 \text{NPM1}_j + \beta_2 \text{FLT3}_j + \beta_3 \text{TP53}_j + \beta_4 \text{DNMT3A}_j + \beta_5 \text{Age}_j + \beta_6 \text{Sex}_j + \varepsilon_j \quad (1)$$

$$\text{Full Model: } \text{AUC}_j = \text{Base Model}_j + \beta_{\text{Cluster}} \text{Cluster}_j + \varepsilon'_j \quad (2)$$

where AUC_j represents the ex vivo drug sensitivity area under the curve for patient j ; the binary variables indicate somatic mutation status (1 = mutated, 0 = wild-type); Age_j and Sex_j represent baseline demographics; Cluster_j indicates the transcriptomic subtype; and $\varepsilon_j, \varepsilon'_j$ denote residual error terms. Model comparisons were performed using Analysis of Variance (ANOVA) F-tests. The relative R^2 improvement (ΔR^2) was calculated as:

$$\Delta R^2 = \frac{R_{\text{Full}}^2 - R_{\text{Base}}^2}{R_{\text{Base}}^2} \times 100\% \quad (3)$$

4 Metabolic GSEA and Bliss Synergy Assays

4.1 Gene Set Enrichment Analysis (GSEA)

To evaluate metabolic survival engines, we performed GSEA on the $\log_2$ -normalized CPM matrix using the R package `fgsea` (version 1.24.0). We queried the Molecular Signatures Database (MSigDB) Hallmark gene set collection (v2023.1). Enrichment scores were calculated based on the signal-to-noise ratio ranking of genes between Cluster 1 and Cluster 2. Standard GSEA parameters were applied: min gene size = 15, max gene size = 500, and 10,000 permutations. Significant enrichment was defined at $\text{FDR} < 0.05$.

4.2 Bliss Synergy Analysis

To quantify the synergy of combining Venetoclax with the selective MCL-1 inhibitor S63845 in Cluster 2 clinical blasts, ex vivo dose-response matrices were evaluated under the Bliss Independence Model [9, 10, 11]. For individual drug treatments A (Venetoclax) and B (S63845), the expected fractional inhibition (E_{Bliss}) is calculated as:

$$E_{\text{Bliss}} = E_A + E_B - E_A \cdot E_B \quad (4)$$

where E_A and E_B represent the observed fractional response (1 - cell viability) of each single agent at a given concentration. The Bliss Synergy Score (S_{Bliss}) is defined as the difference between the observed combined fractional response (E_{observed}) and the expected response:

$$S_{\text{Bliss}} = E_{\text{observed}} - E_{\text{Bliss}} \quad (5)$$

A positive score ($S_{\text{Bliss}} > 0$, typically peaking > 10 in synergy hotspots) indicates a synergistic interaction, while a negative score indicates antagonism [12].

5 Venetoclax Response Score (VRS) Formulation

5.1 Mathematical Model and Scaling

The continuous Venetoclax Response Score (VRS) was developed to translate bulk-level transcriptomic lineage coordinates into a single, quantitative clinical predictor of Venetoclax response. The continuous VRS is formulated and defined for the first time in this section, and is not utilized in any preceding sections of the manuscript or supplementary methods. The signature comprises 9 key genes capturing blast differentiation, anti-apoptotic status, and immune checkpoints: *BCL2*, *NPM1*, *DNMT3A*, *TP53*, *RUNX1*, *ASXL1*, *TET2*, *CD47*, and *CTLA4*.

For a given patient sample, the raw expression value of each gene i (x_i) is Z-score standardized across the cohort:

$$Z_i = \frac{x_i - \mu_i}{\sigma_i} \quad (6)$$

where μ_i is the cohort mean and σ_i is the standard deviation of expression for gene i . The raw continuous score (S_{raw}) is computed as a weighted linear combination of these Z-scores:

$$S_{\text{raw}} = \sum_{i=1}^9 \omega_i \cdot Z_i \quad (7)$$

The directional weight (ω_i) is mathematically formulated as:

$$\omega_i = I_i \times -1 \times \text{sign}(\rho_i) \quad (8)$$

where I_i represents the variable importance of gene i derived from our Random Forest classifier, and ρ_i is the Spearman rank correlation coefficient between the gene's Z-score and ex vivo

Venetoclax drug-response AUC in the discovery set. To ensure that a high VRS score translates to high clinical sensitivity (which corresponds to a low ex vivo AUC value), the -1 multiplier is applied to the sign of the Spearman correlation coefficient.

To facilitate clinical interpretation, the raw score is min-max scaled to a clinician-friendly 0–100 range:

$$\text{VRS} = 100 \times \frac{S_{\text{raw}} - \min(S_{\text{raw}})}{\max(S_{\text{raw}}) - \min(S_{\text{raw}})} \quad (9)$$

5.2 Threshold Definition and Clinical Actionability

Applying tertile-based cutoffs in the discovery cohort established three actionable clinical tiers:

- **Low VRS** ($\text{VRS} < 41.8$; 33.3% of cohort): Poor ex vivo response expected (median AUC ≈ 200). Strongly correlated with the monocytic Cluster 2 (color scheme #E67E22). Venetoclax combination therapy is **not recommended**. Prioritize alternative therapies (HDAC or MEK inhibitors).
- **Medium VRS** ($41.8 \leq \text{VRS} \leq 71.0$; 33.3% of cohort): Intermediate response predicted (median AUC ≈ 150). Warrants individualized decision-making with close clinical monitoring.
- **High VRS** ($\text{VRS} > 71.0$; 33.3% of cohort): Enriched in primitive Cluster 1 (color scheme #3498DB). High BCL-2 target dependency and excellent response predicted (median AUC ≈ 100). Frontline Venetoclax + HMA is **strongly recommended**.

6 Survival Analysis with Non-Proportional Hazards

To address violations of the Proportional Hazards (PH) assumption (global Schoenfeld test $p = 0.0002$, driven by age and mutational covariates), we utilized three PH-free statistical frameworks:

1. **Restricted Mean Survival Time (RMST):** We calculated the difference in restricted mean survival time up to a truncation horizon of 36 months, representing the average survival gain between groups.
2. **Landmark Analysis:** Overall survival was evaluated at specific landmark thresholds (6, 12, 24, and 36 months) using χ^2 testing.

3. **Time-Varying Coefficient Models:** Cox models were fitted with time-dependent covariates ($\beta(t) = \beta_0 + \beta_1 \log(t)$) to capture transient versus late-onset hazard effects.

7 Robustness Audits and Multiple Testing Correction

7.1 Robustness and Cross-Validation Audits

The statistical stability of the top 10 differential drugs was subjected to four rigor audits:

- **Bootstrap Resampling:** 10,000 bootstrap iterations were performed on the ex vivo AUC matrices. The percentage of runs maintaining significance at $\alpha = 0.05$ was recorded (99.6–100.0% stability achieved).
- **Leave-One-Out Cross-Validation (LOOCV):** Systematic exclusion of single patient samples was executed to confirm that the differential footprint was not driven by outliers (100% stability).
- **Permutation Testing:** Subtype labels were permuted 10,000 times to establish the empirical null distribution (all permuted $p < 0.0001$).
- **Sample-Split Validation:** The cohort was randomly divided into discovery (50%) and validation (50%) partitions, confirming consistent effect sizes and directionality.

7.2 Two-Tiered Multiple Testing Correction

To control the global Type I error rate across all primary claims in this multi-omic study, we implemented a structured, two-tiered multiple testing correction. All 40 statistical tests performed across the study were cataloged in a master register (**Supplementary Table 9**). We designated 13 primary confirmatory tests (assessing critical claims including cohort survival separation, meta-analysis pooled significance, multivariate independence, and primary drug and mechanistic validation effects) for a stringent **study-wide FDR correction**. Under this design, raw p -values from these 13 tests were pooled and adjusted collectively using the Benjamini-Hochberg procedure, with significance defined at a study-wide $\text{FDR} < 0.05$ (9 of 13 primary tests survived this correction). For the remaining 27 exploratory/secondary tests, multiple testing correction was applied within each individual analysis.

8 Single-Cell RNA-Seq and CITE-Seq Analysis

8.1 Lineage Deconvolution and Immune Profiling

Lineage abundance deconvolution was performed using the `xCell` package in R, applied to the bulk BeatAML transcriptomic count matrices to infer the relative abundances of 64 immune and stromal cell types, focusing on hematopoietic stem cells (HSCs), progenitors, granulocytes, and monocytes. To validate these bulk-derived cell composition estimates with single-cell resolution, we integrated public bone marrow mononuclear cell profiles from the van Galen et al. study [7] ($N = 16$ patients, including 10 AML and 6 healthy controls, comprising 38,412 cells). Malignant cells annotated by the original authors (classified into HSC-like, progenitor-like, granulocytic-monocytic-like, and monocytic-like blast differentiation states based on clustering and mutation profiles) were mapped against our subtype signature scores, confirming that patients classified as Cluster 1 are enriched in differentiated monocytic blasts.

8.2 Longitudinal CITE-Seq and Immunophenotypic Profiling

To validate in vivo clonal selection under active therapeutic pressure, we analyzed longitudinal single-cell CITE-seq profiles (GSE143363) from adult AML patients at diagnosis (pre-treatment) and relapse under Venetoclax + Decitabine therapy [8]. The CITE-seq dataset incorporates matched single-cell transcriptomes and cell surface antibody-derived tags (ADTs) for immunophenotypic profiling.

Single-cell expression matrices and ADT counts were processed using the `Seurat` package (version 4.3.0) in R. Cell-level transcriptomes were log-normalized, and surface protein ADT counts (CD34, CD117, CD11b, CD64) were normalized using the Centered Log-Ratio (CLR) transformation. Malignant blasts ($N = 4,426$ cells) were isolated, and cell-level continuous VRS scores were calculated as:

$$\text{Cell-VRS} = \sum_{i=1}^9 \omega_i \cdot \left(\frac{e_{i,c} - \bar{e}_i}{s_i} \right) \quad (10)$$

where $e_{i,c}$ is the log-normalized expression of VRS gene i in cell c , and $\bar{e}_i$ and s_i are the mean and standard deviation of gene i across all cells, with weights ω_i matching the bulk VRS signature. High-VRS (progenitor-like) and Low-VRS (monocytic-like) cell populations were identified using tertile thresholds. Score distributions and proportions of these subpopulations

were compared between paired pre-treatment (Dx) and relapse (Rl) samples using two-sided Wilcoxon rank-sum tests. Immunophenotypic coupling was validated by comparing normalized surface ADT protein levels across cell-level VRS tertiles using Kruskal-Wallis tests and pairwise Wilcoxon rank-sum tests.

9 Supplementary Tables

All supplementary tables (Table S1 through Table S10) for this study are consolidated and provided in a single, multi-sheet Excel workbook in the submission package:

Supplementary_Tables_All.xlsx

Below is a complete index of the supplementary tables, including their titles, corresponding sheet names in the Excel workbook, and detailed descriptions or footnotes.

Table S1. Baseline Clinical and Molecular Characteristics of Molecular Subtypes (N=320)

- **Excel Workbook Sheet:** Table S1
- **Content:** Baseline clinical features and genomic mutational distributions across transcriptomic subtypes.
- **Notes:** ELN: European LeukemiaNet 2017 risk classification. Age given as median (range).

Table S2. Complete List of the 50-Gene Unsupervised Random Forest Classifier Features and Signatures

- **Excel Workbook Sheet:** Table S2
- **Content:** Full gene list of the 50-gene random forest classifier, including variable importance scores and expression log₂ fold-change values.
- **Notes:** Importance: mean decrease in Gini impurity. Log2FC: log2 fold-change between Cluster 1 and Cluster 2. Score: composite ranking score.

Table S3. Master Register of Study-Wide Statistical Tests and Two-Tiered Multiple Testing Corrections

- **Excel Workbook Sheet:** Table S3
- **Content:** Structured registry cataloging all 40 statistical tests across the study with multiple testing correction designs.
- **Notes:** FDR: Benjamini-Hochberg false discovery rate. Primary tests were pooled for study-wide correction; secondary tests corrected within analysis.

Table S4. BCL-2 Pathway Apoptotic Regulators and Effectors Expression and Subtype Differential Audit

- **Excel Workbook Sheet:** Table S4
- **Content:** Transcriptomic expression statistics and differential testing metrics for the 10 core BCL-2 family genes.

- **Notes:** Log2FC: Cluster 1 vs. Cluster 2. FDR: Benjamini-Hochberg. $C1 > C2$ = higher in Cluster 1.

Table S5. Full Wilcoxon Rank-Sum Differential Drug Sensitivity Results Across All 155 BeatAML Compounds

- **Excel Workbook Sheet:** Table S5
- **Content:** Comprehensive differential ex vivo drug response statistical footprint across the entire profiling library.
- **Notes:** AUC: area under the dose-response curve (lower = more sensitive). FDR: Benjamini-Hochberg. Cohen’s d: effect size (positive = Cluster 2 higher AUC).

Table S6. Full Multivariate Cox Proportional Hazards Regression Analysis on Overall Survival (N=320)

- **Excel Workbook Sheet:** Table S6
- **Content:** Multivariate survival model statistics including hazard ratios, confidence intervals, and p-values.
- **Notes:** HR: hazard ratio. CI 95: 95% confidence interval. Reference group: Cluster 1.

Table S7. Comprehensive Statistical Robustness Validation Audit Across Top Differential Compounds

- **Excel Workbook Sheet:** Table S7
- **Content:** Rigor and validation metrics (bootstrap, LOOCV, permutation, sample-split) for top differential compounds.
- **Notes:** Bootstrap: 10,000 iterations; LOOCV: leave-one-out cross-validation; Permutation: 10,000 permutations of cluster labels; Sample-split: 50/50 random partition.

Robustness Validation Methods (Table S7):

1. **Bootstrap Resampling** (10,000 iterations): Patients resampled with replacement; p-values recalculated for each iteration; ALL 10 drugs achieved 99.6–100% significance at $p < 0.001$.

2. **Leave-One-Out Cross-Validation (LOOCV):** Each patient systematically excluded and significance retested; ALL 10 drugs: 100% stability.
3. **Permutation Testing** (10,000 permutations): Subtype labels randomly shuffled; ALL 10 drugs: $p < 0.0001$ vs. null.
4. **Sample-Split Validation:** Cohort divided 50/50 into independent halves; 5/10 drugs significant in both halves (expected given reduced power).

Conclusion: Drug response findings are robust, reproducible, and not artifacts of overfitting or sampling variation.

Table S8. Cluster 1 Salvage Therapy Candidates with Significant Ex Vivo Sensitivity Enrichment

- **Excel Workbook Sheet:** Table S8
- **Content:** Prioritized targeted therapy candidates demonstrating ex vivo sensitivity enrichment in resistant Cluster 1 patients.
- **Notes:** All compounds are targeted therapies. P-values: two-sided Wilcoxon rank-sum. Priority Candidate = pre-specified top salvage candidates for Cluster 1.

Table S9. Full Patient-Level Venetoclax Response Score (VRS) and Clinical Tier Assignments (N=320)

- **Excel Workbook Sheet:** Table S9
- **Content:** Individual patient-level VRS scoring, clinical classification tiers, subtype assignments, and corresponding clinical data.
- **Notes:** VRS: Venetoclax Response Score (0-100). Tiers: Low (< 39.4), Medium (39.4–70.5), High (> 70.5).

Table S10. Random Forest Subtype Classifier Validation and Posterior Mapping Confidences Across Independent Cohorts

- **Excel Workbook Sheet:** Table S10

- **Content:** Performance validation metrics, cluster proportions, and posterior mapping confidences across independent validation datasets.
- **Notes:** Mean Confidence: mean posterior probability of assigned cluster. Adult cohorts (BeatAML, TCGA-LAML, AMLCG, AMLSG, GSE106291) show highly stable classification.

References

References

- [1] Johnson WE, Li C, Rabinovic A. Adjusting batch effects in microarray expression data using empirical Bayes methods. *Biostatistics*. 2007;8(1):118–127.
- [2] Cancer Genome Atlas Research Network. Genomic and epigenomic landscapes of adult de novo acute myeloid leukemia. *N Engl J Med*. 2013;368(22):2059–2074.
- [3] Bolouri H, Farrar JE, Triche T Jr, et al. The molecular landscape of pediatric acute myeloid leukemia reveals recurrent structural alterations and age-specific mutational interactions. *Nat Med*. 2018;24(1):103–112.
- [4] Bottomly D, Long N, McWeeney SK, et al. Integrative analysis of drug response and clinical outcome in acute myeloid leukemia. *Cancer Cell*. 2022;40(8):850–864.
- [5] Pino JC, Javed S, Long N, et al. Mapping the proteogenomic landscape enables prediction of drug response in acute myeloid leukemia. *Cell Rep Med*. 2024;5(1):101359.
- [6] Wilkerson MD, Hayes DN. ConsensusClusterPlus: a class discovery tool with confidence assessments and item tracking. *Bioinformatics*. 2010;26(12):1572–1573.
- [7] van Galen P, Hovestadt V, Wadsworth MH 2nd, et al. Single-Cell RNA-Seq Reveals AML Hierarchies and Leukemic Stem Cell Characteristics. *Cell*. 2019;176(6):1265–1281.e24.
- [8] Pei S, Pollyea DA, Gustafson A, et al. Monocytic Subclones Confer Resistance to Venetoclax-Based Therapy in Acute Myeloid Leukemia Patients. *Cancer Discov*. 2020;10(4):536–551.
- [9] Bliss CI. The toxicity of poisons applied jointly. *Ann Appl Biol*. 1939;26(3):585–615.

- [10] Greco WR, Bravo G, Parsons JC. The search for synergy: a critical review from a response surface perspective. *Pharmacol Rev.* 1995;47(2):331–385.
- [11] Foucquier J, Guedj M. Analysis of drug combinations: current methodological landscape. *Pharmacol Res Perspect.* 2015;3(3):e00149.
- [12] Ianevski A, He L, Aittokallio T, Tang J. SynergyFinder: a web application for analyzing drug combination dose–response matrix data. *Bioinformatics.* 2017;33(15):2413–2415.

10 Extended Supplementary Methods and Protocols

10.1 Detailed Cohort Processing and Multi-Omics Integration

BeatAML data (phs001657.v1.p1) were downloaded from dbGaP. Expression profiles were generated via RNA-sequencing of mononuclear cells from bone marrow and peripheral blood aspirates. Sequence reads were aligned to GRCh38, quantified as TPM, and $\log_2$ -transformed. Batch effects across sequencing centers (centerID) were corrected using ComBat empirical Bayes formulation [1]. The 320-patient "Gold Standard" multi-omics cohort represents primary aspirates with complete $\log_2$ expression, targeted variant calls for all 11 driver genes, ex vivo drug profiling across 155 compounds, and complete clinical overall survival data.

10.2 Consensus Clustering Formulation and Stability Analysis

Unsupervised consensus clustering was executed using ConsensusClusterPlus [6] over 1,000 iterations with 80% sample resampling, Euclidean distance metric, and Ward.D2 agglomerative linkage. Top variance genes were selected using Median Absolute Deviation (MAD). Consensus stability metrics included CDF slope area change, cluster consensus scores, and average silhouette width.

10.3 Supervised 50-Gene Classifier and 9-Gene VRS Score Algorithm

The 50-gene Random Forest classifier was built using the 'randomForest' package (ntree=1,000, mtry=7) and evaluated via 10-fold cross-validation. Predictor importance was ranked by mean decrease in Gini impurity. The continuous 9-Gene Venetoclax Response Score (VRS) is calculated as:

$$\text{VRS}_{\text{raw}} = \sum_{i=1}^9 w_i \cdot z(g_i) \quad (11)$$

where g_i represents the 9 key genes: *BCL2*, *NPM1*, *DNMT3A*, *MCL1*, *TP53*, *RUNX1*, *ASXL1*, *CD14*, and *FCGR1A*. Positive weights ($w_i = +1$) are assigned to primitive/BCL-2 addiction markers (*BCL2*, *NPM1*, *DNMT3A*), while negative weights ($w_i = -1$) are assigned to monocytic/resistance markers (*MCL1*, *TP53*, *RUNX1*, *ASXL1*, *CD14*, *FCGR1A*). Raw scores are standardized via reference Z-score transformation and rescaled to a 0–100 continuous scale.

10.4 Multivariable Variance Decomposition (R^2 Improvement Analysis)

To quantify independent predictive value for drug response, hierarchical linear models were fit:

$$\text{Model 1 (Mutational Baseline): } \text{AUC}_k = \beta_0 + \sum_{m=1}^{11} \beta_m \cdot \text{Mutation}_m + \varepsilon \quad (12)$$

$$\text{Model 2 (Combined Model): } \text{AUC}_k = \beta_0 + \sum_{m=1}^{11} \beta_m \cdot \text{Mutation}_m + \gamma \cdot \text{Subtype} + \varepsilon \quad (13)$$

The absolute variance improvement $\Delta R^2 = R^2_{\text{Model 2}} - R^2_{\text{Model 1}}$ and relative percentage improvement $\frac{\Delta R^2}{R^2_{\text{Model 1}}} \times 100\%$ were calculated. Hypothesis testing for $\gamma \neq 0$ was performed using ANOVA F -tests with Benjamini-Hochberg FDR adjustment.

10.5 Single-Cell CITE-seq and DepMap Functional Essentiality Protocols

Single-cell RNA-seq and ADT surface protein profiles from van Galen et al. [7] (38,412 cells) and Pei et al. [8] ($N = 4,426$ blasts) were processed using Seurat v4. Single-cell VRS scores were computed per cell. Cell-surface antibody-derived tags (CD34, CD117, CD11b, CD64) were normalized via CLR (Centered Log-Ratio) transformation. Cancer Dependency Map (DepMap 24Q2) Chronos gene knockout dependency scores for 24 myeloid leukemia lines were evaluated to confirm functional target essentiality.

11 Supplementary Figures




Figure S1: Technical Batch Mitigation and Batch Effect Correction. **A**, Principal Component Analysis (PCA) of raw gene expression data showing clear batch effects across sequencing centers. **B**, PCA of ComBat-seq corrected expression data showing removal of technical batch effects. **C**, Expression levels of top 10 batch-associated genes before correction. **D**, Expression levels of the same 10 genes after ComBat-seq correction, demonstrating that technical variation is successfully removed.




Figure S2: Pediatric TARGET-AML Overall Survival Validation. Kaplan-Meier overall survival analysis comparing molecular subtypes Cluster 1 ($N = 1368$) vs Cluster 2 ($N = 345$) in the pediatric TARGET-AML validation cohort, showing no significant difference ($p = 0.052$).





Figure S3: Baseline Overall Survival Partitioning across ELN 2017 Subgroups. Kaplan-Meier overall survival curves comparing Cluster 1 vs Cluster 2 within: **A**, ELN 2017 Favorable-risk cohort. **B**, ELN 2017 Intermediate-risk cohort. **C**, ELN 2017 Adverse-risk cohort. **D**, Hazard ratio forest plot across ELN risk strata.


A. Immune Landscape Deconvolution (xCell/MCP-counter)

Cluster 2 is significantly enriched for myelomonocytic and cytotoxic signatures



B. Lineage-Specific Enrichment

Significant increase in Neutrophils and CD8+ T cells ($p < 1e-15$)



C. The Mechanistic Link

Monocytic abundance drives Venetoclax resistance




Figure S4: Tumor Microenvironment Deconvolution and Cell Type Fraction Analysis. Immune and stromal cell type fractions deconvoluted across BeatAML transcriptomic subtypes showing monocytic vs progenitor cell population ratios.




Figure S5: Subgroup Benchmarking against LSC17, ROC Discrimination, and Decision Curve Analysis (DCA). **A**, Correlation between the 9-gene VRS and the prognostic LSC17 stemness score, demonstrating biological orthogonality. **B**, ROC curves comparing the discrimination accuracy of the VRS versus LSC17 for predicting ex vivo Venetoclax response. **C**, Decision Curve Analysis (DCA) showing the clinical net benefit of VRS-guided treatment selection. **D**, Bootstrap stability analysis of the ROC-AUC performance.




Figure S6: Metabolic Reprogramming, Dual Apoptotic Priming, and Pathway Integration in Monocytic Blast States. **A**, Signature scores for Oxidative Phosphorylation (OXPHOS) and Glycolysis showing Cluster 2 as a high-energy metabolic hybrid state. **B**, Ex vivo Bliss synergy matrix for Venetoclax + S63845 (MCL-1 inhibitor) combination in primary patient blasts, showing synergy. **C**, Heatmap showing coordinated enrichment of high-energy metabolic circuits in Cluster 2.




Figure S7: Functional Genomics and DepMap CRISPR Knockout Validation of VRS Targets. **A**, BCL2 gene expression versus CRISPR BCL2 dependency probability across cell lines. **B**, BCL2/MCL1 expression ratio versus CRISPR BCL2 dependency. **C**, CRISPR HDAC3 dependency versus VRS, demonstrating that lower VRS (resistant) aligns with greater HDAC3 vulnerability. **D**, CRISPR MAPK1 (ERK2) dependency versus VRS, showing MEK/ERK vulnerability in resistant cells.




Figure S8: Bulk Proteomic Target Validation Rescued by Functional Ratios and Tumor Purity Filtering. **A**, Spearman correlation between bulk protein VRS (pVRS) and ex vivo Venetoclax AUC across BeatAML proteomics ($N = 327$). **B**, BCL2/MCL1 protein ratio vs Venetoclax AUC. **C**, CD64/BCL2 protein ratio vs Venetoclax AUC. **D**, CD64/BCL2 protein ratio filtered for blast purity $\geq 50\%$.




Figure S9: Platform-Independent Validation of the Continuous VRS across Central European Trial Cohorts. Boxplots of continuous 9-gene VRS stratified by molecular subtype across independent European trial cohorts (AMLCG GPL96 $N = 163$, AMLCG GPL570 $N = 79$, AMLSG $N = 211$).




Figure S10: External Validation of the VRS and Molecular Subtypes in the Beat-AML 2.0 Cohort. **A**, Scatter plot of continuous VRS vs ex vivo Venetoclax AUC in independent BeatAML 2.0 validation cohort ($N = 367$). **B**, Boxplot comparing Venetoclax response across predicted subtypes.




Figure S11: Comprehensive Baseline Prognostic Overall Survival Validation. Kaplan-Meier overall survival curves comparing Cluster 1 vs Cluster 2 across discovery BeatAML, TCGA-LAML, and European multicenter cohorts.




Figure S12: Immunophenotypic Coupling of Single-Cell Resistance Hierarchy with ADT Cell Surface Markers. Single-cell CITE-seq ADT surface protein distributions (CD34, CD117, CD11b, CD64) across single-cell VRS tiers.




Figure S13: Longitudinal Selection Dynamics of Single-Cell VRS and Cell Surface ADT Markers under Venetoclax Combination Therapy. Violin plots comparing Diagnosis vs Relapse from matched longitudinal patient CITE-seq profiles under Venetoclax + Decitabine therapy.




Figure S14: UMAP Visualization of Single-Cell VRS and Cell Surface ADT Markers in GSE143363 CITE-seq Profiles. Single-cell UMAP projections colored by: **A**, Clinical timepoint (Diagnosis vs Relapse). **B**, Single-cell VRS. **C**, CD34 surface protein expression. **D**, CD64 surface protein expression.




Figure S15: Validation of the Monocytic Resistance Axis and Age-Independence of the VRS. A, Monocytic differentiation scores between Cluster 1 and Cluster 2. B, VIALE-A trial resistance signature score between clusters. C, Ex vivo Venetoclax response against patient age.




Figure S16: Cox Proportional Hazards Model Diagnostics and Subgroup Audits. **A**, Forest plot of subtype hazard ratio consistency across clinical and mutational subgroups. **B**, Schoenfeld residuals test for checking the proportional hazards assumption. **C**, Calibration plot comparing observed versus predicted 3-year survival. **D**, Restricted Mean Survival Time (RMST) difference analysis between subtypes.




Figure S17: Benchmarking of the 9-Gene VRS against ELN 2017/2022 Risk Groups. **A**, Boxplot of continuous VRS across ELN risk categories. **B**, Scatter plots of continuous VRS against ex vivo Venetoclax response (AUC) within each ELN risk category.




Figure S18: Single-Cell Resolution Validation of the 9-Gene VRS and Developmental Partitioning. Single-cell RNA-seq validation in bone marrow hierarchy (van Galen dataset, GSE116256): **A**, UMAP of cell types. **B**, VRS projection across cells. **C**, VRS by lineage. **D**, BCL2 vs MCL1 trade-off.

9-Gene VRS Noise Sensitivity Analysis

Robustness of clinical stratification under simulated technical noise




Figure S19: VRS Robustness and Technical Noise Sensitivity Analysis. Sensitivity analysis demonstrating classification concordance and Pearson correlation under simulated technical noise levels up to 50% of the true standard deviation.




Figure S20: Classifier Confidence Distribution and Interactive VRS Web Calculator. Poster mapping confidence distribution across external validation cohorts and Streamlit interactive bedside VRS web calculator interface.