

Effective spin-spin coupling in magnon-polarons toward acoustic-assisted probing of magnetic domains

Chongzhou Wang^{1,2}, Chensong Hua^{1,2}, and Weichao Yu^{1,3*}

¹State Key Laboratory of Surface Physics and Institute for Nanoelectronic Devices and Quantum Computing, Fudan University, Shanghai 200433, China

²Department of Physics, Fudan University, Shanghai 200433, China

³Zhangjiang Fudan International Innovation Center, Fudan University, Shanghai 201210, China

Received October 31, 2025; accepted December 26, 2025; published online February 5, 2026

We develop an effective spin-spin coupling framework to describe magnon-phonon hybridization in magnetoelastic systems, where both quasiparticles carry intrinsic spin angular momentum. The strength of the magnon-polaron anticrossing is shown to depend explicitly on the relative orientation of magnon and phonon spins, providing a unified framework for understanding chiral coupling. To validate this theory, we introduce a self-consistent frequency-domain micromagnetic simulation method incorporating full magnetoelastic coupling, which efficiently captures the dispersion and excitation dynamics of hybrid modes with excellent agreement between numerical and analytical results. Based on the polarization-selective magnon excitation governed by magnon-phonon spin alignment, we propose a scheme for acoustic-assisted probing of magnetic domains by exploiting polarization-controlled elastic waves. This work offers a new perspective for spin mechanics, where the spin degrees of freedom of both phonons and magnons are treated on equal footing, revealing their joint role in mediating coherent magnetoelastic couplings.

spin-spin coupling, magnon-phonon hybridization, micromagnetic simulation, acoustic probing

Citation: C. Wang, C. Hua, and W. Yu, Effective spin-spin coupling in magnon-polarons toward acoustic-assisted probing of magnetic domains, *Sci. China-Phys. Mech. Astron.* **69**, 247512 (2026), <https://doi.org/10.1007/s11433-025-2884-7>

1 Introduction

Magnetoelastic systems, where elastic and magnetic degrees of freedom coexist and interact, host rich physics arising from spin-lattice coupling [1-3]. As in other multi-physics platforms, magnetic detection techniques leverage the coupling between magnetization and external probes: optical, electrical, and quantum-microscopy approaches, such as Brillouin light scattering (BLS) [4, 5], the inverse spin Hall effect [6], magneto-optic Kerr effect [7] and nitrogen-vacancy centers [8, 9], which allow char-

acterization of magnonic excitations in real and reciprocal space. Recently, in the acoustic domain, surface acoustic waves (SAWs) can likewise be generated and probed non-destructively with high spatial resolution, for example, by BLS and micropillar-resonator methods [10, 11]. Coherent detection of the oscillating acoustoelectric effect in graphene enables phase-coherent electrical readout of SAW-driven electron dynamics in two-dimensional materials [12]. However, acoustic methods for probing magnetic structures remain underdeveloped, motivating the exploration of sound wave interactions with magnetic quasiparticles, particularly magnons, as a pathway for acoustic-based magnetic detection.

*Corresponding author (email: wcyu@fudan.edu.cn)

A typical quasiparticle in such systems is the magnon-polaron formed by magnon-phonon hybridization via magnetoelastic coupling (MEC) [13–17]. Interdigital transducers (IDTs) enable the excitation of various SAW modes and thereby acoustically driven ferromagnetic resonance, providing a route to drive and detect the hybrid dynamics [18–23]. Breaking symmetries through dipole-dipole interactions, interfacial Dzyaloshinskii-Moriya interaction, or Ruderman-Kittel-Kasuya-Yosida coupling can induce non-reciprocal propagation and directional control of magnon-polaron transport [24–35]. Alternatively, chiral magnon-phonon coupling can emerge from crystallographic polarization [36, 37]. Leveraging magnetoelastic couplings has enabled anisotropic polariton emission [38], acoustic-wave-assisted magnetization and vortex reversal [39, 40], generation and manipulation of domain walls and skyrmions [41–43], and tuning of antiferromagnetic phase transitions [44]. These advances highlight that the interaction of such hybrid excitations is not only a rich field for fundamental physics but also a promising basis for acoustic-assisted probing of magnetic states and domains.

The intrinsic degree of freedom, e.g., spin angular momentum of quasiparticles such as phonons and magnons, has recently attracted significant research interest [45–47]. For phonons, in close analogy to photons, circularly polarized transverse waves possess a longitudinal spin component aligned with the propagation direction, whereas evanescent and Gaussian waves typically exhibit transverse spin originating from amplitude decay mechanisms [48–52]. In SAWs, which inherently break spatial symmetry, the resulting distribution of transverse spin underpins spin-momentum locking [53–55], and has been harnessed to engineer chiral devices and phononic crystals with tailored functionalities [56, 57]. Turning to magnons, their quantum nature is that of elemental spin excitations. In an excited magnetic system, certain electron spins deviate from the ground-state alignment, and the collective contribution of these altered spins gives rise to the spin angular momentum carried by a magnon [58, 59]. Magnon spin has proven important across diverse phenomena, including magnonic band topology, the magnon spin Nernst effect, and magnon spin-orbit coupling effects [60–63]. Taken together, the effective spin-spin coupling framework captures rich information linking a quasiparticle's propagation direction to its intrinsic polarization for both phonons and magnons, offering a fresh perspective for analyzing and exploiting their coupled dynamics.

In this work, we develop a theoretical framework in which magnetization dynamics driven by elastic waves are interpreted through an effective spin-spin coupling between magnon spin and phonon spin. Within this framework, the MEC depends explicitly on both the relative orientation and

the amplitude of the two spins. We show that the anti-crossing (repulsion) magnitude in the magnon-polaron dispersion follows directly from the effective spin-spin coupling strength, and that the phonon-induced magnon response intensity is governed not only by spin orientations but also by the phonon-spin amplitude. This establishes a direct connection between the spatial distribution of magnetization and the local phonon-spin field. Leveraging this connection, we propose a detection protocol whereby the responses of magnons to elastic waves with controlled polarization in a multidomain ferromagnet reveal the magnetization orientation within each domain.

The remainder of this article is organized as follows. In sect. 2, we formulate an effective spin-spin coupling theory for MEC in a cubic lattice. In sect. 3, we introduce a self-developed technique, i.e., frequency-domain micromagnetic simulation with MEC, and validate the hybrid magnon-phonon modes against analytical results. In sect. 4, we compute the phonon-driven magnon response, reveal how phonon-spin amplitude and polarization govern excitation efficiency, and demonstrate its sensitivity to domain-specific magnetization orientations. In sect. 5, we propose a domain-resolved detection scheme based on polarization-controlled elastic-wave excitation and discuss experimental considerations and parameter regimes. We conclude in sect. 6 with a summary and an outlook.

2 Theory

We consider a magnetic film uniformly magnetized in x - y plane. The orientation of magnetization is denoted as (Θ, Φ) as illustrated in Figure 1. We neglect dipolar interactions for simplicity, as their effect can be absorbed into an effective anisotropy field for in-plane configurations. The energy density of magnetization is given by

$$\mathcal{H}_{\text{mag}} = -\mu_0 \mathbf{H} \cdot \mathbf{m} - \mu_0 K_u (\mathbf{m} \cdot \mathbf{u})^2 - \mu_0 \mathbf{H}_{\text{ex}} \cdot \mathbf{m}, \quad (1)$$

where $\mathbf{H}$ represents the external magnetic field, μ_0 denotes the vacuum permeability, and $\mathbf{m} = \mathbf{M}/M_s$ is the normalized magnetization with M_s being the saturation magnetization. The parameter K_u characterizes the uniaxial magnetic anisotropy along the unit vector $\mathbf{u}$, while $\mathbf{H}_{\text{ex}} = A\nabla^2 \mathbf{m}$ describes the Heisenberg exchange field with exchange stiffness constant A .

On the other hand, the energy density of elastic components in a solid is given by [64]

$$\mathcal{H}_{\text{el}} = \frac{1}{2} C_{ijkl} \epsilon_{ij} \epsilon_{kl}, \quad (2)$$

where we use the Einstein summation convention. Here, C_{ijkl}




Figure 1 (Color online) Schematic illustration of spin-spin couplings in MEC. (a) Blue arrows denote lattice vibrational displacements from circularly polarized transverse waves. (b) Red arrows represent magnetic moment precession in spin waves. Green arrows indicate the spin angular momentum of elastic spins, while orange arrows correspond to the spin angular momentum of spin waves. The MEC creates mutual interaction between elastic spin and magnon spin. Right panels illustrate the parameterization of these interactions: elastic wave polarization state (θ, ϕ) on the Bloch sphere (upper) and magnetic moment orientation (Θ, Φ) in the coordinate system (lower). This work focuses on in-plane configuration ($\Theta = \pi/2$) and assumes wave propagation along $+x$. (c) Schematic illustration of magnon-polaron anticrossing for different relative angles Φ between phonon and magnon spins.

is the elastic stiffness tensor and $\varepsilon_{ij} = (\partial u_i / \partial x_j + \partial u_j / \partial x_i) / 2$ are the strain tensor components, where u_i ($i = x, y, z$) represents the displacement fields.

The presence of MEC in a cubic crystal is described by the following energy density [2]

$$\mathcal{H}_{\text{mec}} = b_1 \sum_i m_i^2 \varepsilon_{ii} + b_2 \sum_{ij} m_i m_j \varepsilon_{ij}, \quad (3)$$

where b_1 and b_2 are the MEC constants.

Combining all contributions above, the total energy density of the magnetic film reads as:

$$\mathcal{H} = \mathcal{H}_{\text{mag}} + \mathcal{H}_{\text{el}} / M_s + \mathcal{H}_{\text{mec}}. \quad (4)$$

The dynamics of magnetization is described by the Landau-Lifshitz-Gilbert (LLG) equation [65-67]:

$$\dot{\mathbf{m}} = -\gamma \mathbf{m} \times \mu_0 \mathbf{H}_{\text{eff}} + \alpha \mathbf{m} \times \dot{\mathbf{m}}, \quad (5)$$

where γ is the gyromagnetic ratio and α is Gilbert damping coefficient. The effective field of magnetization is determined by the total energy density

$$\mu_0 \mathbf{H}_{\text{eff}} = -\frac{\partial \mathcal{H}}{\partial \mathbf{m}}. \quad (6)$$

Similarly, the equation of motion for elastic waves is given by

$$\rho \frac{\partial^2 u_i}{\partial t^2} = \frac{\partial \sigma_{ik}}{\partial x_k}, \quad (7)$$

where ρ is the mass density of the thin film and σ_{ik} are the components of the stress tensor. The stress tensor can be obtained from the total energy density according to [68]

$$\sigma_{ik} = \frac{1 + \delta_{ik}}{2} \frac{\partial \mathcal{H}}{\partial \varepsilon_{ik}}. \quad (8)$$

Under an arbitrary magnetic ground-state orientation $\mathbf{m}_0 = (\sin \Theta \cos \Phi, \sin \Theta \sin \Phi, \cos \Theta)$, we linearize the dynamical equations of the magnetic and elastic subsystems, eqs. (5) and (7). The resulting eigenvalue problem yields the dispersion relations of magnetoelastic waves

$$\begin{aligned} \rho \omega^2 u_x &= C_{11} k^2 u_x + 2b_1 ik \sin \Theta \cos \Theta (\sin \Phi M_2 \\ &\quad - \cos \Theta \cos \Phi M_1), \\ \rho \omega^2 u_y &= C_{44} k^2 u_y - b_2 ik \sin \Theta (2 \cos \Theta \sin \Phi \cos \Phi M_1 \\ &\quad + \cos 2\Phi M_2), \\ \rho \omega^2 u_z &= C_{44} k^2 u_z + b_2 ik (\cos \Theta \sin \Phi M_2 \\ &\quad - \cos 2\Theta \cos \Phi M_1), \end{aligned} \quad (9)$$

where for compactness, the rank-4 stiffness tensor C_{ijkl} is recast as a rank-2 matrix in Voigt notation.

In the absence of dipolar interactions and external fields, the bare magnonic dispersion is dominated by exchange interactions and is given by $\omega_m = \gamma \mu_0 (Ak^2 + 2K_u)$. The bare phonon dispersion exhibits a linear dependence on the wave vector in the long-wavelength limit $\omega_p = c_t k$, where $c_t = \sqrt{C_{44}/\rho}$ is the transverse wave velocity and ρ is the mass density of the elastic medium.

For the remainder of this work, we restrict our analysis to thin films with in-plane magnetization, i.e., $\Theta = \pi/2$. When magnons and phonons hybridize through MEC, considering a transverse wave linearly polarized in the y -direction (with displacement constraints $u_x = u_z = 0$) as a representative case and assuming the magnetization remains in-plane, the coupled dispersion relation takes the explicit form [19]

$$\frac{\omega^2}{\omega_p^2} = 1 - \frac{\gamma b_2^2 M_s}{c_t^2 \rho} \frac{\omega_m \cos^2 2\Phi}{\omega_m^2 - \omega^2}. \quad (10)$$

At the resonance condition where elastic and spin waves achieve frequency degeneracy ($\omega_p = \omega_m = \omega_r$), the linear

approximation becomes valid with $\omega^2 - \omega_r^2 \approx 2\omega_r(\omega - \omega_r)$. Under these conditions, the MEC lifts the degeneracy, resulting in two hybridized modes with frequencies

$$\omega_{\pm} = \omega_r \pm \frac{1}{2} \sqrt{\frac{\gamma b_2^2 M_s \omega_r}{c_t^2 \rho}} |\cos 2\Phi|. \quad (11)$$

This frequency splitting occurs at the critical wave vector k_r where $\omega_p(k_r) = \omega_m(k_r) = \omega_r$ and the bare magnon and phonon dispersions intersect, demonstrating the formation of anti-crossing behavior characteristic of hybridized magnetoelastic modes.

This phenomenon demonstrates the hybridization of magnon and phonon dispersion branches through MEC, resulting in characteristic anti-crossing behavior near the intersection point. The energy splitting magnitude Δ , defined as the eigenfrequency difference between the upper and lower hybridized modes at resonance ($\omega_+ - \omega_-$), serves as a quantitative measure of the magnetoelastic coupling strength.

The analysis can be extended to all polarization directions of elastic waves. For each polarization mode, the energy splitting at the resonance frequency ω_r is given by

$$\begin{aligned} \Delta_x &= |\sin 2\Phi| \sqrt{\frac{\gamma b_1^2 M_s \omega_r}{c_l^2 \rho}}, \\ \Delta_y &= |\cos 2\Phi| \sqrt{\frac{\gamma b_2^2 M_s \omega_r}{c_t^2 \rho}}, \\ \Delta_z &= |\cos \Phi| \sqrt{\frac{\gamma b_2^2 M_s \omega_r}{c_t^2 \rho}}, \end{aligned} \quad (12)$$

where Δ_x corresponds to x -polarized longitudinal modes with velocity $c_l = \sqrt{C_{11}/\rho}$, while Δ_y and Δ_z correspond to y - and z -polarized transverse modes with velocity $c_t = \sqrt{C_{44}/\rho}$, respectively.

To analyze the hybridization effects in magnetoelastic systems where orthogonal transverse modes coexist, we adopt a tripartite coupled oscillator framework [69, 70]. In this model, transverse elastic waves decompose into orthogonal y - and z -polarized eigenmodes, each independently coupling to the magnonic system while maintaining zero direct inter-mode coupling. For generic elliptical phonon polarization states characterized by arbitrary displacement component ratios, both polarization components cooperatively contribute to magnon-phonon anticrossing through parallel coupling channels. The absence of direct y - z phonon coupling enables linear superposition of the individual hybridization effects,

yielding a total energy splitting

$$\begin{aligned} \Delta &= \sqrt{\Delta_y^2 + \Delta_z^2} \\ &= \sqrt{\frac{\gamma b_2^2 M_s \omega_r}{c_t^2 \rho}} \sqrt{\cos^2 2\Phi + \cos^2 \Phi}. \end{aligned} \quad (13)$$

The angular dependence of the magnetoelastic energy splitting reflects a fundamental aspect of the underlying physics that the interaction between magnon and phonon systems can be understood as an effective spin-spin coupling mechanism, as schematically shown in Figure 1(a) and (b). This perspective emerges naturally when we recognize that both magnons and phonons carry intrinsic angular momentum, which can be characterized by their respective spin vectors. The spectrum of the magnon-polaron hybrid mode depends critically on the relative angle between the phonon spin ($\mathbf{S}_p$) and the magnon spin ($\mathbf{S}_m$), as illustrated in Figure 1(c).

The magnon spin moment (denoted as $\boldsymbol{\mu}_k$) refers to the magnetic moment carried by a single magnon excitation with wave vector k . Within the classical continuum framework, the moment is approximately given by $\boldsymbol{\mu}_k \simeq -\mu_B |\delta \mathbf{m}_k|^2 \mathbf{m}_0$ [58], where μ_B is the Bohr magneton, $\mathbf{m}_0$ is the unit vector along the equilibrium magnetization and $\delta \mathbf{m}_k$ denotes the transverse component of the spin-wave excitation associated with wave vector k . The negative sign signifies that the presence of a magnon, which corresponds to a reduction of spin angular momentum along the equilibrium direction, leads to a decrease in the magnetic moment parallel to $\mathbf{m}_0$ as illustrated in Figure 1(b). We define the density of magnon spin as:

$$\mathbf{S}_m = \frac{\boldsymbol{\mu}_k}{\gamma V}, \quad (14)$$

where V is the volume of the magnetic material.

Similarly, transverse acoustic phonons carry intrinsic angular momentum due to the rotational motion of the elastic medium. The instantaneous density of phonon spin $\mathbf{S}_p$ can be defined as [48, 49]¹⁾:

$$\mathbf{S}_p = \frac{\rho \omega}{2} \text{Im}(\mathbf{u}^* \times \mathbf{u}), \quad (15)$$

where $\mathbf{u}$ is the displacement field. For circularly polarized transverse waves, the phonon spin aligns parallel (right-handed) or antiparallel (left-handed) to the propagation direction, analogous to photon helicity. The quantity $\mathbf{S}_p$ represents the intrinsic spin angular momentum density of the elastic wave, arising from its local circular polarization state rather than from orbital motion of material points. This definition follows the continuum-field convention established in

1) The term “phonon spin” here refers to the intrinsic spin angular momentum density associated with circularly polarized elastic waves in the continuum description. This should not be confused with intrinsic quantum spin. Its microscopic origin lies entirely in the orbital-like mechanical motion of atoms, which is referred to as orbital angular momentum. The apparent discrepancy arises from differing perspectives and terminological conventions, not from any physical inconsistency.

phononics [48,49] and is directly analogous to the spin angular momentum of circularly polarized light in optics [71-73], ensuring a consistent quasiparticle-level description that parallels the spin of magnons.

In this work, we assume without loss of generality that all elastic waves propagate along the $+x$ -direction. This allows us to define the relative orientation between phonon spin and magnon spin unambiguously in the transverse plane, enabling a simplified angular description of their coupling. The relative orientation angle Φ between $\mathbf{S}_m$ and $\mathbf{S}_p$ thus governs the strength of the effective spin-spin coupling, providing a microscopic foundation for understanding the angular-dependent energy splitting in magnetoelastic systems, and offers a unified framework for describing magnon-phonon hybridization across different material systems and geometric configurations. Building on the established framework of magnon and phonon spins, eq. (13) can be reformulated using their normalized direction vectors $\mathbf{s}_m = \mathbf{S}_m/|\mathbf{S}_m|$ and $\mathbf{s}_p = \mathbf{S}_p/|\mathbf{S}_p|$, such that $\mathbf{s}_p \cdot \mathbf{s}_m = \cos \Phi$, and the hybridization gap can be expressed as:

$$\Delta = \sqrt{\frac{\gamma b_2^2 M_s \omega_r}{c_t^2 \rho} (4(\mathbf{s}_p \cdot \mathbf{s}_m)^4 - 3(\mathbf{s}_p \cdot \mathbf{s}_m)^2 + 1)}. \quad (16)$$

It is noteworthy that the magnitude of the level splitting depends solely on even-order terms of the spin inner product $\mathbf{s}_p \cdot \mathbf{s}_m = \cos \Phi$. This dependence reflects the invariance of the interaction under the transformation $\Phi \rightarrow -\Phi$, indicating that the coupling strength is identical for clockwise and counter-clockwise rotations between the magnon and phonon spins. The functional form $f(\cos \Phi) = 4 \cos^4 \Phi - 3 \cos^2 \Phi + 1$ exhibits the characteristic $\cos(2\Phi)$ periodicity due to the dominance of the quadratic term, resulting in a twofold symmetric angular dependence of the coupling strength. According to the coupling strength, the dispersion relation of the magnon polaron eq. (11) can be given by

$$\omega_{\pm} = \frac{1}{2} \left[(\omega_p + \omega_m) \pm \sqrt{(\omega_p - \omega_m)^2 + \Delta^2} \right]. \quad (17)$$

When the magnon spin aligns parallel with the phonon spin ($\mathbf{s}_p \cdot \mathbf{s}_m = \pm 1$), the repulsion strength reaches its maximum value.

3 Numerical simulation

To validate the theoretical predictions presented above, we develop a comprehensive frequency-domain micromagnetic simulation framework [74] that further incorporates MEC. In contrast to conventional time-domain micromagnetic approaches that require time-evolution calculations followed by Fourier transformation, this frequency-domain method offers

remarkable computational efficiency by directly calculating the system's response under specific frequency excitations.

The simulation is performed in COMSOL Multiphysics using the Micromagnetics Module (Frequency Domain) [74] and the built-in Solid Mechanics Module. In this frequency-domain framework, magnetization dynamics is pre-transformed using temporal Fourier transformation, $\mathbf{m}(\mathbf{r}, t) \rightarrow \mathbf{m}(\mathbf{r}, \omega)$, with the assumption that the magnetic moment $\mathbf{m}$ is perturbed around its equilibrium magnetization $\mathbf{m}_0$ with certain angular frequency ω . The unit magnetization vector can be decomposed as $\mathbf{m} = \mathbf{m}_0 + \delta \mathbf{m} e^{i\omega t}$ under linear approximation $\delta \mathbf{m} \ll \mathbf{m}_0$ and $\mathbf{m}_0 \cdot \delta \mathbf{m} = 0$. Following the same procedures, the effective field can be decomposed into static and dynamic components as $\mathbf{H}_{\text{eff}} = \mathbf{h}_0^{\text{eff}} + \delta \mathbf{h}^{\text{eff}} e^{i\omega t}$. Substituting these expressions for $\mathbf{m}$ and $\mathbf{H}_{\text{eff}}$ into the time-dependent LLG equation yields

$$i\omega \delta \mathbf{m} = -\gamma \mu_0 (\mathbf{m}_0 \times \delta \mathbf{h}^{\text{eff}} + \delta \mathbf{m} \times \mathbf{h}_0^{\text{eff}}) + i\omega \alpha \mathbf{m}_0 \times \delta \mathbf{m}, \quad (18)$$

where high-order terms such as $\delta \mathbf{m}^2$ and $\delta \mathbf{h}^2$ are neglected. With the linear approximation, the effective fields contributed by the MEC $\mu_0 \mathbf{H}_{\text{eff}}^{\text{mec}} = -\partial \mathcal{H}_{\text{mec}} / \partial \mathbf{m}$ can be obtained as:

$$\begin{aligned} \mathbf{H}_{\text{eff}}^{\text{mec}} = & -\frac{2b_1}{\mu_0} \begin{pmatrix} m_{0x} \frac{\partial u_x}{\partial x} \\ m_{0y} \frac{\partial u_y}{\partial y} \\ m_{0z} \frac{\partial u_z}{\partial z} \end{pmatrix} \\ & -\frac{b_2}{\mu_0} \begin{pmatrix} m_{0y} (\frac{\partial u_x}{\partial y} + \frac{\partial u_y}{\partial x}) + m_{0z} (\frac{\partial u_z}{\partial x} + \frac{\partial u_x}{\partial z}) \\ m_{0x} (\frac{\partial u_x}{\partial y} + \frac{\partial u_y}{\partial x}) + m_{0z} (\frac{\partial u_y}{\partial z} + \frac{\partial u_z}{\partial y}) \\ m_{0y} (\frac{\partial u_y}{\partial z} + \frac{\partial u_z}{\partial y}) + m_{0x} (\frac{\partial u_z}{\partial x} + \frac{\partial u_x}{\partial z}) \end{pmatrix}. \end{aligned} \quad (19)$$

On the other hand, for isotropic elastic media, the frequency-domain equation of motion for elastic waves (Navier's equation) [75, 76] can be formally expressed through displacement fields:

$$-\rho \omega^2 \mathbf{u} = \mu \nabla^2 \mathbf{u} + (\lambda + \mu) \nabla (\nabla \cdot \mathbf{u}) + \mathbf{f}_{\text{mec}}, \quad (20)$$

where μ and λ represent Lamé's first and second parameters, respectively. The Lamé's constants and the stiffness matrix in Voigt notation are formally equivalent representations $C_{11} = \lambda + 2\mu$, $C_{44} = \mu$ [49]. The magnetoelastically mediated body forces $\mathbf{f}_{\text{mec}}$ imposed by spin-wave excitations upon the elastic medium can be mathematically formulated through the functional derivative of the magnetoelastic energy density with respect to the displacement gradient [77]. For the cubic MEC considered here, the effective force den-

sity exerted by the magnetization dynamics reads

$$\begin{aligned} \mathbf{f}_{\text{mec}} = & 2b_1 M_s \begin{pmatrix} m_{0x} \frac{\partial \delta m_x}{\partial x} + \delta m_x \frac{\partial m_{0x}}{\partial x} \\ m_{0y} \frac{\partial \delta m_y}{\partial y} + \delta m_y \frac{\partial m_{0y}}{\partial y} \\ m_{0z} \frac{\partial \delta m_z}{\partial z} + \delta m_z \frac{\partial m_{0z}}{\partial z} \end{pmatrix} \\ & + b_2 M_s \begin{pmatrix} m_{0y} \frac{\partial \delta m_x}{\partial y} + \delta m_y \frac{\partial m_{0x}}{\partial y} + m_{0x} \frac{\partial \delta m_y}{\partial x} + \delta m_x \frac{\partial m_{0y}}{\partial x} \\ m_{0z} \frac{\partial \delta m_x}{\partial z} + \delta m_z \frac{\partial m_{0x}}{\partial z} + m_{0x} \frac{\partial \delta m_z}{\partial x} + \delta m_x \frac{\partial m_{0z}}{\partial x} \\ m_{0z} \frac{\partial \delta m_y}{\partial z} + \delta m_z \frac{\partial m_{0y}}{\partial z} + m_{0y} \frac{\partial \delta m_z}{\partial z} + \delta m_y \frac{\partial m_{0z}}{\partial z} \end{pmatrix} \\ & + b_2 M_s \begin{pmatrix} m_{0y} \frac{\partial \delta m_z}{\partial x} + \delta m_y \frac{\partial m_{0z}}{\partial x} + m_{0z} \frac{\partial \delta m_y}{\partial x} + \delta m_z \frac{\partial m_{0y}}{\partial x} \\ m_{0x} \frac{\partial \delta m_z}{\partial y} + \delta m_x \frac{\partial m_{0z}}{\partial y} + m_{0z} \frac{\partial \delta m_x}{\partial y} + \delta m_z \frac{\partial m_{0x}}{\partial y} \\ m_{0x} \frac{\partial \delta m_y}{\partial z} + \delta m_x \frac{\partial m_{0y}}{\partial z} + m_{0y} \frac{\partial \delta m_x}{\partial z} + \delta m_y \frac{\partial m_{0x}}{\partial z} \end{pmatrix}. \end{aligned} \quad (21)$$

Building upon this frequency-domain framework, we further implement a Floquet boundary condition that extends the Fourier transformation from the temporal domain to the spatial domain, enabling efficient calculation of dispersion relations $\mathbf{m}(\mathbf{r}, \omega) \rightarrow \mathbf{m}(\mathbf{k}, \omega)$. This boundary condition can specify arbitrary phase differences of dependent variables between the source (src) and destination (dst) boundaries with the following form:

$$\delta \mathbf{m}_{\text{dst}} = \delta \mathbf{m}_{\text{src}} e^{-i\mathbf{k}_F \cdot (\mathbf{r}_{\text{dst}} - \mathbf{r}_{\text{src}})}, \quad (22)$$

where $\mathbf{k}_F$ is the Floquet wave vector (k -vector for Floquet periodicity), $\mathbf{r}_{\text{src}}$ and $\mathbf{r}_{\text{dst}}$ are the positions of the source and destination boundaries, respectively. In contrast to standard (e.g., open or periodic) boundary conditions, which confine waves within a finite domain and generate spurious standing-wave resonances due to reflections, the Floquet condition enforces a traveling-wave phase relationship across the boundaries. This eliminates artificial finite-size effects that would otherwise distort the continuous magnon-polaron spectrum and degrade k -space resolution. With this Floquet boundary condition established in eq. (22), dispersion relations can be efficiently computed using minimal computational resources and significantly reduced unit cells. This approach enables effective and high-precision calculation of dispersion characteristics in the present work, while also providing a versatile tool for readily computing dispersion relations of other periodic structures, such as magnonic crystals [78, 79]. The following simulations employ material parameters of Yttrium Iron Garnet (YIG), incorporating both its magnetic and elastic properties [47, 67]: $\gamma\mu_0 = 2.21 \times 10^5 \text{ m/(s} \cdot \text{A)}$, $M_s = 0.194 \times 10^6 \text{ A/m}$, $A = 0.328 \times 10^{-10} \text{ A} \cdot \text{m}$, $2K_u = 0.388 \times 10^5 \text{ J/m}^3$, $\rho = 5.17 \times 10^3 \text{ kg/m}^3$, $b_2 M_s = 2b_1 M_s = 6.96 \times 10^5 \text{ J/m}^3$, $\lambda = 1.16 \times 10^{11} \text{ Pa}$, $\mu = 7.64 \times 10^{10} \text{ Pa}$, resulting sound speed $c_l = 7209 \text{ m/s}$, $c_t = 3843 \text{ m/s}$.

To verify the consistency between our numerical framework and theoretical predictions, we compare the simulated

spin wave dispersion with analytical results in Figure 2. The colormap superimposed on the background represents the numerically simulated spatial profile of spin wave amplitudes, while the dashed colored curves correspond to dispersion relations calculated from eq. (17). The quantitative agreement demonstrates that maximum spin wave excitation amplitude occurs exclusively when the driving frequency and wave vector simultaneously satisfy the dispersion criteria, thereby validating our numerical approach.

The energy splitting in the dispersion relation exhibits angular dependence on the relative orientation between the magnon and phonon spins. Maximum splitting in Figure 2(a) is observed under collinear spin configurations (antiparallel alignment of phonon and magnon spins), reaching a magnitude $\sqrt{2}$ times greater than that in the orthogonal configuration shown in Figure 2(b). Notably, the apparent null splitting at parallel alignment (illustrated in Figure 2(c)) arises from chirality mismatch mechanisms that excite dark modes rather than from actual degeneracy. The dependence of resonant frequencies on arbitrary relative angles Φ between magnon and phonon spin, measured at specific resonant wave vectors, is numerically resolved in Figure 2(d). For each wave vector, three eigenmodes are identified, with the 1.37 GHz mode constituting a dark mode that remains decoupled from the external driving field. This dark mode represents an eigen-




Figure 2 (Color online) The dispersion of magnon polarons formed by coupling between right-handed elastic waves and spin waves at different relative angles between magnon and phonon spin. The dashed colored curves represent theoretical dispersion relations obtained from eq. (17), while the background colormaps show numerical results of spin wave amplitude $|\delta m|$ per unit volume. (a) Antiparallel alignment: $\Phi = 0$, $\mathbf{s}_p \cdot \mathbf{s}_m = -1$, showing maximum energy splitting. (b) Orthogonal alignment: $\Phi = \pi/2$, $\mathbf{s}_p \cdot \mathbf{s}_m = 0$, demonstrating reduced splitting. (c) Parallel alignment: $\Phi = \pi$, $\mathbf{s}_p \cdot \mathbf{s}_m = 1$, exhibiting apparent null splitting due to dark mode excitation. (d) Angular dependence of energy splitting at resonant wave vectors as a function of relative angle Φ , confirming the theoretical predictions.

state with negligible coupling strength to the applied excitation, resulting in minimal observable amplitude despite being an intrinsic solution of the eigenvalue spectrum. Beyond the angular-modulated energy splitting, the relative excitation amplitudes of these eigenmodes exhibit pronounced angular sensitivity to spin alignment configurations.

Our analysis as well as micromagnetic simulations neglect explicit dipolar fields, which is valid for thin films or high- k modes where exchange dominates. We have verified that including full magnetostatic interactions introduces nonreciprocity due to surface magnons but leaves the magnon-polaron anticrossing gap unchanged. While this work focuses on bulk transverse phonons whose spin is collinear with propagation, extending the framework to surface acoustic waves (where phonon spin is transverse) represents an important direction for future study.

4 Polarization-selective magnon excitation

The inherent chiral symmetry breaking in ferromagnetic magnon systems establishes spin-dependent selection rules that govern magnon activation efficiency under external driving forces. This polarization selectivity originates from the fundamental property that magnons in ferromagnetic systems possess exclusively right-handed polarization due to the sign of the gyromagnetic ratio. Consequently, phonons with right-handed spin (corresponding to right-handed circularly polarized elastic waves) achieve optimal phase coherence with magnon spin through MEC, enabling strong resonant excitation. In contrast, phonons with left-handed spin experience chirality mismatch with magnon spin, resulting in weak-amplitude forced oscillations with excitation efficiencies reduced by over an order of magnitude, as quantitatively verified in Figure 2(d).

The phonon spin can be parameterized using the relative amplitude and phase of displacement field components. Following the representation on the Bloch sphere [80], we define $u_y \propto \sin(\theta/2)$ and $u_z \propto \cos(\theta/2)e^{i\phi}$, where θ represents the relative amplitude ratio between the two transverse displacement components and ϕ denotes their relative phase difference. This parameterization yields the phonon spin $\mathbf{s}_p = \sin\theta \sin\phi \hat{\mathbf{x}}$. Physically, $\theta = 0$ corresponds to linearly polarized waves along the z -direction, $\theta = \pi/2$ with $\phi = \mp\pi/2$ generates right/left-handed circularly polarized waves, and $\theta = \pi$ represents linear polarization along the y -direction. While altering the phonon spin orientation (i.e., the polarization state of elastic waves) does not affect the energy splitting of magnon polarons, it dramatically modulates the magnon excitation intensity, as demonstrated in Figure 3(a) and (b). The energy splitting is entirely determined by the relative angle between the magnon and phonon spins,

as governed by the effective spin-spin coupling perspective discussed above.

To quantify this polarization-selective magnon excitation, we define the total excitation intensity as:

$$P = \int |\delta\mathbf{m}(k_r)| d\omega, \quad (23)$$

where the integration is performed over the complete resonance spectrum at the critical wave vector k_r , encompassing both the lower and upper hybridized modes. Systematic analysis of phonon spin parameter space reveals distinct modulation patterns: when the relative magnitude parameter θ remains constant, varying the phase ϕ shifts the minimum intensity magnitude while maintaining the minimum position consistently at $2\pi/3$ and $4\pi/3$ (Figure 3(c)). Conversely, when the phase ϕ is held constant, adjusting the relative magnitude θ preserves the minimum intensity magnitude but shifts the minimum position within the range $\pi/3$ to $\pi/2$ and $3\pi/2$ to $5\pi/3$ (Figure 3(d)).

The magnon excitation intensity exhibits strong polarization selectivity that depends on the relative orientation between phonon and magnon spins. Through systematic parametric scanning of phonon polarization configurations, we identify optimal states that maximize magnonic excitation efficiency, as demonstrated in Figure 4(b)-(e). The total excitation intensity is computed using eq. (23) integrated over the frequency range from 0 to 5 GHz. Under collinear spin alignment, right-handed circularly polarized phonons ($\theta = \pi/2, \phi = 3\pi/2$) achieve maximum spin-wave activation, as shown in Figure 4(b). As the magnon spin cants away from the phonon spin direction, the extremal excitation positions shift systematically while remaining fixed at specific phase values $\phi = \pi/2$ and $\phi = 3\pi/2$ across all angular configurations (Figure 4(c)-(e)).

The systematic evolution of these excitation extrema is mapped in Figure 4(a), where each phonon polarization state corresponds to a spin coordinate on the hybridized Bloch sphere. The angular dependence of extremum positions is quantified in Figure 4(f) and (g), which extracts the polar coordinates of maximum and minimum excitation points as functions of the spin canting angle. These data reveal that extremal intensity trajectories trace three-quarter arc segments ($\Delta\theta = 3\pi/2$) upon complete spin reorientation, confirming the geometric correspondence between phonon polarization configurations and magnonic response on the Bloch sphere.

5 Proposal for acoustic-assisted probing of magnetic domains

Motivated by the chiral effective spin-spin coupling between




Figure 3 (Color online) Magnon excitation efficiency under elastic wave driving with variable phonon spin configurations. (a) Magnon excitation spectra driven by elastic waves with fixed amplitude parameter $\theta = \pi/2$ and variable phase parameter $\phi \in \{0, \pi/4, \pi/2\}$, demonstrating phase-dependent magnon resonance activation (top insets show corresponding elastic wave polarization ellipses). (b) Complementary analysis of magnon excitation under elastic wave driving with fixed phase $\phi = 3\pi/2$ and variable amplitude ratio $\theta \in \{0, \pi/4, \pi/2\}$ (polarization geometries shown above). (c) and (d) Total excitation intensity P as a function of spin misalignment angle for configurations in panels (a) and (b), respectively.




Figure 4 (Color online) Polarization-selective magnon excitation under phonon spin configurations. (a) Normalized total excitation intensity maps for spin canting angles $0, \pi/6, \pi/3, \pi/2$, with blue (red) curves tracking minima (maxima) trajectories. Upper insets show corresponding trajectories on the Bloch sphere representation. (b)-(e) Individual excitation intensity distributions for each spin canting angle, with insets indicating optimal chiral phonon states. (f) and (g) Extracted minima and maxima positions as functions of spin canting angle, respectively.

magnons and phonons derived in sect. 2, we propose a non-invasive, polarization-resolved acoustic imaging technique for mapping in-plane magnetic domain textures. By exploiting the angular selectivity of magnon excitation under circularly or elliptically polarized bulk acoustic waves (BAWs) [49, 81], this method transforms elastic wave polarization into a spatially resolved probe of local magnetization orientation, enabling acoustic-assisted magnetic domain visualization with optical or electrical contacts. Unlike surface acoustic waves (SAWs), whose polarization states are geometrically constrained by boundary conditions and surface symmetry [53, 55], BAWs permit full three-dimensional control over phonon spin via independent tuning of transverse displacement components, making them ideal for probing chiral magnon-phonon interactions.

To quantitatively validate this concept, we perform micromagnetic simulations based on the frequency-domain formalism newly developed in this work (sect. 3), which self-consistently solves the coupled LLG and elastodynamic equations under harmonic BAW driving. The simulated structure consists of a $10\ \mu\text{m} \times 10\ \mu\text{m}$ epitaxial YIG thin film ($2K_u = 1 \times 10^5\ \text{A/m}$) on a piezoelectric substrate, hosting 16 magnetic domains with Néel-type spin textures. To mimic realistic polycrystalline or disordered magnetic films while preserving computational tractability, the easy-axis orientations of these domains are randomly sampled within the first quadrant ($\Phi \in [0^\circ, 90^\circ]$), specifically assigned to 0° , 30° , 60° , and 90° for systematic analysis. This restriction to the first quadrant is physically justified by the fourfold rotational symmetry of the MEC energy density (eq. (3)), which ensures identical response patterns under 90° rotations. Each domain has an average area of $\approx 6.25\ \mu\text{m}^2$, with characteristic length of

domain walls $\approx 18\ \text{nm}$, significantly narrower than the BAW wavelength ($1.281\ \mu\text{m}$ at $f = 3\ \text{GHz}$), ensuring intra-domain uniformity. BAWs are launched along $+x$ via interdigitated transducers (IDTs), with polarization (equivalent to phonon spin s_p) controlled through the transverse displacement ratio $\eta = u_y/u_z$ via the body forces $\mathbf{F} = (0, F_y, F_z)e^{i\omega t}$, where $F_y = 2 \times 10^{-5} \sin(\theta/2)\ \text{N}$ and $F_z = 2 \times 10^{-5} \cos(\theta/2)e^{i\phi}\ \text{N}$.

The simulated spatial maps of magnon excitation intensity $|\delta\mathbf{m}|$, shown in Figure 5(b)–(e), reveal pronounced polarization-dependent contrast across domains, with each subfigure corresponding to a distinct BAW polarization state controlled by $\eta = u_y/u_z$. Under right-handed circular polarization ($\eta = 1$, Figure 5(b)), the strongest excitation occurs in domains with $\Phi = 0^\circ$ magnetization, while the weakest response is observed in 60° domains. When η is increased to 5.8 (Figure 5(c)), the response profile inverts: 90° domains now dominate the excitation, while 30° domains become minimally responsive. Reducing η to 0.68 (Figure 5(d)) shifts peak intensity to 60° domains, with 0° domains exhibiting the lowest response. Finally, at $\eta = 0.42$ (Figure 5(e)), maximum excitation is observed in 30° domains, while 90° domains are suppressed. This systematic angular mapping arises directly from the $\cos^4\Phi$ -like dependence of the coupling strength. The total excitation intensity, defined as $P_k = \int |\delta\mathbf{m}(\omega_r)| dk$ (integrated over wave vector at resonance frequency), provides a scalar metric for domain classification. Figure 5(f) and (g) identify the polarization sectors (red/green) that maximize or minimize P_k for 0° and 60° domains, respectively, establishing a predictive framework for reconstructing local magnetization from multi-polarization acoustic measurements. We note that the simulated device size ($10\ \mu\text{m}$) is much smaller than the typical acoustic




Figure 5 (Color online) (a) Schematic of BAW-driven magnon excitation in a multidomain YIG film (inset: domain map with easy-axis orientations). (b) Magnon excitation map under right-handed circular polarization ($\eta = 1$). (c) Response under $\eta = 5.8$. (d) Response under $\eta = 0.68$. (e) Response under $\eta = 0.42$. Insets in (b)–(e): polarization ellipses corresponding to each η . (f) and (g) Polarization sectors (red/green) that maximize or minimize total excitation intensity for domains oriented at 0° and 60° , respectively.

attenuation length in YIG/GGG systems (>1 mm) [82, 83], justifying the neglect of phonon damping. Domain walls (~ 18 nm wide) are also much narrower than both the acoustic wavelength (~ 1 μ m) and domain size (2–3 μ m), minimizing scattering. An absorbing boundary is implemented to reduce reflections. Interference fringes in Figure 5 arise from weak scattering at domain edges but do not affect the main coupling signature.

To implement this probing scheme experimentally, a sequence of elastic waves with controlled polarization states is excited in the ferromagnetic film using IDTs. By adjusting the phase and amplitude of the driving signals, the phonon spin orientation can be systematically varied across the full range of possible configurations. At each measurement location, the resulting magnon response is detected locally using BLS [11]. The measured excitation efficiency as a function of phonon spin forms a distinct signature for the local magnetization direction. This experimental signature is then compared with pre-established response patterns derived from theoretical modeling or calibration measurements. This approach enables spatially resolved, non-invasive characterization of magnetic domain structures through selective coupling between phonon and magnon spins.

Crucially, unlike conventional broadband or off-resonance acoustic probing schemes, our approach operates precisely at the magnon-phonon frequency resonance condition. This technique achieves angular resolution better than 30° and is compatible with standard IDT-driven platforms, enabling dynamic, non-optical magnetic imaging under diverse environmental conditions. Key limitations include sensitivity to film inhomogeneity, the requirement for coherent multi-polarization excitation, and the need for calibration against known domain orientations. Importantly, equivalent domain mapping can be achieved by fixing BAW polarization and rotating the IDT orientation, which offers experimental flexibility. While our simulations focus on the first quadrant for symmetry-based simplification, the full method is generalizable to arbitrary in-plane magnetization angles, as the underlying coupling physics is rotationally covariant. Although this work focuses on BAWs for maximal polarization control, the technique can be adapted to SAW-based systems [84, 85], albeit with reduced polarization degrees of freedom, opening pathways toward nondestructive evaluation of magnetic heterostructures.

6 Conclusion

We have established an effective spin-spin coupling framework to describe magnon-phonon hybridization, demonstrating that the interaction strength and excitation efficiency

are governed by the relative orientation and magnitude of magnon and phonon spins, which are intrinsic angular momentum carriers in their respective subsystems. Validated by fully-coupled simulations in the frequency domain, our results reveal that the anticrossing gaps exhibit a clear angular dependence and that the magnon response shows strong polarization selectivity, where right-handed phonons resonantly drive ferromagnetic magnons as a result of chiral matching between their spins. Based on this mechanism, we propose that spatially resolved acoustic excitation with engineered phonon spin might, in principle, offer a pathway toward probing local magnetic domain configurations, although this approach remains to be experimentally verified and is subject to practical limitations that require further exploration.

While our minimal model focuses on local MEC governed by the relative orientation of magnon and phonon spins in bulk media, it establishes a foundational framework for understanding more complex chiral interactions arising from spin-momentum locking in surface waves or in the presence of non-local interaction such as dipolar fields. Looking ahead, an important next step is to generalize this effective spin-spin coupling perspective to antiferromagnetic systems, where the presence of two magnetic sublattices leads to magnon modes that carry opposite chiralities [86, 87]. Extending this framework to antiferromagnetic systems would provide spin mechanics with a new perspective, where phonons in solids and magnons in antiferromagnets fully harness their chiral degrees of freedom. This enables a more complete description of phenomena such as non-reciprocal transport and chiral mode hybridization, offering new physical insights that go beyond conventional scalar or isotropic coupling models.

This work was supported by the National Key Research and Development Program of China (Grant No. 2022YFA1403300), Quantum Science and Technology-National Science and Technology Major Project (Grant No. 2024ZD0300103), and the National Natural Science Foundation of China (Grant No. 12204107). The authors thank Yang Long for fruitful discussion.

Conflict of interest The authors declare that they have no conflict of interest.

- 1 A. Rückriegel, S. Streib, G. E. W. Bauer, and R. A. Duine, *Phys. Rev. B* **101**, 104402 (2020).
- 2 C. Kittel, *Phys. Rev.* **110**, 836 (1958).
- 3 A. E. Clark, B. DeSavage, W. Coleman, E. R. Callen, and H. B. Callen, *J. Appl. Phys.* **34**, 1296 (1963).
- 4 S. O. Demokritov, B. Hillebrands, and A. N. Slavin, *Phys. Rep.* **348**, 441 (2001).
- 5 C. Banerjee, P. Gruszecki, J. W. Klos, O. Hellwig, M. Krawczyk, and A. Barman, *Phys. Rev. B* **96**, 024421 (2017).
- 6 O. Mosendz, V. Vlaminc, J. E. Pearson, F. Y. Fradin, G. E. W. Bauer, S. D. Bader, and A. Hoffmann, *Phys. Rev. B* **82**, 214403 (2010).
- 7 J. Kerr, *London Edinburgh Dublin Philos. Mag. J. Sci.* **3**, 321 (1877).
- 8 F. Casola, T. van der Sar, and A. Yacoby, *Nat. Rev. Mater.* **3**, 17088 (2018).

- 9 J. R. Maze, P. L. Stanwix, J. S. Hodges, S. Hong, J. M. Taylor, P. Cappellaro, L. Jiang, M. V. G. Dutt, E. Togan, A. S. Zibrov, et al., *Nature* **455**, 644 (2008).
- 10 T. Lee, H. S. Kwon, X. Li, Z. Yu, T. Kimura, Y. Tadokoro, and H. Iizuka, *Phys. Rev. B* **111**, 174101 (2025).
- 11 J. Yang, M. Y. Guo, Z. L. Li, P. Wu, K. M. Cai, X. Z. Liu, Y. G. Peng, Q. Wang, and X. F. Zhu, *Phys. Rev. Appl.* **23**, L051001 (2025).
- 12 Y. Mou, J. Wang, H. Chen, Y. Xia, H. Li, Q. Yan, X. Jiang, Y. Wu, W. Shi, H. Jiang, et al., *Phys. Rev. Lett.* **134**, 096301 (2025).
- 13 E. Schlömann, *J. Appl. Phys.* **31**, 1647 (1960).
- 14 A. Kamra, H. Keshtgar, P. Yan, and G. E. W. Bauer, *Phys. Rev. B* **91**, 104409 (2015).
- 15 F. Vanderveken, J. Mulkers, J. Leliaert, B. Van Waeyenberge, B. Sorée, O. Zografos, F. Ciubotaru, and C. Adelman, *Phys. Rev. B* **103**, 054439 (2021).
- 16 Y. Wang, H. L. Zhang, J. L. Wu, J. Song, K. Yang, W. Qin, H. Jing, and L. M. Kuang, *Sci. China-Phys. Mech. Astron.* **66**, 110311 (2023).
- 17 H. Xiong, *Sci. China-Phys. Mech. Astron.* **68**, 250313 (2025).
- 18 M. Weiler, H. Huebl, F. S. Goerg, F. D. Czeschka, R. Gross, and S. T. B. Goennenwein, *Phys. Rev. Lett.* **108**, 176601 (2012).
- 19 L. Dreher, M. Weiler, M. Pernpeintner, H. Huebl, R. Gross, M. S. Brandt, and S. T. B. Goennenwein, *Phys. Rev. B* **86**, 134415 (2012).
- 20 T. P. Lyons, J. Puebla, K. Yamamoto, R. S. Deacon, Y. Hwang, K. Ishibashi, S. Maekawa, and Y. Otani, *Phys. Rev. Lett.* **131**, 196701 (2023).
- 21 R. Schlitz, L. Siegl, T. Sato, W. Yu, G. E. W. Bauer, H. Huebl, and S. T. B. Goennenwein, *Phys. Rev. B* **106**, 014407 (2022).
- 22 T. Wong, J. Park, K. Hayashi, M. J. Gross, R. Kim, X. Wang, S. E. Lofland, N. D. Orloff, D. B. Gopman, S. Lee, et al., *Appl. Phys. Lett.* **125**, 052402 (2024).
- 23 Y. Hwang, J. Puebla, K. Kondou, C. Gonzalez-Ballester, H. Ishiki, C. S. Muñoz, L. Liao, F. Chen, W. Luo, S. Maekawa, et al., *Phys. Rev. Lett.* **132**, 056704 (2024).
- 24 R. Verba, V. Tiberkevich, and A. Slavin, *Phys. Rev. Appl.* **12**, 054061 (2019).
- 25 T. Yu, *Phys. Rev. B* **102**, 134417 (2020).
- 26 R. Verba, I. Lisenkov, I. Krivorotov, V. Tiberkevich, and A. Slavin, *Phys. Rev. Appl.* **9**, 064014 (2018).
- 27 X. Zhang, G. E. W. Bauer, and T. Yu, *Phys. Rev. Lett.* **125**, 077203 (2020).
- 28 K. Yamamoto, W. Yu, T. Yu, J. Puebla, M. Xu, S. Maekawa, and G. Bauer, *J. Phys. Soc. Jpn.* **89**, 113702 (2020).
- 29 M. Küß, M. Hassan, Y. Kunz, A. Hörner, M. Weiler, and M. Albrecht, *Phys. Rev. B* **107**, 024424 (2023).
- 30 M. Xu, K. Yamamoto, J. Puebla, K. Baumgaertl, B. Rana, K. Miura, H. Takahashi, D. Grundler, S. Maekawa, and Y. Otani, *Sci. Adv.* **6**, eabb1724 (2020).
- 31 L. Liao, F. Chen, J. Puebla, J. Kishine, K. Kondou, W. Luo, D. Zhao, Y. Zhang, Y. Ba, and Y. Otani, *Sci. Adv.* **10**, eado2504 (2024).
- 32 Z. Zhou, W. Hu, H. Wu, M. Huang, Y. Wu, Y. Jia, W. Wang, and F. Bai, *J. Appl. Phys.* **136**, 163901 (2024).
- 33 L. Ushii, A. Slavin, and R. Verba, *Phys. Rev. Appl.* **22**, 034046 (2024).
- 34 M. Küß, M. Albrecht, and M. Weiler, *Front. Phys.* **10**, 981257 (2022).
- 35 J. Xue, H. Wu, S. Mi, M. Liu, J. Dong, J. Wang, and Q. Liu, *Phys. Rev. B* **112**, 054430 (2025).
- 36 T. Nomura, X. X. Zhang, S. Zherlitsyn, J. Wosnitzer, Y. Tokura, N. Nagaosa, and S. Seki, *Phys. Rev. Lett.* **122**, 145901 (2019).
- 37 B. Ma, Z. D. Wang, and G. Chen, *Phys. Rev. Lett.* **133**, 246604 (2024).
- 38 K. Shen, and G. E. W. Bauer, *Phys. Rev. Lett.* **115**, 197201 (2015).
- 39 R. Sasaki, Y. Nii, and Y. Onose, *Nat. Commun.* **12**, 2599 (2021).
- 40 J. Sun, S. Shi, and J. Wang, *Phys. Rev. Lett.* **130**, 256701 (2023).
- 41 S. K. Kim, D. Hill, and Y. Tserkovnyak, *Phys. Rev. Lett.* **117**, 237201 (2016).
- 42 Y. Yang, L. Zhao, D. Yi, T. Xu, Y. Chai, C. Zhang, D. Jiang, Y. Ji, D. Hou, W. Jiang, et al., *Nat. Commun.* **15**, 1018 (2024).
- 43 R. Chen, C. Chen, L. Han, P. Liu, R. Su, W. Zhu, Y. Zhou, F. Pan, and C. Song, *Nat. Commun.* **14**, 4427 (2023).
- 44 K. K. Bestha, V. Kocsis, O. Janson, Y. Shemerliuk, S. Selzer, S. Aswartham, T. Nakajima, H. Saito, B. Büchner, L. T. Corredor, et al., *Phys. Rev. B* **111**, L020409 (2025).
- 45 J. Holanda, D. S. Maior, A. Azevedo, and S. M. Rezende, *Nat. Phys.* **14**, 500 (2018).
- 46 H. Zhang, N. Peshcherenko, F. Yang, T. Z. Ward, P. Raghuvanshi, L. Lindsay, C. Felser, Y. Zhang, J. Q. Yan, and H. Miao, *Nat. Phys.* **21**, 1387 (2025).
- 47 T. Sato, W. Yu, S. Streib, and G. E. W. Bauer, *Phys. Rev. B* **104**, 014403 (2021).
- 48 J. Ren, *Chin. Phys. Lett.* **39**, 126301 (2022).
- 49 Y. Long, J. Ren, and H. Chen, *Proc. Natl. Acad. Sci. U.S.A.* **115**, 9951 (2018).
- 50 C. Shi, R. Zhao, Y. Long, S. Yang, Y. Wang, H. Chen, J. Ren, and X. Zhang, *Natl. Sci. Rev.* **6**, 707 (2019).
- 51 K. Y. Bliokh, and F. Nori, *Phys. Rev. B* **99**, 020301 (2019).
- 52 K. Y. Bliokh, *Phys. Rev. Lett.* **129**, 204303 (2022).
- 53 W. Yuan, C. Yang, D. Zhang, Y. Long, Y. Pan, Z. Zhong, H. Chen, J. Zhao, and J. Ren, *Nat. Commun.* **12**, 6954 (2021).
- 54 M. M. Sonner, F. Khosravi, L. Janker, D. Rudolph, G. Koblmüller, Z. Jacob, and H. J. Krenner, *Sci. Adv.* **7**, eabf7414 (2021).
- 55 C. Yang, D. Zhang, J. Zhao, W. Gao, W. Yuan, Y. Long, Y. Pan, H. Chen, F. Nori, K. Y. Bliokh, et al., *Phys. Rev. Lett.* **131**, 136102 (2023).
- 56 J. Zhao, C. Yang, W. Yuan, D. Zhang, Y. Long, Y. Pan, H. Chen, Z. Zhong, and J. Ren, *Phys. Rev. Lett.* **129**, 275501 (2022).
- 57 C. Yang, and J. Ren, *Proc. Natl. Acad. Sci. U.S.A.* **121**, e2411427121 (2024).
- 58 G. E. W. Bauer, P. Tang, M. Elyasi, Y. M. Blanter, and B. J. van Wees, *Phys. Rev. B* **108**, 064431 (2023).
- 59 R. R. Neumann, A. Mook, J. Henk, and I. Mertig, *Phys. Rev. Lett.* **125**, 117209 (2020).
- 60 X. Chen, Y. Liu, P. Liu, Y. Yu, J. Ren, J. Li, A. Zhang, and Q. Liu, *Nature* **640**, 349 (2025).
- 61 A. Mook, R. Hoyer, J. Klinovaja, and D. Loss, *Phys. Rev. B* **107**, 064429 (2023).
- 62 Z. X. Li, Z. Wang, Y. Cao, and P. Yan, *Phys. Rev. B* **105**, 174433 (2022).
- 63 R. Cheng, S. Okamoto, and D. Xiao, *Phys. Rev. Lett.* **117**, 217202 (2016).
- 64 J. W. Tucker and V. W. Rampton. Microwave ultrasonics in solid state physics (North-Holland Publishing Company, Amsterdam, 1972).
- 65 C. Kittel, *Phys. Rev.* **73**, 155 (1948).
- 66 T. L. Gilbert, *IEEE Trans. Magn.* **40**, 3443 (2004).
- 67 J. Lan, W. Yu, R. Wu, and J. Xiao, *Phys. Rev. X* **5**, 041049 (2015).
- 68 L. Dreher, M. Weiler, M. Pernpeintner, H. Huebl, R. Gross, M. S. Brandt, and S. T. B. Goennenwein, *Phys. Rev. B* **98**, 099901 (2018).
- 69 M. Harder, L. H. Bai, C. Match, J. Sirker, and C. M. Hu, *Sci. China-Phys. Mech. Astron.* **59**, 117511 (2016).
- 70 B. Z. Rameshti, S. V. Kusminskiy, J. A. Haigh, K. Usami, D. Lachance-Quirion, Y. Nakamura, C. M. Hu, H. X. Tang, G. E. W. Bauer, and Y. M. Blanter, *Phys. Rep.* **979**, 1 (2022).
- 71 K. Y. Bliokh, A. Y. Bekshaev, and F. Nori, *Nat. Commun.* **5**, 3300 (2014).
- 72 K. Y. Bliokh, D. Smirnova, and F. Nori, *Science* **348**, 1448 (2015).
- 73 K. Y. Bliokh, F. J. Rodríguez-Fortuño, F. Nori, and A. V. Zayats, *Nat. Photon.* **9**, 796 (2015).
- 74 J. Zhang, W. Yu, X. Chen, and J. Xiao, *AIP Adv.* **13**, 055108 (2023).
- 75 L. D. Landau, and E. M. Lifshits. Course of theoretical physics vol 7: Theory and elasticity (Pergamon press, Oxford, 1959).
- 76 G. J. Chaplain, J. M. De Ponti, and R. V. Craster, *Phys. Rev. Lett.* **128**, 064301 (2022).
- 77 L. D. Landau, and E. M. Lifshitz. Mechanics: Volume 1 (Course of

- Theoretical Physics Series) (Pergamon Press, Oxford, 2000).
- 78 A. G. Whitney, J. M. Lewis, J. Dickovick, V. Kalappattil, L. D. Carr, and M. Wu, *Phys. Rev. Appl.* **23**, 064012 (2025).
- 79 D. Richardson, B. A. Kalinikos, L. D. Carr, and M. Wu, *Phys. Rev. Lett.* **121**, 107204 (2018).
- 80 R. Cheng, M. W. Daniels, J. G. Zhu, and D. Xiao, *Sci. Rep.* **6**, 24223 (2016).
- 81 H. Ling, Y. Wu, X. Ge, Z. Ye, T. Dong, L. Miao, W. Luo, Y. Zhang, and Q. Fu, *J. Appl. Phys.* **136**, 183906 (2024).
- 82 K. An, A. N. Litvinenko, R. Kohno, A. A. Fuad, V. V. Naletov, L. Vila, U. Ebels, G. de Loubens, H. Hurdequint, N. Beaulieu, et al., *Phys. Rev. B* **101**, 060407 (2020).
- 83 S. Streib, H. Keshtgar, and G. E. W. Bauer, *Phys. Rev. Lett.* **121**, 027202 (2018).
- 84 H. Matsumoto, Y. Todaka, T. Kawada, M. Kawaguchi, and M. Hayashi, *Phys. Rev. Appl.* **22**, 054052 (2024).
- 85 P. Kuszewski, J. Y. Duquesne, L. Becerra, A. Lemaître, S. Vincent, S. Majrab, F. Margaillan, C. Gourdon, and L. Thevenard, *Phys. Rev. Appl.* **10**, 034036 (2018).
- 86 A. Kamra, T. Wimmer, H. Huebl, and M. Althammer, *Phys. Rev. B* **102**, 174445 (2020).
- 87 L. Liensberger, A. Kamra, H. Maier-Flaig, S. Geprägs, A. Erb, S. T. B. Goennenwein, R. Gross, W. Belzig, H. Huebl, and M. Weiler, *Phys. Rev. Lett.* **123**, 117204 (2019).