

Random Vectors

Ricardo Dahis

PUC-Rio, Department of Economics

Summer 2023

Contents

1. Joint and marginal distributions
2. Conditional distribution and independence
3. Bivariate transformations
4. Hierarchical models, mixtures and a LIE
5. Covariance and correlation
6. Multivariate distributions
7. Inequalities
8. Exercises

Contents

1. Joint and marginal distributions
2. Conditional distribution and independence
3. Bivariate transformations
4. Hierarchical models, mixtures and a LIE
5. Covariance and correlation
6. Multivariate distributions
7. Inequalities
8. Exercises

random vector

- **definition:** an n -dimensional random vector is a function from the sample space S into the n -dimensional Euclidean space \mathbb{R}^n
- **example:** consider the experiment of tossing two fair dice, and let X and Y denote the sum of the two dice and the absolute difference of the two dice, respectively

$$\mathbb{P}(X = 5, Y = 3) = \mathbb{P}(\{(1, 4), (4, 1)\}) = \frac{2}{36} = \frac{1}{18}$$

- **definition:** let (X, Y) denote a discrete bivariate random vector, then the joint pmf $f_{X,Y}(x, y)$ from \mathbb{R}^2 into \mathbb{R} is given by $f(x, y) = \mathbb{P}(X = x, Y = y)$
- we can now discuss probabilities of events defined in terms of (X, Y) .

joint pmf

- the joint pmf completely characterizes the probability distribution of a random vector (X, Y) just as in the univariate case

$$\mathbb{P}((X, Y) \in A) = \sum_{(x,y) \in A} f_{X,Y}(x, y)$$

- expectations are defined

$$\mathbb{E}[g(X, Y)] = \sum_{(x,y) \in \mathbb{R}^2} g(x, y) f_{X,Y}(x, y)$$

- fortunately, the expectation operator continues to have the same properties as before; in particular

$$\mathbb{E}[a g(X, Y) + b h(X, Y) + c] = a \mathbb{E}[g(X, Y)] + b \mathbb{E}[h(X, Y)] + c$$

properties of joint pdfs

joint pmf satisfies the usual properties (**verify**), namely

(i) $f_{X,Y}(x, y) \geq 0$ for any (x, y)

(ii) $\sum_{(x,y) \in \mathbb{R}^2} f_{X,Y}(x, y) = 1$

and thus it is a well-defined probability distribution.

marginal pmfs

- there may be events, probabilities, moments or expectations that involve only one of the random variables in the vector.
- **theorem** (CB 4.1.6): let (X, Y) denote a discrete bivariate random vector with joint pmf $f_{X,Y}(x, y)$, then the **marginal pmfs** of X and Y are respectively

$$f_X(x) = \mathbb{P}(X = x) = \sum_{y \in \mathbb{R}} f_{X,Y}(x, y)$$

$$f_Y(y) = \mathbb{P}(Y = y) = \sum_{x \in \mathbb{R}} f_{X,Y}(x, y)$$

we use the subscript X in $f_X(x)$ to emphasize the distinction from $f_{X,Y}(x, y)$.

same marginals, different joint pmfs

- same marginal pmfs $\not\Rightarrow$ same joint pmfs.
- counterexample: define

$$\begin{aligned}f_{X,Y}(0,0) &= f_{X,Y}(0,1) = \frac{1}{6} \\f_{X,Y}(1,0) &= f_{X,Y}(1,1) = \frac{1}{3} \\f_{X,Y}(x,y) &= 0 \text{ for any other } (x,y)\end{aligned}$$

the marginals are

$$\begin{aligned}f_X(0) &= \frac{1}{3}, & f_X(1) &= \frac{2}{3} \\f_Y(0) &= \frac{1}{2}, & f_Y(1) &= \frac{1}{2}\end{aligned}$$

same marginals, different joint pmfs

- counterexample (cont'd): now define

$$\begin{aligned}g_{XY}(0,0) &= \frac{1}{12} & g_{XY}(0,1) &= \frac{3}{12} \\g_{XY}(1,0) &= \frac{5}{12} & g_{XY}(1,1) &= \frac{3}{12} \\g_{XY}(x,y) &= 0 \text{ for any other } (x,y)\end{aligned}$$

the marginals are

$$\begin{aligned}g_X(0) &= \frac{1}{3}, & g_X(1) &= \frac{2}{3} \\g_Y(0) &= \frac{1}{2}, & g_Y(1) &= \frac{1}{2}\end{aligned}$$

- $f_X(0) = g_X(0)$, $f_X(1) = g_X(1)$, $f_Y(0) = g_Y(0)$, $f_Y(1) = g_Y(1)$ but $f_{X,Y}(x,y) \neq g_{XY}(x,y)$.
- intuitive since marginals contain less information than joint pmfs.

joint and marginal pdfs

- **definition:** a function $f_{X,Y}(x,y)$ from \mathbb{R}^2 into \mathbb{R} is the joint pdf of the continuous bivariate random vector (X, Y) if, for every $A \subset \mathbb{R}^2$,

$$\mathbb{P}((X, Y) \in A) = \iint_A f_{X,Y}(x,y) dx dy$$

- the joint pdf is such that $f_{X,Y}(x,y) \geq 0$ for all $(x,y) \in \mathbb{R}^2$ and that $\int_{-\infty}^{\infty} \int_{-\infty}^{\infty} f_{X,Y}(x,y) dx dy = 1$
- expectations are just like in the discrete case, but with integrals

$$\mathbb{E}[g(X, Y)] = \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} g(x,y) f_{X,Y}(x,y) dx dy$$

- **definition:** the marginal pdfs are given by (you can also verify that this distribution is proper)

$$f_X(x) = \int_{-\infty}^{\infty} f_{X,Y}(x,y) dy, \quad -\infty < x < \infty$$

$$f_Y(y) = \int_{-\infty}^{\infty} f_{X,Y}(x,y) dx, \quad -\infty < y < \infty$$

example

- **example:** let (X, Y) denote a continuous bivariate random vector with joint pdf $f_{X,Y}(x, y) = 6xy^2$ for (x, y) in the unit square and zero otherwise.

$$\begin{aligned}\int_{-\infty}^{\infty} \int_{-\infty}^{\infty} f_{X,Y}(x, y) \, dx \, dy &= \int_0^1 \int_0^1 6xy^2 \, dx \, dy \\ &= \int_0^1 (3x^2y^2)_0^1 \, dy = \int_0^1 3y^2 \, dy = (y^3)_0^1 = 1\end{aligned}$$

$$f_X(x) = \int_{-\infty}^{\infty} f_{X,Y}(x, y) \, dy = \int_0^1 6xy^2 \, dy = 6x (y^3/3)_0^1 = 2x$$

$$f_Y(y) = \int_{-\infty}^{\infty} f_{X,Y}(x, y) \, dx = \int_0^1 6xy^2 \, dx = 6y^2 (x^2/2)_0^1 = 3y^2$$

example

- example (cont'd): let (X, Y) denote a continuous bivariate random vector with joint pdf $f_{X,Y}(x, y) = 6xy^2$ for (x, y) in the unit square and zero otherwise.

- Consider now calculating the probability that $X + Y \geq 1$.
- The region over which we integrate is

$$\begin{aligned} A &= \{(x, y) : x + y \geq 1, 0 < x < 1, 0 < y < 1\} \\ &= \{(x, y) : x \geq 1 - y, 0 < x < 1, 0 < y < 1\} \\ &= \{(x, y) : 1 - y \leq x < 1, 0 < x < 1, 0 < y < 1\} \end{aligned}$$

- So

$$\mathbb{P}(X + Y \geq 1) = \int_0^1 \int_{1-y}^1 6xy^2 dx dy = 0.9$$


a more complicated example

- example 2: let (X, Y) denote a continuous bivariate random vector with joint pdf $f_{X,Y}(x, y) = e^{-y}$ for $0 < x < y < \infty$.

$$\begin{aligned}\mathbb{P}(X + Y \geq 1) &= 1 - \mathbb{P}(X + Y < 1) \\ &= 1 - \int_0^{1/2} \int_x^{1-x} e^{-y} dy dx \\ &= 1 - \int_0^{1/2} \left(e^{-x} - e^{-(1-x)} \right) dx \\ &= 1 - \left(-e^{-\frac{1}{2}} + e^0 - e^{-\frac{1}{2}} + e^{-1} \right) \\ &= 2e^{-1/2} - e^{-1}\end{aligned}$$

given that $\Omega_{XY} = \{(x, y) : x + y \geq 1, 0 < x < y < \infty\}$ is the unbounded region with three sides given by $x = y$, $x + y = 1$, and $x = 0$

regions from the example



joint cdf

- the joint probability distribution of (X, Y) is also completely described with the joint cdf $F_{X,Y}(x, y) = \mathbb{P}(X \leq x, Y \leq y)$ for all $(x, y) \in \mathbb{R}^2$
- characterization**: not very handy for discrete random vectors, but extremely useful for continuous random vectors given that

$$F_{X,Y}(x, y) = \int_{-\infty}^y \int_{-\infty}^x f_{X,Y}(u, v) \, du \, dv$$

and hence, by the fundamental theorem of calculus,

$$\frac{\partial^2 F_{X,Y}(x, y)}{\partial x \partial y} = f_{X,Y}(x, y)$$

at any continuity point of $f_{X,Y}(x, y)$

Contents

1. Joint and marginal distributions
- 2. Conditional distribution and independence**
3. Bivariate transformations
4. Hierarchical models, mixtures and a LIE
5. Covariance and correlation
6. Multivariate distributions
7. Inequalities
8. Exercises

conditional probability

- **definition:** let (X, Y) denote a discrete bivariate random vector with joint pmf $f_{X,Y}(x, y)$ and marginals $f_X(x)$ and $f_Y(y)$, then the conditional pmf of Y given $X = x$ is

$$f_{Y|X}(y|x) = \mathbb{P}(Y = y|X = x) = \frac{f_{X,Y}(x, y)}{f_X(x)}$$

for any x such that $f_X(x) = \mathbb{P}(X = x) > 0$

- just checking to be on the safe side...
 - (i) $f_{Y|X}(y|x) \geq 0$ for every y given that $f_{X,Y}(x, y) \geq 0$ and $f_X(x) > 0$
 - (ii) $\sum_y f_{Y|X}(y|x) = \frac{\sum_y f_{X,Y}(x, y)}{f_X(x)} = \frac{f_X(x)}{f_X(x)} = 1$

continuous random variables

- if X and Y are continuous random variables, then $\mathbb{P}(X = x) = 0$ for every value of x and hence we cannot divide the joint probability by the probability of the conditioning event
- **however**, we may still define the conditional probability of Y given $X = x$ analogously to the discrete case with pdfs replacing pmfs
- **definition**: let (X, Y) be a continuous bivariate random vector with joint pdf $f_{X,Y}(x, y)$ and marginals $f_X(x)$ and $f_Y(y)$, then the conditional pdf of Y given $X = x$ is

$$f_{Y|X}(y|x) = \frac{f_{X,Y}(x, y)}{f_X(x)}$$

for any x such that $f_X(x) > 0$. Analogously,

$$f_{X|Y}(x|y) = \frac{f_{X,Y}(x, y)}{f_Y(y)}$$

for any y such that $f_Y(y) > 0$.

conditional expectation

- conditional pdfs/pmfs are useful not only to compute conditional probabilities, but also to calculate conditional expectations

$$\mathbb{E}[g(Y)|X = x] = \begin{cases} \sum_y g(y)f_{Y|X}(y|x) & \text{if discrete} \\ \int_{-\infty}^{\infty} g(y)f_{Y|X}(y|x) dy & \text{if continuous} \end{cases}$$

- the conditional expectation satisfies all the properties of the usual expectation operator
- in particular, $\mathbb{E}(Y|X)$ provides the best guess at Y based on knowledge of X in a MSE sense (you can try to show this!)
- note that $f_{Y|X}(y|x)$ is function of x . So we really have a family of distributions, one for each x , possibly with different $\mathbb{E}(Y|X = x)$.
 - the notation $Y|X$ describes the entire family of distributions.

interesting example

- **example:** let (X, Y) have a joint pdf $f_{X,Y}(x, y) = e^{-y}$ for $0 < x < y < \infty$, then

$$f_X(x) = \int_{-\infty}^{\infty} f_{X,Y}(x, y) dy = \int_x^{\infty} e^{-y} dy = e^{-x}$$

$$f_{Y|X}(y|x) = \frac{f_{X,Y}(x, y)}{f_X(x)} = e^{-(y-x)} \text{ for } y > x$$

and hence $X \sim \text{Exp}(1)$ and $Y|X = x$ is also exponential with location parameter x

$$\mathbb{E}(Y|X = x) = \int_x^{\infty} y f_{Y|X}(y|x) dy = \int_x^{\infty} y e^{-(y-x)} dy = 1 + x$$

$$\begin{aligned} \text{var}(Y|X = x) &= \mathbb{E}(Y^2|X = x) - [\mathbb{E}(Y|X = x)]^2 \\ &= \int_x^{\infty} y^2 e^{-(y-x)} dy - (1 + x)^2 = 1 \end{aligned}$$

but why is it interesting?

- conditional variance does not depend on x , but does that mean it is equal to the unconditional variance?

$$f_Y(y) = \int_{-\infty}^{\infty} f_{X,Y}(x,y) dx = \int_0^y e^{-y} dx = ye^{-y}$$

remember: the gamma distribution is given by

$$f(x|\alpha, \beta) = \frac{1}{\Gamma(\alpha)\beta^\alpha} x^{\alpha-1} e^{-\frac{x}{\beta}}, \quad \text{for}$$

for $0 < t < \infty$, $\alpha, \beta > 0$ and $\Gamma(\alpha) = (\alpha - 1)!$. Hence $Y \sim G(\alpha, \beta)$, with $\alpha = 2$ and $\beta = 1$, implying that $\text{var}(Y) = \alpha\beta^2 = 2$.

- even though the conditional variance does not depend on the value of x , knowledge of the latter considerably reduces the variability of Y

$$\text{var}(Y|X = x) = c \not\Rightarrow \text{var}(Y) = c$$

- we will come back to this point later

independence

- $\mathbb{E}[g(Y)|X]$ is a random variable whose values typically depend on the value of X , unless independent ($X \perp\!\!\!\perp Y$)
- **definition:** let (X, Y) denote a bivariate random vector with joint pdf/pmf $f_{X,Y}(x, y)$ and marginals $f_X(x)$ and $f_Y(y)$, then X and Y are independent if

$$f_{X,Y}(x, y) = f_X(x)f_Y(y)$$

- This immediately implies that $f_{Y|X}(y|x) = f_Y(y)$, since

$$f_{Y|X}(y|X=x) = \frac{f_{X,Y}(x, y)}{f_X(x)} = \frac{f_X(x)f_Y(y)}{f_X(x)} = f_Y(y)$$

and the knowledge of x does not inform the distribution of Y

a note of caution

- two pdfs that differ only a zero-measure set define the same probability distribution for (X, Y) .
- so definition may fail to hold on sets with measure zero. But in this case X and Y are still independent.
- to see this, take $f_{X,Y}(x, y)$ and $f_{X,Y}^*(x, y)$ equal everywhere except on A for which $\int_A \int dx dy = 0$.
- let (X, Y) have pdf $f_{X,Y}(x, y)$, (X^*, Y^*) have pdf $f_{X,Y}^*(x, y)$, and $B \subset \mathbb{R}^2$. Then

$$\begin{aligned} P((X, Y) \in B) &= \int_B \int f(x, y) dx dy = \int_{B \cap A^c} \int f(x, y) dx dy \\ &= \int_{B \cap A^c} \int f^*(x, y) dx dy = \int_B \int f^*(x, y) dx dy \\ &= P((X^*, Y^*) \in B) \end{aligned}$$

- for example, take $f_{X,Y}(x, y) = e^{-(x+y)}$ with $x, y > 0$, describing two independent exponential random variables.
- and take $f_{X,Y}^*(x, y) = f_{X,Y}(x, y)$ except that $f_{X,Y}^*(x, y) = 0$ if $x = y$ in $A = \{(x, x), x > 0\}$.

independence

- **theorem** (CB 4.2.7): let (X, Y) be a bivariate random vector with joint pdf $f(x, y)$. Then X and Y are independent if, and only if, there exist functions $g(x)$ and $h(y)$ such that, for every $x \in \mathbb{R}$ and $y \in \mathbb{R}$,

$$f(x, y) = g(x)h(y)$$

- **proof** (\Rightarrow): trivial setting $g(x) = f_X(x)$ and $h(y) = f_Y(y)$.
- **proof** (\Leftarrow): suppose that $f(x, y) = g(x)h(y)$ and define

$$\int_{-\infty}^{\infty} g(x) dx = c \text{ and } \int_{-\infty}^{\infty} h(y) dy = d$$

so cd satisfies

$$\begin{aligned} cd &= \left(\int_{-\infty}^{\infty} g(x) dx \right) \left(\int_{-\infty}^{\infty} h(y) dy \right) \\ &= \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} g(x)h(y) dx dy \\ &= \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} f(x, y) dx dy = 1 \end{aligned}$$

independence

- proof (\Leftarrow) (cont'd): the marginals are given by

$$f_X(x) = \int_{-\infty}^{\infty} g(x)h(y) dy = g(x)d$$

$$f_Y(y) = \int_{-\infty}^{\infty} g(x)h(y) dx = h(y)c$$

\Downarrow

$$f(x, y) = g(x)h(y) = g(x)h(y)cd = f_X(x)f_Y(y)$$

establishing the desired result. ■

- **example:** Consider $f(x, y) = \frac{1}{384}x^2y^4e^{-y-\frac{x}{2}}$ with $x, y > 0$ and

$$g(x) = \begin{cases} x^2e^{-x/2} & x > 0 \\ 0 & x \leq 0 \end{cases} \quad \text{and} \quad h(y) = \begin{cases} \frac{1}{384}y^4e^{-y} & y > 0 \\ 0 & y \leq 0 \end{cases}$$

by theorem above, it follows immediately that X and Y are independent.

independence and support of the joint pdf

- the **support set** matters: independence can be ruled out in simple cases.
- denote the support of the marginals as $A = \{x : f_X(x) > 0\}$ and $B = \{y : f_Y(y) > 0\}$
- if X and Y independent, then $f(x, y) = f_X(x)f_Y(y) > 0$ on the set $\{(x, y) : x \in A, y \in B\}$
 - define $A \times B = \{(x, y) : x \in A, y \in B\}$, denoted **cross-product set**
 - if the set $\{(x, y) : f(x, y) > 0\}$ is not a cross-product, X and Y **cannot be independent**.
 - in one of the examples above, we have support set $0 < x < y < \infty$, so not only $0 < x, y < \infty$ but also $x < y$, so not independent.

independence and support of the joint pdf

- **theorem** (CB 4.2.10): let X and Y be independent variables
 - (i) for any $A \subset \mathbb{R}$ and $B \subset \mathbb{R}$, $\mathbb{P}(X \in A, Y \in B) = \mathbb{P}(X \in A)\mathbb{P}(Y \in B)$. That is, the events $\{X \in A\}$ and $\{Y \in B\}$ are independent
 - (ii) let $g(x)$ be a function of x and $h(y)$ be a function of y . Then

$$\mathbb{E}(g(X)h(Y)) = \mathbb{E}(g(X))\mathbb{E}(h(Y))$$

- **proof** (ii): Notice that

$$\begin{aligned}\mathbb{E}(g(X)h(Y)) &= \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} g(x)h(y)f(x, y) \, dx \, dy \\ &= \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} g(x)h(y)f_X(x)f_Y(y) \, dx \, dy \\ &= \mathbb{E}(g(X))\mathbb{E}(h(Y))\end{aligned}$$

independence and support of the joint pdf

- **proof (i)**: Set $g(X) = 1(x \in A)$, $h(Y) = 1(y \in B)$. Notice that

$$\mathbb{P}(X \in A) = \mathbb{E}[1(x \in A)] = \int_{-\infty}^{\infty} \mathcal{I}_A(x) f_X(x) dx$$

$$\mathbb{P}(Y \in B) = \mathbb{E}[1(y \in B)] = \int_{-\infty}^{\infty} \mathcal{I}_B(y) f_Y(y) dy$$

$$\mathbb{P}(X \in A, Y \in B) = \mathbb{E}[\mathcal{I}_{AB}(x, y)] = \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} \mathcal{I}_{AB}(x, y) f_X(x) f_Y(y) dx dy$$

and apply (ii). ■

independence and support of the joint pdf

- **theorem** (CB 4.2.12): let X and Y be independent random variables with moment generating functions $M_X(t)$ and $M_Y(t)$. Then the mgf of $Z = X + Y$ is

$$M_Z(t) = M_X(t)M_Y(t)$$

- **proof:**

$$M_Z(t) = \mathbb{E}\left(e^{tZ}\right) = \mathbb{E}\left(e^{t(X+Y)}\right) = \left(\mathbb{E}e^{tX}\right)\left(\mathbb{E}e^{tY}\right) = M_X(t)M_Y(t)$$



independence and support of the joint pdf

- **example/corollary** (CB 4.2.14): let $X \sim N(\mu, \sigma^2)$ and $Y \sim N(\gamma, \tau^2)$, independent. Then $Z = X + Y \sim N(\mu + \gamma, \sigma^2 + \tau^2)$.
- **proof**: X and Y have mgf representations

$$M_X(t) = e^{\mu t + \sigma^2 t^2 / 2}$$

$$M_Y(t) = e^{\gamma t + \tau^2 t^2 / 2}$$

then

$$M_Z(t) = e^{(\mu + \gamma)t + (\sigma^2 + \tau^2)t^2 / 2}$$

which is the mgf of a normal random variable with mean $\mu + \gamma$ and variance $\sigma^2 + \tau^2$ ■

Contents

1. Joint and marginal distributions
2. Conditional distribution and independence
- 3. Bivariate transformations**
4. Hierarchical models, mixtures and a LIE
5. Covariance and correlation
6. Multivariate distributions
7. Inequalities
8. Exercises

discrete random vectors

- let (X, Y) be a bivariate random vector with known probability distribution.
- Consider a new bivariate random vector (U, V) such that $U = g_1(X, Y)$ and $V = g_2(X, Y)$
 - $(U, V) \in B \Leftrightarrow (X, Y) \in A, A = \{(x, y) : (g_1(x, y), g_2(x, y)) \in B\}$
 - $\mathbb{P}((U, V) \in B) = \mathbb{P}((X, Y) \in A)$
 - keeping track of the support: from $\Omega_{X,Y} = \{(x, y) : f_{X,Y}(x, y) > 0\}$ to
$$\Omega_{U,V} = \{(u, v) : u = g_1(x, y), v = g_2(x, y) \text{ for some } (x, y) \in \Omega_{X,Y}\}$$
 - In the discrete case,

$$\begin{aligned} f_{UV}(u, v) &= \mathbb{P}(U = u, V = v) = \mathbb{P}\left((X, Y) \in \Omega_{X,Y}^{(uv)}\right) \\ &= \sum_{(x,y) \in \Omega_{X,Y}^{(uv)}} f_{X,Y}(x, y) \end{aligned}$$

where $\Omega_{X,Y}^{uv} = \{(x, y) \in \Omega_{X,Y} : g_1(x, y) = u, g_2(x, y) = v\}$.

sum of Poisson variables

example: let X and Y be independent Poisson random variables with joint pmf given by $f_{X,Y}(x,y) = \frac{e^{-\theta}\theta^x}{x!} \frac{e^{-\lambda}\lambda^y}{y!}$. also let $U = X + Y$ and $V = Y$.

- the support of the Poisson is $\Omega_{X,Y} = \{(x,y) : x \in \mathbb{N}, y \in \mathbb{N}\}$
- then $\Omega_{U,V} = \{(u,v) : v = 0, 1, 2, \dots \text{ and } u = v, v+1, v+2, \dots\}$
- $\Omega_{X,Y}^{(uv)}$ consists of only the single point $(u-v, v)$ and

$$f_{U,V}(u,v) = f_{X,Y}(u-v, v) = \frac{e^{-\theta}\theta^{u-v}}{(u-v)!} \frac{e^{-\lambda}\lambda^v}{v!}$$

- theorem/application: $X \sim P(\theta)$, $Y \sim P(\lambda)$ and $X \perp Y \Rightarrow X + Y \sim P(\theta + \lambda)$

$$\begin{aligned} f_U(u) &= \sum_{v=0}^u \frac{e^{-\theta}\theta^{u-v}}{(u-v)!} \frac{e^{-\lambda}\lambda^v}{v!} = e^{-(\theta+\lambda)} \sum_{v=0}^u \frac{\theta^{u-v}\lambda^v}{v!(u-v)!} \\ &= \frac{e^{-(\theta+\lambda)}}{u!} \sum_{v=0}^u \binom{u}{v} \theta^{u-v}\lambda^v = \frac{e^{-(\theta+\lambda)}}{u!} (\theta + \lambda)^u \end{aligned}$$

binomial theorem is used in the last equality: $(x+y)^n = \sum_{k=0}^n \binom{n}{k} x^{n-k} y^k$.

continuous random vector

- let X and Y be continuous random variables with joint pdf $f_{X,Y}(x, y)$.
- as before, the support set $\Omega_{X,Y} = \{(x, y) : f_{X,Y}(x, y) > 0\}$ maps into

$$\Omega_{U,V} = \{(u, v) : u = g_1(x, y), v = g_2(x, y) \text{ for some } (x, y) \in \Omega_{X,Y}\}$$

- for now, assume that transformation $g : \Omega_{X,Y} \rightarrow \Omega_{U,V}$ is bijective: for each $(u, v) \in \Omega_{U,V}$ there is only one pair $(x, y) \in \Omega_{X,Y}$.
- we can solve the inverse transformation $x = h_1(u, v)$ and $y = h_2(u, v)$.

continuous random vector

- **theorem:** the pdf of (U, V) is given by

$$f_{U,V}(u, v) = f_{X,Y}(h_1(u, v), h_2(u, v)) \cdot |J|$$

where J is the Jacobian of the transformation

$$J = \begin{vmatrix} \frac{\partial x}{\partial u} & \frac{\partial x}{\partial v} \\ \frac{\partial y}{\partial u} & \frac{\partial y}{\partial v} \end{vmatrix} = \frac{\partial x}{\partial u} \frac{\partial y}{\partial v} - \frac{\partial x}{\partial v} \frac{\partial y}{\partial u}$$

$x = h_1(u, v)$, $y = h_2(u, v)$ and $|\cdot|$ is the determinant.

- the term $|J|$ gives a "magnification factor" for area in going from u - v coordinates to x - y coordinates, just like in the univariate case.
- **intuition for proof:** draw rectangles in both coordinates and compute equivalent areas accounting for magnification.

product of betas

- **example** (CB 4.3.3): we want to find the distribution of the **product** of independent betas $X \sim B(\alpha, \beta)$ and $Y \sim B(\alpha + \beta, \gamma)$.
- each $B(\alpha, \beta)$ distribution is given by

$$f(x|\alpha, \beta) = \frac{\Gamma(\alpha + \beta)}{\Gamma(\alpha)\Gamma(\beta)} x^{\alpha-1} (1-x)^{\beta-1}$$

with $0 < x < 1$. So the joint distribution of X and Y is

$$f_{X,Y}(x, y) = \frac{\Gamma(\alpha + \beta + \gamma)}{\Gamma(\alpha)\Gamma(\beta)\Gamma(\gamma)} x^{\alpha-1} (1-x)^{\beta-1} y^{\alpha+\beta-1} (1-y)^{\gamma-1}$$

- we really don't care about V , but we choose one such that the mapping is bijective: let $U = XY$ and $V = X$, then $\Omega_{U,V} = \{(u, v) : 0 < u < v < 1\}$
- then we obtain the marginal for U to get the final answer.

product of betas

- so

$$\begin{aligned}f_{U,V}(u, v) &= f_{X,Y}(v, u/v) \left| \frac{\partial x}{\partial u} \frac{\partial y}{\partial v} - \frac{\partial y}{\partial u} \frac{\partial x}{\partial v} \right| \\&= f_{X,Y}(v, u/v) |0(-u/v^2) - 1(1/v)| = \frac{1}{v} f_{X,Y}(v, u/v) \\&= \frac{\Gamma(\alpha + \beta + \gamma)}{\Gamma(\alpha)\Gamma(\beta)\Gamma(\gamma)} v^{\alpha-2}(1-v)^{\beta-1} \left(\frac{u}{v}\right)^{\alpha+\beta-1} \left(1 - \frac{u}{v}\right)^{\gamma-1} \\&= \frac{\Gamma(\alpha + \beta + \gamma)}{\Gamma(\alpha)\Gamma(\beta)\Gamma(\gamma)} u^{\alpha-1} \left(\frac{u}{v} - u\right)^{\beta-1} \left(1 - \frac{u}{v}\right)^{\gamma-1} \frac{u}{v^2}\end{aligned}$$

marginal is also beta

- Taking the marginal for U ,

$$\begin{aligned}f_U(u) &= \int_u^1 f_{U,V}(u, v) \, dv \\&= \frac{\Gamma(\alpha + \beta + \gamma)}{\Gamma(\alpha)\Gamma(\beta)\Gamma(\gamma)} u^{\alpha-1} \int_u^1 \left(\frac{u}{v} - u\right)^{\beta-1} \left(1 - \frac{u}{v}\right)^{\gamma-1} \frac{u}{v^2} \, dv \\&= \frac{\Gamma(\alpha + \beta + \gamma)}{\Gamma(\alpha)\Gamma(\beta)\Gamma(\gamma)} u^{\alpha-1} (1-u)^{\beta+\gamma-1} \\&\quad \times \int_u^1 \left(\frac{u/v - u}{1-u}\right)^{\beta-1} \left(\frac{1-u/v}{1-u}\right)^{\gamma-1} \frac{u}{v^2(1-u)} \, dv \\&= \frac{\Gamma(\alpha + \beta + \gamma)}{\Gamma(\alpha)\Gamma(\beta)\Gamma(\gamma)} u^{\alpha-1} (1-u)^{\beta+\gamma-1} \int_0^1 z^{\beta-1} (1-z)^{\gamma-1} \, dz\end{aligned}$$

defining

$$z = \frac{u/v - u}{1-u} \Rightarrow dz = -\frac{u}{v^2(1-u)} dv$$

marginal is also beta

- so

$$\begin{aligned} f_U(u) &= \frac{\Gamma(\alpha + \beta + \gamma)}{\Gamma(\alpha)\Gamma(\beta)\Gamma(\gamma)} u^{\alpha-1}(1-u)^{\beta+\gamma-1} \underbrace{\int_0^1 z^{\beta-1}(1-z)^{\gamma-1} dz}_{=\frac{\Gamma(\beta)\Gamma(\gamma)}{\Gamma(\beta+\gamma)}} \\ &= \frac{\Gamma(\alpha + \beta + \gamma)}{\Gamma(\alpha)\Gamma(\beta + \gamma)} u^{\alpha-1}(1-u)^{\beta+\gamma-1} \quad U \sim B(\alpha, \beta + \gamma) \end{aligned}$$

- where the last identity comes from recognizing the integrand as the kernel of a Beta pdf and using CB 3.3.17.

sum and difference of standard normals

- **example** (CB 4.3.4): let $X \sim N(0, 1)$ and $Y \sim N(0, 1)$ be independent, then $U = X + Y$ and $V = X - Y$ are also normal random variables

$$\begin{aligned}f_{U,V}(u, v) &= f_{X,Y}\left(\frac{u+v}{2}, \frac{u-v}{2}\right) \left| \frac{1}{2} \begin{pmatrix} 1 & 1 \\ 1 & -1 \end{pmatrix} \right| = \frac{1}{2} f_{X,Y}\left(\frac{u+v}{2}, \frac{u-v}{2}\right) \\&= \frac{1}{2} \frac{1}{2\pi} e^{-\frac{(u+v)^2}{8}} e^{-\frac{(u-v)^2}{8}} = \frac{1}{4\pi} e^{-\frac{u^2+2uv+v^2}{8} - \frac{u^2-2uv+v^2}{8}} \\&= \frac{1}{4\pi} e^{-\frac{u^2}{4}} e^{-\frac{v^2}{4}} = \left(\frac{1}{\sqrt{2\pi}\sqrt{2}} e^{-\frac{u^2}{4}} \right) \left(\frac{1}{\sqrt{2\pi}\sqrt{2}} e^{-\frac{v^2}{4}} \right)\end{aligned}$$

- $U \sim N(0, 2)$ and $V \sim N(0, 2)$ are also independent (it actually holds as long as same variance)

independence

- there is a much simpler, but very important, situation in which the new variables U and V are independent
- **theorem** (CB 4.3.5): let X and Y be independent random variables, then $U = g(X)$ and $V = h(Y)$ are also independent.
- **proof**: consider the continuous case and define $\Omega_u = \{x : g(x) \leq u\}$ and $\Omega_v = \{y : h(y) \leq v\}$, then

$$\begin{aligned}F_{U,V}(u, v) &= \mathbb{P}(U \leq u, V \leq v) = \mathbb{P}(X \in \Omega_u, Y \in \Omega_v) \\&= \mathbb{P}(X \in \Omega_u) \mathbb{P}(Y \in \Omega_v) \\f_{U,V}(u, v) &= \frac{\partial^2}{\partial u \partial v} F_{U,V}(u, v) = \left(\frac{d}{du} \mathbb{P}(X \in \Omega_u) \right) \left(\frac{d}{dv} \mathbb{P}(Y \in \Omega_v) \right)\end{aligned}$$

the first term is a function only of u and the second term is a function only of v ■

find a partition if necessary

- in some situations of interest the transformation is not bijective...
- find a partition A_0, A_1, \dots, A_k of $\Omega_{X,Y}$, for which the set A_0 is such that $\mathbb{P}((X, Y) \in A_0) = 0$, whereas $(U, V) = (g_1(X, Y), g_2(X, Y))$ is one-to-one from A_i to $\Omega_{U,V}$ for each $i = 1, \dots, k$
- Then...

$$f_{U,V}(u, v) = \sum_{i=1}^k f_{X,Y}(h_{1i}(u, v), h_{2i}(u, v)) |J_i|$$

just like in the univariate case.

ratio of independent normal variables

example: let $U = X/Y$ and $V = |Y|$, with $X \sim N(0, 1) \perp\!\!\!\perp Y \sim N(0, 1)$

- $\Omega_{U,V} = \{(u, v) : v > 0\}$
- $A_0 = \{(x, y) : y = 0\}$, $A_1 = \{(x, y) : y > 0\}$, $A_2 = \{(x, y) : y < 0\}$
- $h_{11}(u, v) = uv$, $h_{21}(u, v) = v \Rightarrow |J_1| = |v \cdot 1 - u \cdot 0| = |v|$
- $h_{12}(u, v) = -uv$, $h_{21}(u, v) = -v \Rightarrow |J_2| = |(-v) \cdot (-1) + u \cdot 0| = |v|$
- Using the result above,

$$\begin{aligned} f_{U,V}(u, v) &= \frac{1}{2\pi} e^{-(uv)^2/2} e^{-v^2/2} |v| + \frac{1}{2\pi} e^{-(-uv)^2/2} e^{-(-v)^2/2} |v| \\ &= (v/\pi) e^{-(1+u^2)v^2/2} \quad -\infty < u < \infty \quad 0 < v < \infty \end{aligned}$$

ratio of independent normal variables

- the distribution of the ratio of independent normals is the marginal of U :

$$\begin{aligned}f_U(u) &= \int_0^\infty (v/\pi) e^{-(u^2+1)v^2/2} dv \\&= \frac{1}{2\pi} \int_0^\infty e^{-z(1+u^2)/2} dz\end{aligned}$$

where we used $z = v^2 \Rightarrow dz = 2v dv$. By noticing that the integrand is kernel of exponential pdf with $\beta = \frac{2}{u^2+1}$, we get that

$$\int_0^\infty e^{-z(1+u^2)/2} dz = \frac{2}{1+u^2}$$

and therefore

$$f_U(u) = \frac{1}{2\pi} \frac{2}{1+u^2} = \frac{1}{\pi(1+u^2)} \quad -\infty < u < \infty$$

which is a Cauchy distribution. (intuitive, right?...)

Contents

1. Joint and marginal distributions
2. Conditional distribution and independence
3. Bivariate transformations
- 4. Hierarchical models, mixtures and a LIE**
5. Covariance and correlation
6. Multivariate distributions
7. Inequalities
8. Exercises

hierarchy

- we have so far seen probability models in which a random variable has a single distribution, possibly depending on some fixed parameters
- **however...** it is sometimes useful to think about distributions with random parameters that follow themselves some known distribution
- **advantage** the main benefit is to handle intricate structures by means of a sequence of relatively simple models in a hierarchy
- **classic example:** how many eggs will survive on average if an insect lays a large number of eggs, each surviving with probability p ?

binomial-Poisson hierarchy

- let's make some assumptions...
- large number of eggs is a random variable $N \sim \text{Poisson}(\lambda)$
- each egg's survival is independent and hence we may model their survival as Bernoulli trials $X|N \sim \text{Bin}(N, p)$

$$\begin{aligned}\mathbb{P}(X = x) &= \sum_{n=0}^{\infty} \mathbb{P}(X = x, N = n) = \sum_{n=0}^{\infty} \mathbb{P}(X = x|N = n) \mathbb{P}(N = n) \\&= \sum_{n=x}^{\infty} \binom{n}{x} p^x (1-p)^{n-x} \frac{e^{-\lambda} \lambda^n}{n!} \\&= e^{-\lambda} (\lambda p)^x \sum_{n=x}^{\infty} \frac{n!}{(n-x)! x!} (1-p)^{n-x} \frac{\lambda^{n-x}}{n!} \\&= \frac{e^{-\lambda} (\lambda p)^x}{x!} \sum_{n=x}^{\infty} \frac{[(1-p)\lambda]^{n-x}}{(n-x)!} = \frac{e^{-\lambda} (\lambda p)^x}{x!} \sum_{t=0}^{\infty} \frac{[(1-p)\lambda]^t}{t!} \\&= \frac{e^{-\lambda} (\lambda p)^x}{x!} e^{(1-p)\lambda} = \frac{(\lambda p)^x e^{-\lambda p}}{x!} \Rightarrow X \sim \text{Poisson}(\lambda p)\end{aligned}$$

law of iterated expectations (LIE)

- **theorem:** if X and Y are any two random variables, then

$$\mathbb{E}(X) = \mathbb{E}[\mathbb{E}(X|Y)]$$

as long as the expectations exist.

- **proof:** $f_{X,Y}(x,y) = f_{X|Y}(x|y)f_Y(y)$ by definition, and hence

$$\begin{aligned}\mathbb{E}(X) &= \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} x f_{X,Y}(x,y) dx dy \\ &= \int_{-\infty}^{\infty} \left[\int_{-\infty}^{\infty} x f_{X|Y}(x|y) dx \right] f_Y(y) dy \\ &= \int_{-\infty}^{\infty} \mathbb{E}(X|y) f_Y(y) dy = \mathbb{E}[\mathbb{E}(X|Y)]\end{aligned}$$



- **binomial-Poisson hierarchy:** $\mathbb{E}(X) = \mathbb{E}[\mathbb{E}(X|N)] = \mathbb{E}(Np) = \lambda p$

mixture of distributions

- **definition:** a random variable X has a mixture distribution if the distribution of X depends on a quantity that also has a distribution
- any distribution arising from a hierarchy meets this definition
- **example:** $\text{Poisson}(\lambda p)$ is a mixture distribution as it results from the combination of a binomial distribution $\text{Bin}(N, p)$ and $N \sim \text{Poisson}(\lambda)$
- **example:** there is nothing to stop the hierarchy at two layers of structure there are now a large number of mother insects from which we draw one at random
 - $X|N \sim \text{Bin}(N, p)$
 - $N|\Lambda \sim \text{Poisson}(\lambda)$
 - $\Lambda \sim \text{Exp}(\theta)$

noncentral chi-squared distribution

- apart from aiding understanding, the hierarchical structure also helps with some moment calculations
- **example:** let X have a noncentral chi-squared distribution with p degrees of freedom and noncentrality parameter λ , then

$$f_X(x|\lambda, p) = \sum_{k=0}^{\infty} \frac{x^{p/2+k-1} e^{-x^2}}{\Gamma(p/2 + k) 2^{p/2+k}} \frac{\lambda^k e^{-\lambda}}{k!}$$

it is not so messy to compute $\mathbb{E}(X)$ if one realizes that $X|K \sim \chi_{p/2+K}^2$ and $K \sim \text{Poisson}(\lambda)$

$$\mathbb{E}(X) = \mathbb{E}[\mathbb{E}(X|K)] = \mathbb{E}(p + 2K) = p + 2\lambda$$

conditional variance identity

- theorem: $\text{var}(X) = \mathbb{E}[\text{var}(X|Y)] + \text{var}[\mathbb{E}(X|Y)]$

- proof:

$$\begin{aligned}\text{var}(X) &= \mathbb{E}[X - \mathbb{E}(X)]^2 \\&= \mathbb{E}[X - \mathbb{E}(X|Y) + \mathbb{E}(X|Y) - \mathbb{E}(X)]^2 \\&= \mathbb{E}[X - \mathbb{E}(X|Y)]^2 + \mathbb{E}[\mathbb{E}(X|Y) - \mathbb{E}(X)]^2 \\&\quad + 2\mathbb{E}\{[X - \mathbb{E}(X|Y)][\mathbb{E}(X|Y) - \mathbb{E}(X)]\} \\&\stackrel{LIE}{=} \mathbb{E}(\mathbb{E}\{[X - \mathbb{E}(X|Y)]^2|Y\}) + \text{var}[\mathbb{E}(X|Y)] \\&\quad + 2\mathbb{E}(\mathbb{E}\{[X - \mathbb{E}(X|Y)][\mathbb{E}(X|Y) - \mathbb{E}(X)]|Y\}) \\&= \mathbb{E}[\text{var}(X|Y)] + \text{var}[\mathbb{E}(X|Y)] \\&\quad + 2\mathbb{E}\{[\mathbb{E}(X|Y) - \mathbb{E}(X|Y)][\mathbb{E}(X|Y) - \mathbb{E}(X)]\} \\&= \mathbb{E}[\text{var}(X|Y)] + \text{var}[\mathbb{E}(X|Y)] \quad \blacksquare\end{aligned}$$

Contents

1. Joint and marginal distributions
2. Conditional distribution and independence
3. Bivariate transformations
4. Hierarchical models, mixtures and a LIE
- 5. Covariance and correlation**
6. Multivariate distributions
7. Inequalities
8. Exercises

how to gauge the strength of a relationship?

- let X and Y measure the weight and volume of a sample of water
 - if we gauge the pair (X, Y) in several samples and plot them
 - then data points should fall on a straight line in the absence of measurement errors
- let X and Y measure the body weight and height of a person
 - if we gauge the pair (X, Y) in several samples and plot them
 - then data points should also exhibit an upward trend, though not exactly a straight line

definitions

- the covariance between X and Y is

$$\text{cov}(X, Y) = \mathbb{E}[(X - \mu_X)(Y - \mu_Y)] = \mathbb{E}(XY) - \mu_X \mu_Y,$$

with $\mu_X = \mathbb{E}(X)$ and $\mu_Y = \mathbb{E}(Y)$, whereas the correlation is

$$\text{corr}(X, Y) = \mathbb{E}\left(\frac{X - \mu_X}{\sigma_X} \frac{Y - \mu_Y}{\sigma_Y}\right) = \frac{1}{\sigma_X \sigma_Y} \text{cov}(X, Y),$$

with $\sigma_X = \sqrt{\text{var}(X)}$ and $\sigma_Y = \sqrt{\text{var}(Y)}$

- independence** (CB 4.5.5): if X and Y are independent random variables then $\text{cov}(X, Y) = \text{corr}(X, Y) = 0$

counterexamples

Independence implies $\mathbb{E}(XY) = \mathbb{E}(X)\mathbb{E}(Y)$, but not vice-versa.

1. let X be -1 or 1 with probability 0.5. Let Y be 0 if $X = -1$. If $X = 1$, Y is randomly -1 or 1 with probability 0.5. X and Y are not independent
– however...

$$\begin{aligned}\mathbb{E}(XY) &= -1 \cdot 0 \cdot \mathbb{P}(X = -1) + 1 \cdot 1 \cdot \mathbb{P}(X = 1, Y = 1) \\ &\quad + 1 \cdot -1 \cdot \mathbb{P}(X = 1, Y = -1) \\ &= 0\end{aligned}$$

and $\mathbb{E}(X) = \mathbb{E}(Y) = 0$.

2. A standard normal distribution is such that $\mathbb{E}(X) = \mathbb{E}(X^3) = 0$. Take $Y = X^2$. Then

$$\text{cov}(X, Y) = \mathbb{E}(XY) - \mathbb{E}(X) \cdot \mathbb{E}(Y) = 0$$

example of linear relationship

- example: let $X \sim U(0, 1) \perp\!\!\!\perp Z \sim U(0, 1/10)$ and $Y = X + Z$.
- then the joint pdf of (X, Y) is $f_{X,Y}(x, y) = 10$ for $0 < x < 1$ and $x < y < x + 1/10$, with

$$\begin{aligned}\mathbb{E}(Y) &= \mathbb{E}(X) + \mathbb{E}(Z) = 1/2 + 1/20 = 11/20 \\ \text{cov}(X, Y) &= \mathbb{E}(XY) - \mathbb{E}(X)\mathbb{E}(Y) = \mathbb{E}[X(X + Z)] - \mathbb{E}(X)\mathbb{E}(X + Z) \\ &= \mathbb{E}(X^2) + \mathbb{E}(XZ) - [\mathbb{E}(X)]^2 - \mathbb{E}(X)\mathbb{E}(Z) \\ &= \text{var}(X) = \frac{1}{12}(1 - 0)^2 = \frac{1}{12} \\ \text{var}(Y) &= \text{var}(X + Z) = \text{var}(X) + \text{var}(Z) = \frac{1}{12} + \frac{1}{1200} = \frac{101}{1200} \\ \text{corr}(X, Y) &= \frac{\text{cov}(X, Y)}{\sqrt{\text{var}(X)\text{var}(Y)}} = \frac{1/12}{\sqrt{1/12 \times 101/1200}} = \sqrt{\frac{100}{101}}\end{aligned}$$

example of nonlinear relationship


- example: let $X \sim U(-1, 1) \perp\!\!\!\perp Z \sim U(0, 1/10)$ and $Y = X^2 + Z$, then the joint pdf of (X, Y) is $f_{X,Y}(x, y) = 5$ for $-1 < x < 1$ and $x^2 < y < x^2 + 1/10$, with

$$\begin{aligned}\text{cov}(X, Y) &= \mathbb{E}(XY) - \mathbb{E}(X)\mathbb{E}(Y) \\ &= \mathbb{E}[X(X^2 + Z)] - \mathbb{E}(X)\mathbb{E}(X^2 + Z) \\ &= \mathbb{E}(X^3) + \mathbb{E}(XZ) - \mathbb{E}(X)\mathbb{E}(X^2) - \mathbb{E}(X)\mathbb{E}(Z) \\ &= \mathbb{E}(X^3) + \mathbb{E}(X)\mathbb{E}(Z) - \mathbb{E}(X)\mathbb{E}(X^2) - \mathbb{E}(X)\mathbb{E}(Z) \\ &= 0\end{aligned}$$

given that $\mathbb{E}(X) = \mathbb{E}(X^3) = 0$ due to the symmetric nature of X

- there is a strong dependence between X and Y , but it is not linear...

how does it look like?



linear dependence

- **theorem** (CB 4.5.7): For any random variables X and Y ,
 - (i) $|\text{corr}(X, Y)| \leq 1$
 - (ii) $|\text{corr}(X, Y)| = 1$ if and only if there exist numbers $a \neq 0$ and b such that $\mathbb{P}(Y = aX + b) = 1$, with $a > 0$ if $\text{corr}(X, Y) > 0$ and $a < 0$ if $\text{corr}(X, Y) < 0$
- **proof of (i)**: define $h(t) = \mathbb{E}[(X - \mu_X)t + (Y - \mu_Y)]^2$, so $h(t) \geq 0, \forall t$

$$\begin{aligned}h(t) &= t^2 \mathbb{E}(X - \mu_X)^2 + 2t \mathbb{E}(X - \mu_X)(Y - \mu_Y) + \mathbb{E}(Y - \mu_Y)^2 \\&= t^2 \sigma_X^2 + \sigma_Y^2 + 2t \text{cov}(X - \mu_X, Y - \mu_Y)\end{aligned}$$

and hence it can have at most one real root, implying a nonpositive discriminant,

$$\begin{aligned}[2 \text{cov}(X, Y)]^2 - 4\sigma_X^2 \sigma_Y^2 &\leq 0 \quad \Rightarrow \quad -\sigma_X \sigma_Y \leq \text{cov}(X, Y) \leq \sigma_X \sigma_Y \\&\Rightarrow \quad |\text{corr}(X, Y)| \leq 1\end{aligned}$$

linear dependence

- **proof of (ii):** now, $\text{corr}(X, Y) = 1 \Leftrightarrow [2t \text{cov}(X, Y)]^2 - 4t^2 \sigma_X^2 \sigma_Y^2 = 0$, i.e., $h(t)$ has a single root. Given that $[(X - \mu_X)t + (Y - \mu_Y)]^2 \geq 0$ for all t , $h(t) = 0$ if and only if

$$\mathbb{P}\left([(X - \mu_X)t + (Y - \mu_Y)]^2 = 0\right) = 1$$

\Downarrow

$$\mathbb{P}((X - \mu_X)t + (Y - \mu_Y) = 0) = 1$$

which is equivalent to $\mathbb{P}(Y = aX + b) = 1$ with $a = -t = \frac{\text{cov}(X, Y)}{\sigma_X^2}$ and $b = \mu_X t + \mu_Y$ ■

- we will see that the **Cauchy-Schwartz inequality** considerably shortens the proof above

variance decomposition

- **theorem** (CB 4.5.6): if X and Y are any two random variables, and a and b are any two constants, then

$$\text{var}(aX + bY) = a^2 \text{var}(X) + b^2 \text{var}(Y) + 2ab \text{cov}(X, Y)$$

- **proof:** it follows from $\mathbb{E}(aX + bY) = a\mu_X + b\mu_Y$ that

$$\begin{aligned} \text{var}(aX + bY) &= \mathbb{E}[(aX + bY) - (a\mu_X + b\mu_Y)]^2 \\ &= \mathbb{E}[a(X - \mu_X) + b(Y - \mu_Y)]^2 \\ &= \mathbb{E}[a^2(X - \mu_X)^2 + b^2(Y - \mu_Y)^2 + 2a(X - \mu_X)b(Y - \mu_Y)] \\ &= a^2[\mathbb{E}(X - \mu_X)^2] + b^2\mathbb{E}[(Y - \mu_Y)^2] \\ &\quad + 2ab\mathbb{E}[(X - \mu_X)(Y - \mu_Y)] \\ &= a^2 \text{var}(X) + b^2 \text{var}(Y) + 2ab \text{cov}(X, Y) \quad \blacksquare \end{aligned}$$

- the variation in $X + Y$ is inferior to the sum of the variations in X and Y if $\text{cov}(X, Y) < 0$ because large values of X are more likely to occur with small values of Y

bivariate normal

- **definition:** the **bivariate normal distribution** with parameters $\mu_X, \mu_Y, \sigma_X^2 > 0, \sigma_Y^2 > 0$ and $|\rho| < 1$

$$f_{X,Y}(x,y) = \frac{1}{2\pi\sigma_X\sigma_Y\sqrt{\rho}} \exp \left\{ -\frac{1}{2(1-\rho^2)} \left[\left(\frac{x-\mu_X}{\sigma_X} \right)^2 + \left(\frac{y-\mu_Y}{\sigma_Y} \right)^2 - 2\rho \frac{x-\mu_X}{\sigma_X} \frac{y-\mu_Y}{\sigma_Y} \right] \right\}$$

for $-\infty < x < \infty$ and $-\infty < y < \infty$.

- the following properties hold (**proofs left as exercise**):
 - $\text{corr}(X, Y) = \rho$
 - $X \sim N(\mu_X, \sigma_X^2)$ and $Y \sim N(\mu_Y, \sigma_Y^2)$
 - $X|Y \sim N\left(\mu_X + \rho \frac{\sigma_X}{\sigma_Y} (Y - \mu_Y), \sigma_X^2(1 - \rho^2)\right)$

Contents

1. Joint and marginal distributions
2. Conditional distribution and independence
3. Bivariate transformations
4. Hierarchical models, mixtures and a LIE
5. Covariance and correlation
- 6. Multivariate distributions**
7. Inequalities
8. Exercises

joint, marginal and conditional probabilities

- **discrete:** the joint pmf of $\mathbf{X} = (X_1, \dots, X_n) \subset \mathbb{R}^n$ is a function $f_{\mathbf{X}}(\mathbf{x})$ such that

$$\mathbb{P}(\mathbf{X} \in A) = \sum_{\mathbf{x} \in A} f_{\mathbf{X}}(\mathbf{x})$$

for any $A \subset \mathbb{R}^n$

- **continuous:** the joint pdf of $\mathbf{X} = (X_1, \dots, X_n) \subset \mathbb{R}^n$ is a function $f_{\mathbf{X}}(\mathbf{x})$ such that

$$\mathbb{P}(\mathbf{X} \in A) = \int \cdots \int_A f(x_1, \dots, x_n) dx_1 \cdots dx_n$$

for any $A \subset \mathbb{R}^n$

- **expectation:**

$$\mathbb{E}[g(\mathbf{x})] = \begin{cases} \int_{-\infty}^{\infty} \cdots \int_{-\infty}^{\infty} g(\mathbf{x}) f_{\mathbf{X}}(\mathbf{x}) d\mathbf{x} & \text{if continuous} \\ \sum_{\mathbf{x} \in \mathbb{R}^n} g(\mathbf{x}) f_{\mathbf{X}}(\mathbf{x}) & \text{if discrete} \end{cases}$$

joint, marginal and conditional probabilities

- **marginals** with respect to a subset of the variables can be obtained integrating with respect to the other variables

$$f(x_1, \dots, x_k) = \int_{-\infty}^{\infty} \cdots \int_{-\infty}^{\infty} f(x_1, \dots, x_n) dx_{k+1} \cdots dx_n$$

- similarly, the **conditional pdf** is

$$f(x_{k+1}, \dots, x_n | x_1, \dots, x_k) = \frac{f(x_1, \dots, x_n)}{f(x_1, \dots, x_k)}$$

example

- example: let

$$f(x_1, x_2, x_3, x_4) = \begin{cases} \frac{3}{4}(x_1^2 + x_2^2 + x_3^2 + x_4^2) & 0 < x_i < 1, i = 1, 2, 3, 4 \\ 0 & \text{o.w.} \end{cases}$$

- verify that:

- (i) $\int_0^1 \int_0^1 \int_0^1 \int_0^1 f(x_1, x_2, x_3, x_4) dx_1 dx_2 dx_3 dx_4 = 1$
- (ii) $\mathbb{P}(X_1 < \frac{1}{2}, X_2 < \frac{3}{4}, X_4 > \frac{1}{2}) = \frac{3}{256}$
- (iii) $f(x_1, x_2) = \frac{3}{4}(x_1^2 + x_2^2) + \frac{1}{2}$
- (iv) $\mathbb{E}X_1 X_2 = \frac{5}{16}$
- (v) $f(x_3, x_4 | x_1, x_2) = \frac{x_1^2 + x_2^2 + x_3^2 + x_4^2}{x_1^2 + x_2^2 + \frac{2}{3}}$

multinomial distribution

- Bernoulli trials now have n distinct outcomes, with probabilities p_1, \dots, p_n , common across trials. X_i represents the number of times that the i th outcome happened among m trials.
- **example:** toss a six-sided dice and let Z be the outcome. The dice is unbalanced and $\mathbb{P}(Z = z) = \frac{z}{21}$. Consider now tossing the dice ten times, and X_i counts the number of times i came up. Then $X = (X_1, X_2, \dots, X_6)$ has a multinomial distribution with $m = 10$ trials, $n = 6$ possible outcomes, and

$$\begin{aligned} f(0, 0, 1, 2, 3, 4) &= \frac{10!}{0!0!1!2!3!4!} \left(\frac{1}{21}\right)^0 \left(\frac{2}{21}\right)^0 \left(\frac{3}{21}\right)^1 \left(\frac{4}{21}\right)^2 \left(\frac{5}{21}\right)^3 \left(\frac{6}{21}\right)^4 \\ &= 0.0059 \end{aligned}$$

multinomial distribution

- **definition:** let n and m denote positive integers, then the discrete random vector $\mathbf{X} = (X_1, \dots, X_n)$ has a multinomial distribution with m trials and cell probabilities $0 \leq p_1, \dots, p_n \leq 1$ such that $\sum_{i=1}^n p_i = 1$ if the joint pmf of \mathbf{X} is given by

$$f_{\mathbf{X}}(x_1, \dots, x_n) = \frac{m!}{x_1! \cdots x_n!} p_1^{x_1} \cdots p_n^{x_n} = m! \prod_{i=1}^n \frac{p_i^{x_i}}{x_i!}$$

for $\mathbf{x} = (x_1, \dots, x_n)$ such that each integer $x_i \geq 0$ and $\sum_{i=1}^n x_i = m$

marginal and conditional pmfs of a multinomial

if the discrete random vector $\mathbf{X} = (X_1, \dots, X_n)$ is multinomial with m trials and cell probabilities $0 \leq p_1, \dots, p_n \leq 1$, (you may try to show these properties)

- the marginal of X_i is binomial $\text{Bin}(m, p_i)$
- the conditional distribution of $(X_1, \dots, X_{i-1}, X_{i+1}, \dots, X_n)$ given $X_i = x_i$ is multinomial with $m - x_i$ trials and cell probabilities $p_j / (1 - p_i)$ for $1 \leq j \neq i \leq n$
- there is some negative correlation given that $\sum_{i=1}^n X_i = m$ $\text{corr}(X_i, X_j) = -mp_i p_j$ for $1 \leq i \neq j \leq n$

independence

- **definition:** let $\mathbf{X}_1, \dots, \mathbf{X}_n$ denote random vectors with joint pdf/pmf $f_{\mathbf{X}}(\mathbf{x}_1, \dots, \mathbf{x}_n)$ and marginal pdf/pmf $f_{\mathbf{X}_i}(\mathbf{x}_i)$, then they are mutually independent random vectors if, for every $(\mathbf{x}_1, \dots, \mathbf{x}_n)$,

$$f_{\mathbf{X}}(\mathbf{x}_1, \dots, \mathbf{x}_n) = f_{\mathbf{X}_1}(\mathbf{x}_1) \cdots f_{\mathbf{X}_n}(\mathbf{x}_n) = \prod_{i=1}^n f_{\mathbf{X}_i}(\mathbf{x}_i)$$

- we now need to generalize the results we had for independent bivariate distributions

independence

if X_1, \dots, X_n are independent,

(1) let g_1, \dots, g_n be real-valued functions such that $g_i(x_i)$ is a function only of x_i .

$$\mathbb{E}[g_1(X_1) \cdots g_n(X_n)] = \prod_{i=1}^n \mathbb{E}[g_i(X_i)]$$

(2) let $M_{X_1}(t), \dots, M_{X_N}(t)$ be the mgfs of X_1, \dots, X_N and $Z = \sum_{i=1}^n X_i$. Then

$$M_Z(t) = \prod_{i=1}^n M_{X_i}(t)$$

(3) let $a_1, \dots, a_n, b_1, \dots, b_n$ be fixed constants and $Z = \sum_{i=1}^n a_i X_i + b_i$. Then

$$M_Z(t) = \left(e^{t \sum b_i} \right) \prod_{i=1}^n M_{X_i}(a_i t)$$

independence

if X_1, \dots, X_n are independent,

(4) X_1, \dots, X_n are independent if, and only if, there exists functions $g_i(x_i)$ such that

$$f(x_1, \dots, x_n) = \prod_{i=1}^n g_i(x_i)$$

(5) $U_1 = g_1(X_1), \dots, U_n = g_n(X_n)$ are also mutually independent

independence and normality

- **example** (CB 3.6.10): $X_i \sim N(\mu_i, \sigma_i^2)$, mutually independent. Let $a_1, \dots, a_n, b_1, \dots, b_n$ be fixed constants. Then

$$Z = \sum_{i=1}^n (a_i X_i + b_i) \sim N\left(\sum_{i=1}^n (a_i \mu_i + b_i), \sum_{i=1}^n a_i^2 \sigma_i^2\right)$$

- **proof**: the mgf of a normal random variable is $M(t) = e^{\mu t + \sigma^2 t^2 / 2}$. Then

$$\begin{aligned} M_Z(t) &= \left(e^{t \sum b_i}\right) \prod_{i=1}^n e^{\mu_i a_i t + \sigma_i^2 a_i^2 t^2 / 2} \\ &= e^{t \sum (a_i \mu_i + b_i) + (\sum a_i^2 \sigma_i^2) t^2 / 2} \end{aligned}$$

which is the mgf of a $N\left(\sum_{i=1}^n (a_i \mu_i + b_i), \sum_{i=1}^n a_i^2 \sigma_i^2\right)$. ■

multivariate normal

- the pdf of multivariate normal distributions is

$$f_X(x) = \frac{1}{(2\pi)^{n/2}} |\Sigma|^{-\frac{1}{2}} e^{-\frac{1}{2}(x-\mu)'\Sigma^{-1}(x-\mu)}$$

for n -dimensional X . Denote $X \sim N(\mu, \Sigma)$.

- lemma:** let $Z \sim N(0, I_n)$ and $X = \mu + \Sigma^{1/2}Z$. Then $X \sim N(\mu, \Sigma)$.
- proof:** the distribution of Z is

$$f_Z(z) = \frac{1}{(2\pi)^{n/2}} e^{-\frac{1}{2}z'z}$$

and the transformation $x = \mu + \Sigma^{1/2}z$ has Jacobian $|\Sigma|^{-1/2}$.

- lemma:** if $Y = AX + b$, then $Y \sim N(A\mu + b, A\Sigma A')$.
- proof:** follows from previous slide.

multivariate normal

- take a partition $X = [X_1', X_2']'$, with $X \sim N(\mu, \Sigma)$ and let

$$\mu = \begin{bmatrix} \mu_1 \\ \mu_2 \end{bmatrix} \text{ and } \Sigma = \begin{bmatrix} \Sigma_{11} & \Sigma_{12} \\ \Sigma_{21} & \Sigma_{22} \end{bmatrix}$$

- **theorem:** X_1 and X_2 are independent if and only if $\Sigma_{12} = 0$.
- **proof (\Rightarrow):** this is immediate (independent random variables imply zero correlation)
- **proof (\Leftarrow):** let $\Sigma_{12} = 0$ and write

$$\Sigma = \begin{pmatrix} \Sigma_{11} & 0 \\ 0 & \Sigma_{22} \end{pmatrix}$$

then

$$\begin{aligned} f_X(x) &= \frac{1}{(2\pi)^{n/2}} |\Sigma|^{-\frac{1}{2}} \exp \left\{ -\frac{1}{2} (x - \mu)' \Sigma^{-1} (x - \mu) \right\} \\ &= \frac{1}{(2\pi)^{n_1/2}} |\Sigma_{11}|^{-\frac{1}{2}} \exp \left\{ -\frac{1}{2} (x_1 - \mu_1)' \Sigma_{11}^{-1} (x_1 - \mu_1) \right\} \\ &\quad \times \frac{1}{(2\pi)^{n_2/2}} |\Sigma_{22}|^{-\frac{1}{2}} \exp \left\{ -\frac{1}{2} (x_2 - \mu_2)' \Sigma_{22}^{-1} (x_2 - \mu_2) \right\} \\ &= f_{X_1}(x_1) \cdot f_{X_2}(x_2) \end{aligned}$$

multivariate normal

- **theorem:** the conditional distribution of $X_1|X_2$ is $N(\mu_{1.2}, \Sigma_{11.2})$ with

$$\begin{aligned}\mu_{1.2} &= \mu_1 + \Sigma_{12}\Sigma_{22}^{-1}(X_2 - \mu_2) \\ \Sigma_{11.2} &= \Sigma_{11} + \Sigma_{12}\Sigma_{22}^{-1}\Sigma_{21}\end{aligned}$$

- **proof:** consider a random vector given by

$$\begin{bmatrix} X_1 - \Sigma_{12}\Sigma_{22}^{-1}X_2 \\ X_2 \end{bmatrix} = \begin{bmatrix} I & -\Sigma_{12}\Sigma_{22}^{-1} \\ 0 & I \end{bmatrix} \begin{bmatrix} X_1 \\ X_2 \end{bmatrix}$$

which is a linear transformation of a normal random vector X . The two subvectors $X_1 - \Sigma_{12}\Sigma_{22}^{-1}X_2$ and X_2 are uncorrelated,

$$\begin{aligned}\text{Var} \begin{bmatrix} X_1 - \Sigma_{12}\Sigma_{22}^{-1}X_2 \\ X_2 \end{bmatrix} &= \begin{bmatrix} I & -\Sigma_{12}\Sigma_{22}^{-1} \\ 0 & I \end{bmatrix} \begin{bmatrix} \Sigma_{11} & \Sigma_{12} \\ \Sigma_{21} & \Sigma_{22} \end{bmatrix} \begin{bmatrix} I & 0 \\ -\Sigma_{12}'\Sigma_{22}^{-1} & I \end{bmatrix} \\ &= \begin{bmatrix} \Sigma_{11} - \Sigma_{12}\Sigma_{22}^{-1}\Sigma_{21} & 0 \\ \Sigma_{21} & \Sigma_{22} \end{bmatrix} \begin{bmatrix} I & 0 \\ -\Sigma_{12}'\Sigma_{22}^{-1} & I \end{bmatrix} \\ &= \begin{bmatrix} \Sigma_{11} - \Sigma_{12}\Sigma_{22}^{-1}\Sigma_{21} & 0 \\ 0 & \Sigma_{22} \end{bmatrix}\end{aligned}$$

therefore independent.

multivariate normal

- proof (cont'd): write

$$X_1 = \Sigma_{12}\Sigma_{22}^{-1}X_2 + (X_1 - \Sigma_{12}\Sigma_{22}^{-1}X_2)$$

where the term in brackets is independent of X_2 , so its conditional distribution given X_2 is consequently the same as its unconditional distribution, which is normal with mean $\mu_1 - \Sigma_{12}\Sigma_{22}^{-1}\mu_2$ and variance $\Sigma_{11} - \Sigma_{12}\Sigma_{22}^{-1}\Sigma_{21}$.

- then

$$\begin{aligned} E(X_1|X_2) &= E(\Sigma_{12}\Sigma_{22}^{-1}X_2|X_2) + E(X_1 - \Sigma_{12}\Sigma_{22}^{-1}X_2) \\ &= \mu_1 + \Sigma_{12}\Sigma_{22}^{-1}(X_2 - \mu_2) \end{aligned}$$

$$\begin{aligned} \text{Var}(X_1|X_2) &= \text{Var}(\Sigma_{12}\Sigma_{22}^{-1}X_2|X_2) + \text{Var}(X_1 - \Sigma_{12}\Sigma_{22}^{-1}X_2) \\ &= \Sigma_{11} - \Sigma_{12}\Sigma_{22}^{-1}\Sigma_{21} \end{aligned}$$



transformations of random vectors

- denote $\mathbf{U} = (U_1, \dots, U_n)$, with $U_i = g_i(X_1, \dots, X_n)$ for $i = 1, \dots, n$.
- let the support set be $\Omega_X = \{x : f_X(x) > 0\}$
- find partitions $A_0, A_1, A_2, \dots, A_k$ such that $\mathbb{P}(X \in A_0) = 0$ and g is a bijective transformation within each A_j , $j > 0$
- we then have inverse transformations $x_1 = h_{1j}(u_1, \dots, u_n)$, \dots , $x_n = h_{nj}(u_1, \dots, u_n)$ for each $j > 0$
- the Jacobian term is given by

$$J_j = \begin{vmatrix} \frac{\partial x_1}{\partial u_1} & \frac{\partial x_1}{\partial u_2} & \dots & \frac{\partial x_1}{\partial u_n} \\ \frac{\partial x_2}{\partial u_1} & \frac{\partial x_2}{\partial u_2} & \dots & \frac{\partial x_2}{\partial u_n} \\ \vdots & \ddots & \dots & \vdots \\ \frac{\partial x_n}{\partial u_1} & \frac{\partial x_n}{\partial u_2} & \dots & \frac{\partial x_n}{\partial u_n} \end{vmatrix} = \begin{vmatrix} \frac{\partial h_{1j}(\mathbf{u})}{\partial u_1} & \frac{\partial h_{1j}(\mathbf{u})}{\partial u_2} & \dots & \frac{\partial h_{1j}(\mathbf{u})}{\partial u_n} \\ \frac{\partial h_{2j}(\mathbf{u})}{\partial u_1} & \frac{\partial h_{2j}(\mathbf{u})}{\partial u_2} & \dots & \frac{\partial h_{2j}(\mathbf{u})}{\partial u_n} \\ \vdots & \ddots & \dots & \vdots \\ \frac{\partial h_{nj}(\mathbf{u})}{\partial u_1} & \frac{\partial h_{nj}(\mathbf{u})}{\partial u_2} & \dots & \frac{\partial h_{nj}(\mathbf{u})}{\partial u_n} \end{vmatrix}$$

with $x_i = h_{ij}(\mathbf{u})$ for any $x_i \in A_j$ with $i = 1, \dots, n$ and $j = 1, \dots, k$

transformations of random vectors

- then...

$$f_U(u_1, \dots, u_n) = \sum_{j=1}^k f_X(h_{1j}(u_1, \dots, u_n), \dots, h_{nj}(u_1, \dots, u_n)) |J_j|,$$

- example: joint pdf $f_X(x_1, x_2, x_3, x_4) = 24e^{-x_1-x_2-x_3-x_4}$ with $0 < x_1 < x_2 < x_3 < x_4 < \infty$ and $U_1 = X_1$, $U_2 = X_2 - X_1$, $U_3 = X_3 - X_2$ and $U_4 = X_4 - X_3$
 - $X_1 = U_1$, $X_2 = U_1 + U_2$, $X_3 = U_1 + U_2 + U_3$, $X_4 = U_1 + U_2 + U_3 + U_4$
 - Jacobian

$$J = \begin{vmatrix} 1 & 0 & 0 & 0 \\ 1 & 1 & 0 & 0 \\ 1 & 1 & 1 & 0 \\ 1 & 1 & 1 & 1 \end{vmatrix} = 1$$

- so $f_U(u_1, \dots, u_4) = 24e^{-4u_1-3u_2-2u_3-u_4}$

Contents

1. Joint and marginal distributions
2. Conditional distribution and independence
3. Bivariate transformations
4. Hierarchical models, mixtures and a LIE
5. Covariance and correlation
6. Multivariate distributions
- 7. Inequalities**
8. Exercises

a lemma

- **lemma:** let $a, b > 0$ and $p, q > 1$ such that $\frac{1}{p} + \frac{1}{q} = 1$, then $\frac{1}{p}a^p + \frac{1}{q}b^q \geq ab$ with equality if and only if $a^p = b^q$.
- **sketch of proof:** fix b and minimize

$$g(a) = \frac{1}{p}a^p + \frac{1}{q}b^q - ab$$

with respect to a . We get

$$\frac{dg(a)}{da} = 0 \Rightarrow a^{p-1} - b = 0 \Rightarrow b = a^{p-1}$$

The second derivative $\frac{d^2g(a)}{da^2} = (p-1)a^{p-2} > 0$, indeed a minimum. The value at the minimum is

$$\frac{1}{p}a^p + \frac{1}{q}a^{q(p-1)} - a^p = \frac{1}{p}a^p + \frac{1}{q}a^p - a^p = 0$$

since $\frac{1}{p} + \frac{1}{q} = 1 \Rightarrow q + p = pq \Rightarrow q(p-1) = p$. Equality holds if $b = a^{p-1} \Rightarrow a^p = b^q$. ■

Hölder's inequality

- **theorem:** let X and Y denote any two random variables and let p and q satisfy $\frac{1}{p} + \frac{1}{q} = 1$, then

$$|\mathbb{E}(XY)| \leq \mathbb{E}|XY| \leq (\mathbb{E}|X|^p)^{1/p} (\mathbb{E}|Y|^q)^{1/q}$$

- **proof:** the first inequality follows from the fact that

$$-|XY| \leq XY \leq |XY| \Rightarrow -\mathbb{E}|XY| \leq \mathbb{E}(XY) \leq \mathbb{E}|XY|.$$

to prove the second inequality, choose

$$a = \frac{|X|}{(\mathbb{E}|X|^p)^{1/p}} \quad \text{and} \quad b = \frac{|Y|}{(\mathbb{E}|Y|^q)^{1/q}}$$

which, using the lemma, implies

$$\begin{aligned} \frac{1}{p} \frac{|X|^p}{(\mathbb{E}|X|^p)} + \frac{1}{q} \frac{|X|^q}{(\mathbb{E}|X|^q)} &\geq \frac{|X|}{(\mathbb{E}|X|^p)^{1/p}} \frac{|Y|}{(\mathbb{E}|Y|^q)^{1/q}} \\ &= \frac{|XY|}{(\mathbb{E}|X|^p)^{1/p} (\mathbb{E}|Y|^q)^{1/q}} \end{aligned}$$

Hölder's inequality

- proof (cont'd): taking expectations on both sides,

$$\underbrace{\frac{1}{p} \frac{\mathbb{E}|X|^p}{(\mathbb{E}|X|^p)} + \frac{1}{q} \frac{\mathbb{E}|X|^q}{(\mathbb{E}|X|^q)}}_{=\frac{1}{p}+\frac{1}{q}=1} \geq \frac{\mathbb{E}|XY|}{(\mathbb{E}|X|^p)^{1/p}(\mathbb{E}|Y|^q)^{1/q}}$$
$$\Downarrow$$
$$\mathbb{E}|XY| \leq (\mathbb{E}|X|^p)^{1/p}(\mathbb{E}|Y|^q)^{1/q}$$

which completes the proof. ■

Hölder: $|\mathbb{E}(XY)| \leq \mathbb{E}|XY| \leq (\mathbb{E}|X|^p)^{1/p}(\mathbb{E}|Y|^q)^{1/q}$

- selecting $p = q = 2$, we obtain the **Cauchy-Schwarz inequality**: for any random variables X and Y ,

$$|\mathbb{E}(XY)| \leq \mathbb{E}|XY| \leq \sqrt{\mathbb{E}(X^2)}\sqrt{\mathbb{E}(Y^2)}$$

- **covariance inequality**: applying the Cauchy-Schwartz inequality to $X - \mu_X$ and $Y - \mu_Y$ yields

$$|\text{cov}(X, Y)| \leq \sigma_X \sigma_Y$$

or, equivalently, that $|\text{corr}(X, Y)| \leq 1$.

- **Lyapunov's inequality**: set $Y = 1$, replace $|X|$ by $|X|^r$ for $1 < r < p$ and define $s = pr$ to obtain

$$(\mathbb{E}|X|^r)^{1/r} \leq (\mathbb{E}|X|^s)^{1/s}$$

for $1 < r < s < \infty$

Minkowski's inequality

- **theorem:** let X and Y denote any two random variables, then

$$(\mathbb{E}|X + Y|^p)^{1/p} \leq (\mathbb{E}|X|^p)^{1/p} + (\mathbb{E}|Y|^p)^{1/p} \quad 0 \leq p < \infty$$

- **proof:** triangular inequality $|X + Y| \leq |X| + |Y|$ ensures that

$$\begin{aligned} \mathbb{E}|X + Y|^p &= \mathbb{E}(|X + Y||X + Y|^{p-1}) \\ &\leq \mathbb{E}(|X||X + Y|^{p-1}) + \mathbb{E}(|Y||X + Y|^{p-1}) \\ &\leq (\mathbb{E}|X|^p)^{1/p} \left(\mathbb{E}|X + Y|^{q(p-1)} \right)^{1/q} \\ &\quad + (\mathbb{E}|Y|^p)^{1/p} \left(\mathbb{E}|X + Y|^{q(p-1)} \right)^{1/q} \end{aligned}$$

for $1/p + 1/q = 1$ where Hölder's inequality was applied twice.

Minkowski's inequality

- proof (cont'd): dividing by $\left(\mathbb{E}|X+Y|^{q(p-1)}\right)^{1/q}$,

$$\frac{\mathbb{E}|X+Y|^p}{\left(\mathbb{E}|X+Y|^{q(p-1)}\right)^{1/q}} \leq (\mathbb{E}|X|^p)^{1/p} + (\mathbb{E}|Y|^p)^{1/p}$$

and since $\frac{1}{p} + \frac{1}{q} = 1 \Rightarrow p + q = pq \Rightarrow qp - q = p$,

$$\begin{aligned} \frac{\mathbb{E}|X+Y|^p}{\left(\mathbb{E}|X+Y|^{q(p-1)}\right)^{1/q}} &= \frac{\mathbb{E}|X+Y|^p}{(\mathbb{E}|X+Y|^p)^{1/q}} \\ &= (\mathbb{E}|X+Y|^p)^{1-\frac{1}{q}} \\ &= (\mathbb{E}|X+Y|^p)^{\frac{1}{p}} \end{aligned}$$

which completes the proof. ■

Contents

1. Joint and marginal distributions
2. Conditional distribution and independence
3. Bivariate transformations
4. Hierarchical models, mixtures and a LIE
5. Covariance and correlation
6. Multivariate distributions
7. Inequalities
8. Exercises

Reference:

- Casella and Berger, Ch. 4

Exercises:

- 4.1, 4.4–4.7, 4.9, 4.10, 4.13, 4.15, 4.22, 4.24, 4.26, 4.30, 4.32, 4.37, 4.38, 4.41–4.43, 4.47, 4.58, 4.59.