

Lecture #18: Power Series

Date: Mon. 12/3/18

Power Series

Recall the basics of a geometric series

$$\sum_{k=0}^{\infty} ar^k = \frac{a}{1-r} \quad \text{for } |r| < 1$$

This is a power series if we replace r with x . i.e.

$$\sum_{n=0}^{\infty} ax^n = \frac{a}{1-x}, \quad \text{for } |x| < 1$$

In this case we say the power series

$$\sum_{n=0}^{\infty} ax^n$$

Converges for x in the interval $(-1, 1)$

This is a power series centered @ 0.

A general power series is centered @ a by replacing x with $x-a$

Def A series of the form

$$\sum_{n=0}^{\infty} c_n(x-a)^n = c_0 + c_1(x-a) + c_2(x-a)^2 + \dots$$

is a power series centered @ $x=a$

Lecture # 18: Power Series


Date: Mon. 12/3/18

Not every power series is geometric, but there will be x values for which the series will converge.

Convergence of a Power Series

Def a power series $\sum_{n=0}^{\infty} C_n(x-a)^n$ has coeff.s C_n & center a . The set of values of x for which the series converges is the interval of convergence.

The radius of convergence written R , is the distance from the center to the edge of the interval of convergence



Lecture # 18: Power Series

Date: Mon. 12/3/18

The points on the edges of the interval of convergence may or may not be in the interval of convergence. This must be checked in each case.

How do we find the int. of convergence?

Generally, we need the ratio or root test most of the time i.e. testing for absolute convergence.

Thm a power series $\sum C_k(x-a)^k$ can converge in one of 3 ways:

(i) The series converges for all x
The int. of conv. is $(-\infty, \infty)$; $R = \infty$

(ii) The series converges on a finite interval $(a-R, a+R)$ for $0 < R < \infty$, where R is the radius of convergence. The pts $a-R$ & $a+R$ need to be checked individually.

(iii) The series only converges @ the center $x=a$. The radius of conv. is $R=0$.

Lecture # 18: Power Series

Date: Mon. 12/3/18

Ex. 1) $\sum_{k=0}^{\infty} \frac{x^k}{k!}$

Ratio Test:

$$a_k = \frac{x^k}{k!}$$

$$a_{k+1} = \frac{x^{k+1}}{(k+1)!}$$

$$\frac{a_{k+1}}{a_k} = \frac{x^{k+1}}{(k+1)!} \cdot \frac{k!}{x^k} = \frac{x \cdot \cancel{x^k} \cdot \cancel{k!}}{(k+1) \cdot \cancel{k!} \cdot \cancel{x^k}} = \frac{x}{k+1}$$

$$\Rightarrow \lim_{k \rightarrow \infty} \left| \frac{a_{k+1}}{a_k} \right| = \lim_{k \rightarrow \infty} \left| \frac{x}{k+1} \right| = |x| \lim_{k \rightarrow \infty} \frac{1}{k+1} = 0$$

By the Ratio Test, since $\lim_{k \rightarrow \infty} \left| \frac{a_{k+1}}{a_k} \right| < 1$
the series Converges.

So $\sum \left| \frac{x^k}{k!} \right|$ converges $\Rightarrow \sum \frac{x^k}{k!}$ converges for all x

Radius of Convergence : $R = \infty$

Interval of Convergence : $(-\infty, \infty)$

Lecture # 18: Power Series

Date: Mon. 12/3/18

$$\text{Ex. 2)} \sum_{k=0}^{\infty} \frac{(-1)^k (x-2)^k}{4^k}$$

Root Test:

$$a_k = \frac{(-1)^k (x-2)^k}{4^k} = \left(-\frac{(x-2)}{4} \right)^k$$

$$\sqrt[k]{|a_k|} = \left[\left| -\frac{(x-2)}{4} \right|^k \right]^{1/k} = \left| -\frac{(x-2)}{4} \right| = \frac{|x-2|}{4}$$

$$\Rightarrow \rho = \lim_{k \rightarrow \infty} \sqrt[k]{|a_k|} = \lim_{k \rightarrow \infty} \frac{|x-2|}{4} = \frac{|x-2|}{4}$$

For series to converge need $\rho < 1$

$$\Rightarrow \frac{|x-2|}{4} < 1 \Rightarrow |x-2| < 4$$

$$\text{Note: } |x-2| = \begin{cases} x-2, & x \geq 2 \\ -(x-2), & x < 2 \end{cases}$$

$$\Rightarrow x-2 < 4 \quad \& \quad -(x-2) < 4$$

$$x < 6$$

$$x-2 > -4$$

$$x > -2$$

So series converges for $-2 < x < 6$

Lecture #18: Power Series

Date: Mon. 12/3/18

Ex. 2 (cont'd)Need to check $x = -2$ & $x = 6$ For $x = -2$:

$$\sum_{k=0}^{\infty} \frac{(-1)^k (-2-2)^k}{4^k} = \sum_{k=0}^{\infty} (-1)^k \left(\frac{-4}{4}\right)^k = \sum_{k=0}^{\infty} (-1)^k (-1)^k$$

$$= \sum_{k=0}^{\infty} 1^k$$

By Divergence test since


$$\lim_{k \rightarrow \infty} a_k = \lim_{k \rightarrow \infty} 1 = 1 \neq 0 \quad \text{this series diverges}$$

i.e. $x = -2$ is not included in interval of convergenceFor $x = 6$:

$$\sum (-1)^k \frac{(6-2)^k}{4^k} = \sum (-1)^k \frac{4^k}{4^k} = \sum (-1)^k$$

Again, diverges by divergence test

$$\lim_{k \rightarrow \infty} a_k = \lim_{k \rightarrow \infty} (-1)^k = \text{DNE}$$

So interval of convergence is $(-2, 6)$ 

Lecture #18: Power Series

Date: Mon. 12/3/18

Ex. 3) $\sum_{k=1}^{\infty} k! x^k$

Ratio Test:

$$a_k = k! x^k$$

$$a_{k+1} = (k+1)! x^{k+1}$$

$$\frac{a_{k+1}}{a_k} = \frac{(k+1)! x^{k+1}}{k! x^k} = \frac{(k+1) \cancel{k!} x \cdot \cancel{x^k}}{\cancel{k!} \cancel{x^k}} = x(k+1)$$

So

$$\lim_{k \rightarrow \infty} \left| \frac{a_{k+1}}{a_k} \right| = \lim_{k \rightarrow \infty} |x(k+1)| = \infty$$

By ratio test since $\lim_{k \rightarrow \infty} \left| \frac{a_{k+1}}{a_k} \right| > 1$

Series diverges

Since center @ $x=0$ & no other value of x will make $\lim_{k \rightarrow \infty} \left| \frac{a_{k+1}}{a_k} \right|$ finite

\Rightarrow interval of convergence: $x=0$

Radius of convergence: $R=0$

Lecture # 18: Power Series

Date: Mon. 12/3/18

Ex. 4 Find the interval & radius of
Conv. for

$$\sum_{k=1}^{\infty} \frac{(x-2)^k}{\sqrt{k}}$$

Ratio Test:

$$a_k = \frac{(x-2)^k}{\sqrt{k}}$$

$$a_{k+1} = \frac{(x-2)^{k+1}}{\sqrt{k+1}}$$

$$\begin{aligned} \frac{a_{k+1}}{a_k} &= \frac{(x-2)^{k+1}}{\sqrt{k+1}} \cdot \frac{\sqrt{k}}{(x-2)^k} = \frac{(x-2)\sqrt{k}}{\sqrt{k+1}} = (x-2) \frac{\sqrt{k}}{\sqrt{k+1}} \\ &= (x-2) \left(\frac{k}{k+1} \right)^{1/2} \end{aligned}$$

$$\begin{aligned} \lim_{k \rightarrow \infty} \left| \frac{a_{k+1}}{a_k} \right| &= \lim_{k \rightarrow \infty} \left| (x-2) \left(\frac{k}{k+1} \right)^{1/2} \right| = \left| (x-2) \left(\lim_{k \rightarrow \infty} \frac{k}{k+1} \right)^{1/2} \right| \\ &\stackrel{H}{=} \left| (x-2) \left(\lim_{k \rightarrow \infty} \frac{1}{1} \right)^{1/2} \right| \\ &= |x-2| \end{aligned}$$

So series converges if $|x-2| < 1 \Rightarrow 1 < x < 3$

For $x=1$

$$\sum_{k=1}^{\infty} \frac{(1-2)^k}{\sqrt{k}} = \sum_{k=1}^{\infty} \frac{(-1)^k}{\sqrt{k}} = \sum_{k=1}^{\infty} (-1)^k \left(\frac{1}{\sqrt{k}} \right)$$

By Alt. series test, Since

$$\lim_{k \rightarrow \infty} a_k = \lim_{k \rightarrow \infty} \frac{1}{\sqrt{k}} = 0 \quad \text{so series converges}$$

Lecture # 18: Power Series

Date: Mon. 12/3/18

Ex. 4 (cont'd)For $x = 3$

$$\sum_{k=1}^{\infty} \frac{(3-2)^k}{1^k} = \sum_{k=1}^{\infty} \frac{1^k}{1^k} = \sum_{k=1}^{\infty} \frac{1}{k^{1/2}}$$

By p-series test since $p < 1$
Series Diverges.

\Rightarrow interval of convergence: $[1, 3)$

Radius of convergence: $R = 1$

Representing Functions as Power Series

We can create new power series thru mult. & composition. This is best illustrated w/ examples.

Ex. 5 $f(x) = \frac{x^5}{1-x}$ Write this as a power series

$$f(x) = \frac{x^5}{1-x} = x^5 \underbrace{\left(\frac{1}{1-x} \right)}_{\text{Has form } \frac{1}{1-r}}$$

$$= x^5 \sum_{n=0}^{\infty} x^n = \sum_{n=0}^{\infty} x^5 x^n$$

$$= \sum_{n=0}^{\infty} x^{n+5}, \quad |x| < 1$$

Lecture # 18: Power Series

Date: Mon. 12/3/18

Ex. 6 | $g(x) = \frac{1}{1-2x}$

$$g(x) = \frac{1}{1-2x} = \sum_{k=0}^{\infty} (2x)^k \quad \text{for } |2x| < 1$$

$$\Rightarrow |x| < \frac{1}{2}$$

Ex. 7 | $h(x) = \frac{1}{1+x^2}$

$$h(x) = \frac{1}{1+x^2} = \frac{1}{1-(-x^2)} = \sum_{n=0}^{\infty} (-x^2)^n, \quad \begin{array}{l} | -x^2 | < 1 \\ | x^2 | < 1 \end{array}$$

$$|x^2| = \begin{cases} -x^2, & x < 0 \\ x^2, & x \geq 0 \end{cases}$$

$$-x^2 < 1 \quad x^2 > 1$$

$$x^2 > -1$$

$$x > \sqrt{-1}$$

$$\frac{x > \pm 1}{x > -1 \text{ \& } x < 1}$$

$$|x| < 1$$

Lecture # 18: Power Series

Date: Mon. 12/3/18

Operations on Power Series

One of the key reasons why power series are so useful is that if

$$f(x) = \sum c_k (x-a)^k, \quad |x-a| < R$$

i.e. we have a power series rep. for a f , then we get term-by-term

Differentiation:

$$f'(x) = \sum k c_k (x-a)^{k-1}, \quad |x-a| < R$$

Integration:

$$\int f(x) dx = \underset{\substack{\uparrow \\ \text{constant of integration}}}{C} + \sum_{n=0}^{\infty} \frac{c_n}{n+1} (x-a)^{n+1}$$

i.e. we treat power series like a polyn.

Lecture #18: Power Series

Date: Mon. 12/3/18

Ex. 8 | $f(x) = \frac{1}{1-x} \Rightarrow f(x) = \sum_{k=0}^{\infty} x^k, |x| < 1$

Find $f'(x)$.

$$f'(x) = \sum_{k=0}^{\infty} k x^{k-1}, |x| < 1$$

Ex. 9 | Integrate $g(x) = \frac{1}{1+x}$ using a power series

Rep. as power series:

$$g(x) = \frac{1}{1+x} = \frac{1}{1-(-x)} = \sum_{k=0}^{\infty} (-x)^k, |x| < 1$$

Integrate:

$$\int \frac{1}{1+x} dx = C + \sum_{k=0}^{\infty} \frac{(-x)^{k+1}}{k+1}, |x| < 1$$