

CLINICAL STUDY REPORT - SUMMARY OF RESULTS

Study CB207-C301 - Calibra Biologics, Inc.

Version 1.0 18 July 2025

Prepared by Halcyon Clinical Research Organization

DERIVED DEMONSTRATION CORPUS
CORPUS DE DEMONSTRATION DERIVE

Synthetic corpus - generated by cross-doc-qc - not a regulatory document

APPROVAL

Sponsor Head of Clinical Development
PPD

DERIVED DEMONSTRATION CORPUS
CORPUS DE DEMONSTRATION DERIVE

Synthetic corpus - generated by cross-doc-qc - not a regulatory document

1 SYNOPSIS

Study CB207-C301 was a randomised, double-blind, parallel-group, multicentre study comparing CB-207 with the reference product in the neoadjuvant treatment of HER2-positive early breast cancer. The study was conducted at 148 sites in 21 countries between February 2022 and March 2025.

DERIVED DEMONSTRATION CORPUS
CORPUS DE DEMONSTRATION DERIVE

Synthetic corpus - generated by cross-doc-qc - not a regulatory document

2 SUBJECT DISPOSITION

A total of 811 subjects were randomised, of whom 406 were assigned to CB-207 and 405 to the reference product. Enrolment completed in November 2023.

Of the randomised subjects, 803 subjects received at least one dose of study treatment and constituted the Safety Set. The Per-Protocol Set comprised 744 subjects, 374 in the CB-207 group and 370 in the reference product group.

Nine subjects enrolled at two sites in Ukraine were excluded from the Per-Protocol Set in accordance with the analysis plan.

DERIVED DEMONSTRATION CORPUS
CORPUS DE DEMONSTRATION DERIVE

3 DEMOGRAPHICS AND BASELINE CHARACTERISTICS

Demographic and baseline characteristics were balanced between treatment groups. Median age was 51 years and hormone receptor positive disease was documented in 58% of randomised subjects.

DERIVED DEMONSTRATION CORPUS
CORPUS DE DEMONSTRATION DERIVE

Synthetic corpus - generated by cross-doc-qc - not a regulatory document

4 PRIMARY EFFICACY RESULT

In the Per-Protocol Set, the breast pathological complete response rate was 49.8% in the CB-207 group and 41.3% in the reference product group.

The observed ratio of breast pathological complete response rates was 1.206 with a two-sided 90% confidence interval of 1.068 to 1.362.

The observed difference in breast pathological complete response rates was 8.50% with a two-sided 95% confidence interval of 1.93 to 15.07.

DERIVED DEMONSTRATION CORPUS
CORPUS DE DEMONSTRATION DERIVE

5 EQUIVALENCE ASSESSMENT

The equivalence assessment was performed against the two acceptance criteria pre-specified in the statistical analysis plan. The outcome of each assessment is reported in Section 4 and discussed in Section 9.

DERIVED DEMONSTRATION CORPUS
CORPUS DE DEMONSTRATION DERIVE

6 SECONDARY EFFICACY RESULTS

The total pathological complete response rate was 43.6% in the CB-207 group and 36.8% in the reference product group. Overall clinical response rates were comparable between groups.

DERIVED DEMONSTRATION CORPUS
CORPUS DE DEMONSTRATION DERIVE

7 SAFETY RESULTS

Adverse events were graded using the Common Terminology Criteria for Adverse Events version 4.0. Treatment-emergent adverse events of any grade were reported in 94% of subjects in each treatment group.

Asymptomatic decreases in left ventricular ejection fraction were reported in 4.1% of subjects in the CB-207 group and 3.8% in the reference product group.

All subjects who received study treatment were scheduled for a safety follow-up assessment 30 days after the last dose of study treatment.

DERIVED DEMONSTRATION CORPUS
CORPUS DE DEMONSTRATION DERIVE

8 IMMUNOGENICITY RESULTS

Treatment-emergent anti-drug antibodies were detected in 1.2% of subjects in the CB-207 group and 1.5% in the reference product group. No neutralising antibody was associated with loss of response.

DERIVED DEMONSTRATION CORPUS
CORPUS DE DEMONSTRATION DERIVE

9 DISCUSSION AND CONCLUSIONS

The study enrolled more subjects than the number planned in the analysis plan. The primary analysis was performed as pre-specified and the observed treatment effect was consistent across the pre-specified subgroups.

The safety and immunogenicity profiles of CB-207 and the reference product were comparable. The sponsor considers the totality of evidence to support biosimilarity.

DERIVED DEMONSTRATION CORPUS
CORPUS DE DEMONSTRATION DERIVE