

INVESTIGATOR'S BROCHURE

CB-207 (proposed biosimilar HER2-directed monoclonal antibody)

Edition 4 27 February 2024

Sponsor: Calibra Biologics, Inc.

DERIVED DEMONSTRATION CORPUS
CORPUS DE DEMONSTRATION DERIVE

Synthetic corpus - generated by cross-doc-qc - not a regulatory document

1 SUMMARY

CB-207 is a proposed biosimilar to the reference HER2-directed monoclonal antibody. This edition supersedes Edition 3 and incorporates clinical data available through the data cut-off of 31 December 2023.

CB-207 is not approved for marketing in any jurisdiction and is to be administered only within the context of an approved clinical protocol.

DERIVED DEMONSTRATION CORPUS
CORPUS DE DEMONSTRATION DERIVE

2 PHYSICAL, CHEMICAL AND PHARMACEUTICAL PROPERTIES

CB-207 drug product is supplied as a lyophilised powder for reconstitution in single-use vials containing 150 mg and 420 mg of active substance. Reconstituted solution is stable for up to 24 hours at 2 to 8 degrees Celsius.

DERIVED DEMONSTRATION CORPUS
CORPUS DE DEMONSTRATION DERIVE

3 NON-CLINICAL PHARMACOLOGY

CB-207 demonstrated binding affinity, antibody-dependent cellular cytotoxicity, and inhibition of proliferation equivalent to the reference product across the analytical comparability programme.

DERIVED DEMONSTRATION CORPUS
CORPUS DE DEMONSTRATION DERIVE

4 NON-CLINICAL TOXICOLOGY

Repeat-dose toxicology in cynomolgus monkeys identified no target organ toxicity distinguishable from that of the reference product. Cardiac findings were absent at all dose levels evaluated.

DERIVED DEMONSTRATION CORPUS
CORPUS DE DEMONSTRATION DERIVE

5 HUMAN PHARMACOKINETICS

CB-207 exhibits target-mediated disposition with a terminal elimination half-life supporting administration every 21 days. Steady state is reached after approximately four cycles.

DERIVED DEMONSTRATION CORPUS
CORPUS DE DEMONSTRATION DERIVE

6 CLINICAL EXPERIENCE - DEVELOPMENT PROGRAMME

The clinical development programme comprises a single-dose comparative pharmacokinetic study in healthy volunteers, a first-in-patient dose-finding study, and the ongoing confirmatory study CB207-C301.

Across the programme, eligible subjects were adults with an Eastern Cooperative Oncology Group (ECOG) performance status of 0 to 2 at study entry and adequate organ function. Requirements applicable to an individual study are specified in the protocol for that study.

DERIVED DEMONSTRATION CORPUS
CORPUS DE DEMONSTRATION DERIVE

7 CLINICAL EXPERIENCE - STUDY CB207-C101 (DOSE FINDING)

Study CB207-C101 was an open-label Phase 1 dose-escalation study in subjects with HER2-positive advanced solid tumours. On the basis of the observed tolerability and exposure data from this Phase 1 dose-escalation study, the dose selected for further evaluation was 4 mg/kg administered once weekly.

Adverse events in study CB207-C101 were graded using the Common Terminology Criteria for Adverse Events version 4.03, which was the version in force at the time that study was conducted.

DERIVED DEMONSTRATION CORPUS
CORPUS DE DEMONSTRATION DERIVE

8 CLINICAL EXPERIENCE - ONGOING CONFIRMATORY STUDY

Study CB207-C301 is an ongoing randomised, double-blind confirmatory study in the neoadjuvant treatment of HER2-positive early breast cancer. Subjects receive a loading dose of 8 mg/kg followed by a maintenance dose of 6 mg/kg every 21 days.

DERIVED DEMONSTRATION CORPUS
CORPUS DE DEMONSTRATION DERIVE

9 SUMMARY OF SAFETY

The most frequently reported adverse reactions are infusion-related reactions, neutropenia, fatigue, and asymptomatic decrease in left ventricular ejection fraction. The safety profile observed to date is consistent with that of the reference product.

DERIVED DEMONSTRATION CORPUS
CORPUS DE DEMONSTRATION DERIVE

10 IDENTIFIED AND POTENTIAL RISKS

Cardiac dysfunction is an identified risk of the pharmacological class. Left ventricular ejection fraction should be determined before initiation of treatment and monitored during treatment in accordance with the applicable protocol.

Infusion-related reactions and pulmonary toxicity are identified risks. Immunogenicity has been observed at a low incidence and has not been associated with loss of efficacy.

DERIVED DEMONSTRATION CORPUS
CORPUS DE DEMONSTRATION DERIVE

11 GUIDANCE FOR THE INVESTIGATOR

The dose, schedule, permitted modifications, and monitoring requirements applicable to an individual subject are those specified in the protocol under which that subject is enrolled. Where this brochure and the applicable protocol differ, the protocol governs the conduct of the study.

DERIVED DEMONSTRATION CORPUS
CORPUS DE DEMONSTRATION DERIVE

12 DOCUMENT CONTROL

This edition was approved by the sponsor Safety Management Team on 27 February 2024 and is distributed to all participating investigators and ethics committees. The next periodic review is due within twelve months.

DERIVED DEMONSTRATION CORPUS
CORPUS DE DEMONSTRATION DERIVE

Synthetic corpus - generated by cross-doc-qc - not a regulatory document