

1 Exercises for Gravimetry

1.1 The shell theorem

1.2 Gravitational potential inside and outside a sphere with constant density

1.3 Detection of a spherical object in the sub-surface

(a) In the lecture we have discussed the general shape of a gravity anomaly for a sphere with radius R located at depth z below the surface. Formulate the problem in a relative sense using density contrasts ($\Delta\rho$). Derive an analytical expression for the vertical anomaly as a function of the lateral distance x and depth z . Calculate the maximum anomalies that you would expect for some realistic settings (e.g., a limestone cave.)

Solution: Because we have a spherical object, there is not much difference to the point mass scenario discussed in class. The gravitational force exhibited by the sphere is:

$$\vec{g} = G \frac{M}{r^2} \hat{r}$$

The mass anomaly M (formulated in a relative sense) of the sphere is given by:

$$M = \frac{4}{3}\pi R^3 \Delta\rho$$

At distance x the distance to the sphere is $r = \sqrt{x^2 + z^2}$. The gravimeter only measures the vertical component of \vec{g} :

$$g_z = |\vec{g}| \sin \alpha = |\vec{g}| \frac{z}{r}$$

which brings us to:

$$\begin{aligned} g_z &= \frac{4}{3}\pi R^3 \Delta\rho \frac{G}{r^2} \frac{z}{r} \\ &= \frac{4}{3}\pi R^3 \Delta\rho \frac{G}{r^3} z \\ &= \frac{4}{3}\pi R^3 \Delta\rho G \frac{z}{(x^2 + z^2)^{\frac{3}{2}}} \end{aligned}$$

We will encounter the maximum value of the anomaly at $x = 0$ (directly above the target):

$$g_{z,max} = \frac{4}{3}\pi R^3 \Delta\rho G \frac{1}{z^2}$$

which for the limestone cave example is approximately -0.35 mGal (see below).

(b) Use a piece of software of your choice (e.g., Excel, SciDAVis, Matlab, Python) and plot the expected vertical gravity anomaly for a specific setting as a function of lateral distance x . Visualize how this profile changes as you vary, e.g., the depth of the object. Label your axis and post a picture in the forum alongside with a comment which software you used.

```

1 clear all
2 close all
3
4 %This is vector x for the lateral distance
5 x = -150:0.01:150;
6 %Depth of the sphere
7 z = 50;
8 %Radius of the sphere
9 R = 25;
10 %Constants for the limestone cave scenario
11 %(SI stands for System International units)
12 DensityLimestone_InGramsPerCentimeterCubed = 2.0;
13 DensityLimestone_SI = DensityLimestone_InGramsPerCentimeterCubed *1/1000 * 100^3;
14 DenistyAir_SI = 1.2;
15 %Density Contrast in kg/m^3
16 DeltaRho_SI = DenistyAir_SI-DensityLimestone_SI;
17 %Gravitational constants
18 G_SI = 6.674e-11;
19
20
21 %This is the formula we want to plot
22 gz_SI = 4/3*pi*R^3*DeltaRho_SI*G_SI*z./((z.^2+x.^2).^(3/2));
23 gz_SI_max = 4/3*pi*R^3*DeltaRho_SI*G_SI*1/(z^2);
24 gz_mGal = gz_SI*100*1000;
25
26 fig=figure(1)
27 plot(x,gz_mGal-gz_mGal(1),'r-','LineWidth',3);hold on
28
29 %And for a different depths
30 z=75;
31 gz_SI = 4/3*pi*R^3*DeltaRho_SI*G_SI*z./((z.^2+x.^2).^(3/2));
32 gz_mGal = gz_SI*100*1000;
33 plot(x,gz_mGal-gz_mGal(1),'b-','LineWidth',3);hold on
34
35 z=100;
36 gz_SI = 4/3*pi*R^3*DeltaRho_SI*G_SI*z./((z.^2+x.^2).^(3/2));
37 gz_mGal = gz_SI*100*1000;
38 plot(x,gz_mGal-gz_mGal(1),'g-','LineWidth',3);hold on
39
40 legend('Cave at 50 m', 'Cave at 75 m','Cave at 100 m','Location','SouthEast')
41 xlabel('Distance (m)')
42 ylabel('g_z anomaly (mGal)')
43 %Export to a png. (This can be done much better.)
44 set(findall(fig,'-property','FontSize'),'FontSize',15,'fontWeight','bold')
45 print('-dpng','-r300','GravityAnomaly.png')
46
47
48 -2*G_SI*6.0e24/((6370*1000)^3)*100*1000

```

../Src/Exercises/Gravimetry/Gravimetry01.m