

Introduction to Geophysics
R. Drews


Gravimetry

Learning goals today:

- ▶ The gravitational force, its potential field, and how to measure it.
- ▶ The reference gravitational field of the Earth.


Example: Global variability

Introduction to Geophysics



Example: Global variability

Your mass is constant but your weight is not.



What is a force?

Introduction to Geophysics



[Newton (1642-1726) / G. Johnson.]

$$\vec{F} = m\vec{g}$$

\vec{F} : Force (N; kg m s^{-2})

\vec{g} : Acceleration (m s^{-2})

m : Mass (kg)

The gravitational force

Introduction to Geophysics




[Newton (1642-1726) / G. Johnson.]

$$\vec{F} = G \frac{mM}{r^2} \hat{r}$$

$$G = 6.674 \cdot 10^{-11} \text{ (m}^3\text{kg}^{-1}\text{s}^{-2}\text{)}$$


\hat{r} : unit vector

r : distance between point masses



Example: The gravitational constant


Introduction to Geophysics



Cavendish, PNAS, 1798

Example: The gravitational constant

Introduction to Geophysics



©nature

Westphal et al., Nature, 2021

G is the worst known constant in physics.
Why?

Example: Measuring acceleration

Introduction to Geophysics



$$\vec{F} = m\vec{g}$$

$$\vec{F} = G \frac{mM}{r^2} \hat{r}$$

$$\rightarrow \vec{g} = G \frac{M}{r^2} \hat{r}$$

$$\rightarrow \frac{d^2\vec{x}}{dt^2} = G \frac{M}{r^2} \hat{r}$$

This is a differential equation.


Example: Measuring acceleration



$$\frac{d^2 \vec{x}}{dt^2} = G \frac{M}{R_E^2} \approx \text{const.}$$

At the Earth's surface (R_E) g is close to constant and only vertical. (Later we will see that none of this is not quite true).


Example: Measuring acceleration




Example: Measuring acceleration

$$v = \int g dt = \frac{d}{dt} x(t) = \frac{GM}{R_e^2} t + c_1$$

$v(m s^{-1})$



Example: Measuring acceleration



Example: Measuring acceleration


Introduction to Geophysics

$$x(t) = \frac{GM}{2R_e^2}t^2 + c_1t + c_2$$

- ▶ Setting, e.g., $c_1 = 0$ (initial velocity) and $c_2 = 0$ (initial position) is quite convenient.
- ▶ This is the principle of a free-fall gravimeter.

- ▶ Thanks to the Greeks we know the radius R_E for the Earth. However, its mass was unknown for a while.
- ▶ Go ahead and determine the mass of the Earth M with your Smartphone!
- ▶ There is an important first-order finding in Earth Sciences that you can (re-) discover. Which one?


Beyond point masses



$$\vec{F} = G \frac{dM}{r^2} \hat{r}$$

For a small mass dM the point mass approximation holds.

Beyond point masses




$$\vec{F} = G \frac{dM}{r^2} \hat{r}$$

Profiling across a sub-surface target results in a gravity anomaly (\rightarrow Exercises).

Beyond point masses

Introduction to Geophysics




$$\vec{F}(\vec{r}) = \sum_i G \frac{dM_i}{r_i^2} \hat{r}_i$$

For i point masses the effect adds up.

Beyond point masses

Introduction to Geophysics


Surface



$$\vec{F}(\vec{r}) = \sum G \frac{dM_i}{r_i^2} \hat{r}_i$$

Beyond point masses


Surface



$$\vec{F}(\vec{r}) = \sum G \frac{dM_i}{r_i^2} \hat{r}_i$$

Beyond point masses


Introduction to Geophysics



$$\vec{F}(\vec{r}) = \sum G \frac{dM_i}{r_i^2} \hat{r}_i$$

Surface


$$\vec{F}(\vec{r}) = G \int \rho \frac{1}{r^2} \hat{r} dV$$



The summation can be replaced by an integration over a volume enclosing a continuous density.


Surface

$$\vec{F}(\vec{r}) = G \int \rho \frac{1}{r^2} \hat{r} dV$$



The integration is a triple integral. Integration limits and coordinates depend on the viewpoint. Example is a Bouger plate, in general not easy to solve (\rightarrow Exercises).

Example: Shell




[Xaononl CC BY-SA 4.0]

Newton's shell theorem solves the volume integral inside and outside spherical objects (\rightarrow Ex.-Discussion)


- ▶ The field outside a shell is the same as the one from an equivalent point mass
- ▶ The field inside a shell is zero. Everywhere.

- ▶ There are analytical solutions for other shapes (e.g., Nagy 1966 for Prism).




Numerical forward modelling (\rightarrow Ex)

Introduction to Geophysics




Vector fields

$$\vec{g} = G \frac{M}{r^2} \hat{r}$$




Potential Field

$$\vec{g} = G \frac{M}{r^2} \hat{r}$$



What is the amount of work required?



$$\begin{aligned} U(r) &= - \int_{\infty}^r \vec{g} d\vec{r} \\ &= - \int_{\infty}^r g dr \\ &= -GM \int_{\infty}^r \frac{1}{r^2} dr \\ &= -GM \left[-\frac{1}{r} \right]_{\infty}^r \\ &= GM \frac{1}{r} \end{aligned}$$

Potential for a point mass.


$$\vec{g}(r) = -\nabla U(r)$$

- ▶ It is sometimes easier to calculate the potential of an anomaly and to infer the acceleration via the gradient.
- ▶ Equipotential lines are perpendicular to the field direction.
- ▶ Equipotential lines are in general NOT lines of equal field strength (cf. with down-hill slope force in landscape)

Gravitational field of a spherical Earth

Introduction to Geophysics

The Earth's rotation minimizes gravitational acceleration at the equator. At the poles it does nothing.



Gravitational field of a spherical Earth

Introduction to Geophysics

Centripetal acceleration at P perpendicular to rotation axis parallel to O'-P:

$$g_r = \omega^2 R \cos(\theta)$$

Centripetal acceleration at P perpendicular to rotation axis parallel to O'-P:


$$g_{r,proj.} = \omega^2 R \cos^2(\theta)$$

Angular Frequency: ω

Angular Velocity: $\vec{v}_r = \vec{\omega} \times \vec{R} \cos(\theta)$

Angular Acceleration: $\vec{g}_r = \dot{\vec{v}}_r = \vec{\omega} \times \vec{\omega} \times \vec{R} \cos(\theta)$


An ellipsoidal Earth



- ▶ Rotation induces ellipsoidal shape approximate with reference ellipsoid.
- ▶ Latitudinal correction of gravitation is adjusted accordingly.


An ellipsoidal Earth

Introduction to Geophysics



- ▶ Geoid is a real-world equipotential line approximating sea level.
- ▶ It is referenced to the geometric ellipsoid.


An ellipsoidal Earth



- ▶ 2 Geoid is a real-world equipotential line approximating sea level.
- ▶ 2 It is referenced to the geometric ellipsoid.

An ellipsoidal Earth

Introduction to Geophysics



- ▶ Upwarping of geoid indicates mass excess.
- ▶ Downwarping of geoid indicates mass deficit.


Learning goals today:

- ▶ Principles of gravimeters.
- ▶ Gravity survey types (absolute & relative) and general considerations of survey layouts and structure types.
- ▶ Reduction of gravity data.

Spring-based gravimeters

Introduction to Geophysics


[cc Reyko, CC-BY-SA3.0]



Springconstant & Extension.

Pendulum-based gravimeters

Introduction to Geophysics




Eigenfrequency & Length.

$$\omega = \sqrt{\frac{g}{l}}$$

Free-fall gravimeters (\rightarrow Ex.)

Introduction to Geophysics

[FG Gravimeter from MicroGLaCoste]




Traveltimes & Distance.

- ▶ Unit used is *Gal* ($=0.01 \text{ ms}^{-2}$)
- ▶ Top *absolute* gravimeters $\sim 1\mu\text{Gal}$ ($10^{-9}g$)
- ▶ Top *relative* gravimeters $\sim 10\mu\text{Gal}$ ($10^{-9}g$)
- ▶ Typically only g_z is measured.


Instrument drift / temporal variability

Introduction to Geophysics



Instrument drift / temporal variability

Introduction to Geophysics



Absolute gravimeters are needed

- ▶ if loop closure is impossible (e.g. intercontinental surveys),
- ▶ for long-term changes such as isostatic uplift,
- ▶ as basestations for relative surveys.

Relative surveys are always easier to conduct and loop closure can cancel many error sources (e.g., instrument drift).


Every gravity survey measures:

- ▶ latitudinal variability,
- ▶ dependency on elevation,
- ▶ the surrounding terrain,
- ▶ excess mass above anomaly,
- ▶ earth & ocean tides,
- ▶ (instr. drift, motion compons.).
- ▶ density variability in the subsurface.

Latitudinal variability (\rightarrow Ex.)

Introduction to Geophysics

Extension to ellipsoid contains the same physics.




Max 5 Gal (this is large!).


Every gravity survey measures:

- ▶ latitudinal variability,
- ▶ dependency on elevation,
- ▶ the surrounding terrain,
- ▶ excess mass above anomaly,
- ▶ earth & ocean tides,
- ▶ (instr. drift, motion compons.).
- ▶ density variability in the subsurface.

Elevation correction (i.e. free-air)



Elevation correction (i.e. free-air)



The elevation correction references the gravity anomaly to the same datum (e.g., the geoid).

How does the gravitational acceleration change with elevation near the Earth's surface?

Taylor expansion near $r = R_E$:

$$g(r) = g(R_E) + \frac{dg}{dr}\Big|_{R_E}$$

Taylor expansion near $r = R_E$:

$$g(r) \approx G \frac{M}{R_E^2} - 2G \frac{M}{R_E^3} (r - R_E) + \dots$$

Taylor expansion near $r = R_E$:

$$g(r) \approx \underbrace{G \frac{M}{R_E^2}}_{\text{g at Earth's surface}} - 2G \frac{M}{R_E^3} (r - R_E) + \dots$$

Taylor expansion near $r = R_E$:

$$g(r) \approx G \frac{M}{R_E^2} - \underbrace{2G \frac{M}{R_E^3}(r - R_E)}_{\text{change with elevation}} + \dots$$

Taylor expansion near $r = R_E$:

$$g(r) \approx G \frac{M}{R_E^2} - \underbrace{2G \frac{M}{R_E^3}(r - R_E)}_{\text{change with elevation}} + \dots$$

Evaluation at let's say $r = R_E + 1$ (m)
returns a change of $\delta g(r) \approx -0.3$ mGal
per m.

Taylor expansion near $r = R_E$:

$$g(r) \approx G \frac{M}{R_E^2} - \underbrace{2G \frac{M}{R_E^3}(r - R_E)}_{\text{change with elevation}} + \dots$$


$\delta g(r) \approx -0.3$ mGal per m is large compared to the sensitivity of gravimeters, therefore the gravimeter elevation needs to be determined within centimeters using GNSS.

Every gravity survey measures:

- ▶ latitudinal variability,
- ▶ dependency on elevation,
- ▶ the surrounding terrain,
- ▶ excess mass above anomaly,
- ▶ earth & ocean tides,
- ▶ (instr. drift, motion compons.).
- ▶ density variability in the subsurface.


Reduction of gravity data: Terrain correction

Introduction to Geophysics



Reduction of gravity data: Terrain correction


Introduction to Geophysics



A neighboring mountain will reduce the measured g_z independent of target properties.

Reduction of gravity data: Terrain correction

Introduction to Geophysics



A neighboring valley will reduce the measured g_z independent of target properties.


- ▶ The terrain correction requires an elevation model and assumptions about the broad-scale sub-surface density.
- ▶ The terrain correction is positive both for surrounding valleys and mountains.

Every gravity survey measures:

- ▶ latitudinal variability,
- ▶ dependency on elevation,
- ▶ the surrounding terrain,
- ▶ excess mass above anomaly,
- ▶ earth & ocean tides,
- ▶ (instr. drift, motion compons.).
- ▶ density variability in the subsurface.

Reduction of gravity data: Terrain correction


Introduction to Geophysics



Elevation and terrain correction to not account for the mass between the measurement surface and the reference surface.

Reduction of gravity data: Terrain correction


Introduction to Geophysics



What is the effect δg_z of a horizontal plate with constant density?

Reduction of gravity data: Terrain correction


Introduction to Geophysics



$$g_z = G \rho \iiint \frac{1}{r^2} \cos(\phi) dV = ?$$

Reduction of gravity data: Terrain correction

Introduction to Geophysics



$$g_z = G\rho \iiint \frac{1}{r^2} \cos(\phi) dV = 2\pi G\rho h$$

Every gravity survey measures:


- ▶ latitudinal variability,
- ▶ dependency on elevation,
- ▶ the surrounding terrain,
- ▶ excess mass above anomaly,
- ▶ earth & ocean tides,
- ▶ (instr. drift, motion compensation),
- ▶ density variability in the subsurface.

- ▶ Tides are caused by gravity celestial bodies (i.e. Sun & Moon).
- ▶ Tidal forces vary across a spatially extended body.
- ▶ Tidal forces are balanced by centrifugal forces of two (three) body rotations.

The origin of tides




The origin of tides




Gravitational attraction is stronger on the nearside than the farside.

The origin of tides



The origin of tides




Moon

Earth

Centrifugal force can be projected into radial (i.e. parallel to Earth's gravitation) and parallel component. This leads to the force balance.

The origin of tides



Moon

The lunar gravity differential field is responsible for two tidal bulges (i.e. tides twice a day).

- ▶ Tidal forces vary across a spatially extended body.
- ▶ Tidal forces are balanced by centrifugal forces of two (three) body rotations.
- ▶ There is lots of confusion regarding the origin of tides (cf. Matsuda et al. 2015 → Ilias).
- ▶ Tide models can be used for correction

Every gravity survey measures:

- ▶ latitudinal variability,
- ▶ dependency on elevation,
- ▶ the surrounding terrain,
- ▶ excess mass above anomaly,
- ▶ earth & ocean tides,
- ▶ (instr. drift, motion compensation),
- ▶ density variability in the subsurface.

Every gravity survey measures:

- ▶ latitudinal variability,
- ▶ dependency on elevation,
- ▶ the surrounding terrain,
- ▶ excess mass above anomaly,
- ▶ earth & ocean tides,
- ▶ (instr. drift, motion compensation),
- ▶ density variability in the subsurface.