Participant Flow — BRFSS 2011–2015

BRFSS respondents assessed

Annual surveys, 2011–2015

N = 2,380,047

Excluded: all six outcomes missing

n = 535

Final analytic cohort

≥ 1 outcome non-missing

N = 2,379,512

Six disease-specific models, each fitted on respondents with non-missing outcome data for that condition

Myocardial Infarction

Included (outcome non-missing):	**2,368,389**
Excluded (missing / refused / DK):	**11,123**
Positive cases (MI reported):	**141,437**
Negative cases:	**2,226,952**
Prevalence:	**5.97%**

Coronary Heart Disease

Included (outcome non-missing):	**2,359,299**
Excluded (missing / refused / DK):	**20,213**
Positive cases (CHD reported):	**142,475**
Negative cases:	**2,216,824**
Prevalence:	**6.04%**

Stroke

Included (outcome non-missing):	**2,373,243**
Excluded (missing / refused / DK):	**6,269**
Positive cases (stroke reported):	**97,621**
Negative cases:	**2,275,622**
Prevalence:	**4.11%**

Chronic Kidney Disease

Included (outcome non-missing):	**2,370,766**
Excluded (missing / refused / DK):	**8,746**
Positive cases (CKD reported):	**79,285**
Negative cases:	**2,291,481**
Prevalence:	**3.34%**

Depression

Included (outcome non-missing):	**2,367,914**
Excluded (missing / refused / DK):	**11,598**
Positive cases (depression reported):	**446,544**
Negative cases:	**1,921,370**
Prevalence:	**18.86%**

Diabetes

Included (outcome non-missing):	**2,374,814**
Excluded (missing / refused / DK):	**4,698**
Positive cases (diabetes reported):	**302,943**
Negative cases:	**2,071,871**
Prevalence:	**12.76%**