

Modular Arithmetic

- Addition: O(n)
- Multiplication: O(n²) (*naive*)
- Multiplication: O(nlogn) (*FFT*)
- Euclid’s Rule: gcd(x, y) = gcd(x mod y, y)
- # of bits in x^y = ylog₂x ≤ n·2ⁿ
- $\frac{n}{2} \leq n! \leq n^n$
- f: S → T is 1-to-1 (injective) & onto (surjective) ⇒ | S |=| T |
- f: S → T is 1-to-1 (injective) ⇒ | T | ≥| S |
- $\sum_{i=0}^{\infty} r^i = \frac{1}{1-r}$, if r < 1
- $\sum_{i=0}^n i^2 = \frac{n(n+1)(2n+1)}{6}$
- $\sum_{i=0}^n \frac{1}{i} = O(\log_2 n)$

Extended Euclid’s GCD(x,y)

O(n³); gcd(x,y) = d = xi + yb; x ≥ y; # mod x
ext-gcd(x, y) :
if y == 0: return (x, 1, 0)
else:

#		X	Y	X/Y	X%Y		#	d	a	b	
1.		26	15	1	11		6.	1	1	0	
2.		15	11	1	4		5.	1	0	1-(3*0)	
3.		11	4	2	3		4.	1	1	0-(1*1)	
4.		4	3	1	1		3.	1	-1	1-(2*-1)	
5.		3	1	3	0		2.	1	3	-1-(1*3)	
6.		1	0				1.	1	-4	3-(1*-4)	

Fermat’s Little Theorem

if p is prime, then $\forall 1 \leq a < p$
 $a^{p-1} = 1 \bmod p$
Proof: Start by listing first p-1 positive multiples of a:
 $S = \{a, 2a, 3a, \dots (p-1)a\}$
Suppose that ra and sa are the same mod p, $\Rightarrow r = s \bmod p$
 \therefore set S of p-1 multiples of a are distinct and nonzero, that is, they must be congruent to 1, 2, 3, ... p-1 after being sorted. Multiply all congruences together and we find
 $a \cdot 2a \cdot 3a \cdot \dots (p-1) \cdot a = 1 \cdot 2 \cdot 3 \cdot \dots (p-1) \pmod p$ or better,
 $a^{(p-1)} (p-1)! = (p-1)! \bmod p$. Divide both side by (p-1)! ■

Primality Testing

any $a \rightarrow a^{N-1} = 1 \bmod N$? $\begin{cases} \text{yes} \Rightarrow \text{”prime”} \\ \text{no} \Rightarrow \text{”composite”} \end{cases}$
if N is not prime $a^{N-1} = 1 \bmod N \leq$ half values of a < N

Lagrange’s Prime Theorem

Let $\pi(x)$ be the # of primes *leq* x, then
 $\pi(x) \approx \frac{x}{\ln(x)}$, or more precisely $\lim_{x \rightarrow \infty} \frac{\pi(x)}{(\frac{x}{\ln(x)})} = 1$

Modular Exponentiation

$x^y \bmod N \rightarrow$ start with repeated squaring mod N
 $x \bmod N \rightarrow x^2 \bmod N \rightarrow (x^2)^2 \dots x^{\log_2 y} \bmod N$
each step takes O(log²N) times to compute and there are log₂y steps, $\therefore \in O(n^3)$,
where n is the # of bits in N

Formal Limit Proof

$$\lim_{n \rightarrow \infty} \frac{f(n)}{g(n)} \begin{cases} \geq 0 \ (\infty) \Rightarrow f(n) \in \Omega(g(n)) \\ < \infty \ (0) \Rightarrow f(n) \in O(g(n)) \\ = c_{|0 < c < \infty} \Rightarrow f(n) \in \Theta(g(n)) \end{cases}$$

Logarithm Tricks

$$\log_b x^p = p \log_b x$$
$$\frac{\ln(x)}{\ln(m)} = \log_m x$$
$$x^{\log_b y} = y^{\log_b x}$$

Complexity

- $f \in O(g)$ if $f \leq c \cdot g$
- $f \in \Omega(g)$ if $f \geq c \cdot g$
- $f \in \Theta(g)$ if $f \in O(g)$ & $\Omega(g)$

Hierarchy:

- Exponential
- Polynomial
- Logarithmic
- Constant

Master’s Theorem

$$T(n) = aT(\frac{n}{b}) + O(n^d), \text{ if } a > 0, b > 1, d \geq 0$$

$$T(n) = \begin{cases} O(n^d) & \text{if } d > \log_b a \\ O(n^d \log_b n) & \text{if } d = \log_b a \\ O(n^{\log_b n}) & \text{if } d < \log_b a \end{cases}$$

Volker Strassen

faster matrix multiplication...

$$X = \begin{bmatrix} A & B \\ C & D \end{bmatrix} \times Y = \begin{bmatrix} E & F \\ G & H \end{bmatrix} = \begin{bmatrix} AE + BG & AF + BH \\ CE + DG & CF + DH \end{bmatrix}$$

$\in O(n^3)$ with recurrence $T(n)=8T(\frac{n}{2})+O(n^2)$

but thanks to Stassen...

$$XY = \begin{bmatrix} P_5 + P_4 - P_2 + P_6 & P_1 + P_2 \\ P_3 + P_4 & P_1 + P_5 - P_3 + P_7 \end{bmatrix}$$

$P_1 = A(F-H) \quad P_2 = (A+B)H \quad P_3 = (C+D)E \quad P_4 = D(G-E)$
 $P_5 = (A+D)(E+H) \quad P_6 = (B-D)(G+H) \quad P_7 = (A-C)(E+F)$
 $\in O(n^{\log_2 7}) \approx O(n^{2.81})$ with recurrence $T(n)=7T(\frac{n}{2})+O(n^2)$

Polynomial Multiplication

$$A(x) = a_0 + a_1x + \dots + a_dx^d \quad B(x) = b_0 + b_1x + \dots + b_dx^d$$
$$C(x)=A(x) \times B(x) = a_0b_k + a_1b_{k-1} + \dots + a_kb_0 = \sum_{i=0}^k a_ib_{k-i}$$

Fast Fourier Transform

complex nth roots of unity are given by $\omega = e^{\frac{2\pi i}{n}}, \omega^2, \omega^3, \dots$
< values > = FFT(< coefficients >, ω)
< coefficients > = $\frac{1}{n}$ FFT(< values >, ω^{-1}) $\in O(n \log n)$

Vandermonde Matrix, $M_n(\omega)=$

$$\begin{bmatrix} 1 & 1 & 1 & \dots & 1 \\ 1 & \omega & \omega^2 & \dots & \omega^{n-1} \\ 1 & \omega^2 & \omega^4 & \dots & \omega^{2(n-1)} \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 1 & \omega^{n-1} & \omega^{2(n-1)} & \dots & \omega^{(n-1)(n-1)} \end{bmatrix} \text{ where } (j,k)^{th} \text{ entry is } \omega^{jk}$$

Graphs

- **graph** – set of nodes & edges between select nodes
- **tree** – a connected graph with no cycles
- **back edge** – edge leading back to previously visited node
- **graph** – set of nodes & edges between select nodes

Depth First Search

discovers what nodes are reachable from a vertex $\in O(|V|+|E|)$

explore(G, v) :
 v.visit = true
 previsit(v)
 for each edge (v, u) in E:
 if u.visit = false: explore(G, u)
 postvisit(v)
dfs(G) :
 for all v \in V: v.visit = false
 for all v \in V: if v.visit = false:
 explore(G, v)