

PONV Data Extraction Template

Recording Post-Operative Nausea and Vomiting Outcomes

Instructions

Complete one row per patient. Record PONV risk factors on admission, intervention details peri-operatively, and outcomes at Recovery, Ward, and Discharge. Note: rescue antiemetic use is not equivalent to PONV — record both independently.

Patient Details and Risk Factors

Patient ID	Date	Surgery Type	Female	Non-smoker	Prior PONV / Motion Sick	Post-op Opioid	Risk Score (0–4)
			☐	☐	☐	☐	

Intervention and Outcomes

Patient ID	Intervention Type	Timing	Duration (min)	PONV Recovery	PONV Ward	PONV Discharge	Rescue Anti-emetic?	Drug / Dose	Notes
	WB / BA / EA / AU / MA / EAP	Pre / Post / Symptom		N / Mild / Mod / Sev	N / Mild / Mod / Sev	N / Mild / Mod / Sev	Y / N		

Legend

Intervention Type Codes:

- **WB** = Acupressure wristbands

PONV Severity Scale:

- **N** = None

- **BA** = Body acupuncture (PC6+)
- **EA** = Electroacupuncture
- **AU** = Auricular acupuncture
- **MA** = Manual acupressure
- **EAP** = Electroacupressure

- **Mild** = Nausea only, no vomiting
- **Mod** = Nausea with 1–2 vomiting episodes
- **Sev** = 3+ vomiting episodes or intractable nausea

Timing:

- **Pre** = Pre-operative (prevention)
- **Post** = Post-operative (prevention)
- **Symptom** = At symptom onset (management)

Collected	by:	Department:	Date	range:
-----------	-----	-------------	------	--------

PONV risk factors: Apfel et al. 1999. Note: rescue antiemetic use is not equivalent to PONV (track both). Template adapted from POPA4Ease data extraction tool.