

ArtSet

Art

Jrn

PubrN: DOVE Medical Press
JrnT: Int J Chron Obstruct Pulmon Dis
IsnO: 1178-2005
Vol: 12
Iss: 3
PubD [PubStatus="ppublish"]
Yr: 2017
Mo: 03

ArtTi: A case series of an "off-the-shelf" online health resource with integrated nurse:coaching to support self-management in COPD.

FPg: 2955

LPg: 2967

ELoc: 10.2147/COPD.S139532 [EIdType="doi" ValidYN="N"]

Lang: EN

AuL

Au

FName: Frances
LName: Early
Aff: Centre for Self Management Support, Cambridge University Hospitals NHS Foundation Trust, Cambridge, UK.

Au

FName: Jane
MName: S
LName: Young
Aff: Faculty of Health, Social Care and Education, School of Nursing and Midwifery, Anglia Ruskin University, Cambridge, UK.

Au

FName: Elizabeth
LName: Robinshaw
Affi
Aff: Gloucestershire Hospitals NHS Foundation Trust, Gloucester, UK.
Affi
Aff: Faculty of Health, Social Care and Education, School of Nursing and Midwifery, Anglia Ruskin University, Cambridge, UK.
Id: 0000-0001-6439-7613 [Source="ORCID"]

Au

FName: Ella Z
LName: Mi
Affi
Aff: School of Clinical Medicine, University of Cambridge, Cambridge, UK.
Id: <https://ror.org/013meh722> [Source="ROR"]

Au

FName: Jonathan P
LName: Fuld
Aff: Centre for Self Management Support, Cambridge University Hospitals NHS Foundation Trust, Cambridge, UK.

PubTy: Journal Article

ArtIdL

ArtId: copd-12-2955 [IdType="pii"]

ArtId: 10.2147/COPD.S139532 [IdType="doi"]

Hist

PubD [PubStatus="received"]
Yr: 2014
Mo: 09
Day: 30

PubD [PubStatus="accepted"]

Yr: 2014

Mo: 10

Day: 14

PubD [PubStatus="aheadofprint"]

Yr: 2015

Mo: 01

Day: 17

Abst

AbstTx: COPD has significant psychosocial impact. Self-management support improves quality of life, but programs are **not universally available**. IT-based self-management interventions can provide home-based support, but have mixed results. *We conducted a case series of an off-the-shelf Internet^{-based} health_{-promotion} program, The Preventive Plan (TPP), coupled with Nurse-coach support, which aimed to increase patient activation and provide self-management benefits.* [Label="BACKGROUND"]

AbstTx: A total of **19 COPD patients** were recruited, and 14 completed 3-month follow-up in two groups: **groups 1 and 2 or 3 with more and less advanced COPD**, respectively. Change in patient activation was determined with paired t-tests and Wilcoxon signed-rank tests. Benefits and user experience were explored in semistructured interviews, analyzed thematically. [Label="METHODS"]

AbstTx: Only group 1 improved significantly in activation, from a lower baseline than group 2; group 1 also improved significantly in mastery and anxiety. Both groups felt significantly more informed about COPD and reported physical functioning improvements. Group 1 reported improvements in mood and confidence. Overall, group 2 reported fewer benefits than group 1. Both groups valued nurse-coach support; for group 1, it was more important than TPP in building confidence to self-manage. The design of TPP and lack of motivation to use IT were barriers to use, but disease severity and poor IT skills were not. [Label="RESULTS"]

AbstTx: Our findings demonstrate the feasibility of combining nurse-coach support aligned to an Internet-based health resource, TPP, in COPD and provide learning about the challenges of such an approach and the importance of the nurse-coach role. [Label="DISCUSSION"]

OthAbst: An important part of managing COPD is self-management by the patient. With IT-based self-management programs, patients can do this from their own home, which is easier for them and more cost-effective for the health care provider. Our goal was to explore how a generic Internet-based health-promotion program, The Preventive Plan (TPP), containing programs to support behavior change and health information, combined with support from a specialist nurse, could help patients to manage their COPD... Our findings demonstrate that a low-intensity intervention can produce meaningful benefits, and provides learning about the challenges of this approach and the importance of the nurse coach role. [Language="eng" Type="plain-language-summary"]

Cpyr: Copyright© 2017 Early et al. This work is published and licensed by Dove Medical Press limited.

Co: The authors report no conflicts of interest in this work.

ObjL

Obj [Type="keyword"]

Prm: COPD [Name="value"]

Obj [Type="keyword"]

Prm: Internet [Name="value"]

Obj [Type="keyword"]

Prm: coaching [Name="value"]

Obj [Type="keyword"]

Prm: patient activation [Name="value"]

Obj [Type="grant"]

Prm: RO1DK561234 [Name="id"]

Prm: National Institutes of Health [Name="grantor"]

Obj [Type="grant"]

Prm: 2456797AB [Name="id"]

Prm: The British Granting Agency [Name="grantor"]

Prm: BGA [Name="acronym"]

Prm: England [Name="country"]

RefL

Ref

Cit: British Lung Foundation Chronic obstructive pulmonary disease (COPD) statistics. [Accessed January 27, 2017].

ArtIdL

ArtId: PMC4153410 [IdType="pmcid"]

ArtId: 24768240 [IdType="pubmed"]

Ref

Cit: Yohannes AM, Baldwin RC, Connolly MJ. Depression and anxiety in elderly patients with chronic obstructive pulmonary disease. Age Ageing. 2006;35(5):457–459.

ArtIdL

ArtId: PMC3020244 [IdType="pmcid"]

ArtId: 20932581 [IdType="pubmed"]