



# Table of Contents

## Confirmatory methods

### Generalized Order-Restricted Information Criterion (GORIC)

### GORIC weights

### Failsafe: Unconstrained or Complement

### GORICA

### Multiple Studies: Evidence synthesis

### End & Extra

# Confirmatory methods

## Confirmation

Limited set: Compare only prespecified hypotheses including order restrictions ( $<$ ,  $>$ , but also  $=$ ).

Most researchers are able to specify “order-restricted” / “informative” / “theory-based” hypotheses.

Use prior knowledge and/or expertise in hypothesis.

# ANOVA Example: Comparisons of 3 Means

Examine the difference in happiness  
between three types of “treatments”:  
(1) new treatment, (2) current treatment, and (3) no treatment.

Theory-based hypothesis:

$$H_1 : \quad \mu_1 > \mu_2 > \mu_3,$$

where “>” denotes “larger than”.

# Confirmatory methods

## Methods to evaluate theory-based hypotheses

- Hypothesis testing:  $F_{\text{bar}} (\bar{F})$  test  
(renders p-value and can test only one theory-based hypothesis)
- (Confirmatory) Bayesian model selection (BMS)
- Confirmatory model selection using information criteria:  
GORIC and GORICA

Confirmation more power than exploration.

- Kuiper, R. M., and Hoijtink, H. (2010). Comparisons of Means Using Exploratory and Confirmatory Approaches. *Psychological Methods*, 15(1), 69–86.
- Kuiper, R.M., Nederhof, T., and Klugkist, I. (2015). Properties of hypothesis testing techniques and (Bayesian) model selection for exploration-based and theory-based (order-restricted) hypotheses. *British Journal of Mathematical and Statistical Psychology*, 68(2), 220 – 245.

# Table of Contents

Confirmatory methods

Generalized Order-Restricted Information Criterion (GORIC)

GORIC weights

Failsafe: Unconstrained or Complement

GORICA

Multiple Studies: Evidence synthesis

End & Extra

# Information criteria (ICs)

IC, like GORIC, balances fit and complexity.

Describe data as good as possible (fit)  
with fewest number of parameters (simplicity / non-complexity).

# EXTRA: Generalized Order-Restricted Information Criterion

## GORIC

$$\text{'IC'} = -2 \text{ fit} + 2 \text{ complexity}$$

Fit = Maximized order-restricted log likelihood

Maximized log likelihood based on parameters in agreement with  $H_m$ .

Complexity = Penalty

Represents: Expected number of distinct parameters.

Here, expected number of distinct mean values plus 1 (because of the unknown variance term).

Details: Function of level probabilities.



# EXTRA: Idea complexity

loose interpretation

$$H_1 : \mu_1 > \mu_2 > \mu_3$$

contains 1 ordering of three means, 1-2-3.

Thus, not complex (i.e., parsimonious).

$$H_2 : \mu_1 > \mu_2, \mu_3$$

contains 2 orderings of three means: 1-2-3 and 1-3-2.

Thus, more complex (less parsimonious).

$$H_u : \mu_1, \mu_2, \mu_3$$

contains all six possible orderings of three means.

Thus, is most complex one (least parsimonious).

# EXTRA: Example GORIC

Palmer & Gough (2007) Data

$$H_0 : \mu_1 = \mu_2 = \mu_3,$$

$$H_1 : \mu_1 > \mu_2 > \mu_3,$$

$$H_2 : \mu_1 > \mu_2 < \mu_3,$$

$$H_3 : \mu_1 < \mu_2 < \mu_3,$$

$$H_u : \mu_1, \mu_2, \mu_3.$$

## GORIC

Model	Fit	Complexity	GORIC
$H_0$	-196.36	2.00	396.71
$H_1$	-191.89	2.81	389.41
$H_2$	-192.34	3.19	391.05
$H_3$	-196.36	2.81	398.34
$H_u$	-191.89	4.00	391.79

# EXTRA: GORIC

$$IC_m = -2 \text{ fit}_m + 2 \text{ complexity}_m$$

## Broad type of restrictions

More or less: any linear restriction.

e.g., the interaction  $H_1 : \mu_1 - \mu_2 < \mu_3 - \mu_4$ .

## Note

If no inequalities ( $<$  and/or  $>$ ), then (G)ORIC = AIC.

Reference:

- Kuiper, R.M., Hoijtink, H. and Silvapulle, M.J. (2011). An Akaike type information criterion for model selection under inequality constraints. *Biometrika*, 98, 495-501.
- Kuiper, R. M., Klugkist, I., and Hoijtink, H. (2010). A Fortran 90 Program for Confirmatory Analysis of Variance. *Journal of Statistical Software*, 34(8), 1–31.

# GORIC: Lowest value is best

GORIC is like AIC expected distance from the truth (KL-distance).  
Hence, smallest value is best.

# Table of Contents

Confirmatory methods

Generalized Order-Restricted Information Criterion (GORIC)

**GORIC weights**

Failsafe: Unconstrained or Complement

GORICA

Multiple Studies: Evidence synthesis

End & Extra

# Interpretation: GORIC weights

## GORIC values

GORIC values cannot be interpreted, only compared:  
Smallest is best.

## GORIC weights ( $w_m$ ) and ratios ( $w_m/w_{m'}$ )

$w_m$  quantifies how much  $H_m$  is more supported than others in set.  
 $w_m/w_{m'}$  quantifies relative support of  $H_m$  vs  $H_{m'}$ .  
The bigger, the better.

Reference:

Kuiper, R.M., Hoijtink, H. and Silvapulle, M.J. (2012). Generalization of the order restricted information criterion for multivariate normal linear models. *Journal of Statistical Planning and Inference*, 142, 2454-2463.

# Example GORIC weights ( $w_m$ )

Palmer & Gough (2007) Data

$$H_1 : \mu_1 > \mu_2 > \mu_3,$$

$$H_2 : \mu_2 > \mu_1 > \mu_3,$$

$$H_u : \mu_1, \mu_2, \mu_3.$$

## GORIC

Model	Fit	Complexity	GORIC	GORIC weights
$H_1$	-191.89	2.81	389.41	0.68
$H_2$	-193.70	2.81	393.03	0.11
$H_u$	-191.89	4.00	391.79	0.21

$H_1$  is  $.68/.11 \approx 6.1$  times more supported than competing hypothesis  $H_2$ .

# Table of Contents

Confirmatory methods

Generalized Order-Restricted Information Criterion (GORIC)

GORIC weights

Failsafe: Unconstrained or Complement

GORICA

Multiple Studies: Evidence synthesis

End & Extra



## Include “unconstrained” hypothesis

If set of hypotheses does not contain a reasonable/good one:  
Select the best of set of weak hypotheses.  
E.g.:  $w_1 = .8$  and  $w_2 = .2$ .

### Prevent choosing a weak hypothesis

Include unconstrained hypothesis  $H_u$  (highest fit but also most complex).  
E.g.:  $w_1 = .08$ ,  $w_2 = .02$ , and  $w_u = .90$ .

If at least one informative hypothesis not weak ( $w_1 > w_u$  or  $w_1/w_u > 1$ ),  
then compare informative hypotheses.

Hence:  $H_u$  is only a failsafe not another hypothesis of interest.  
See ‘Guidelines\_output\_GORIC.html’ on  
<https://github.com/rebeccakuiper/Tutorials>.  
On github site, go to Code (green button) and download zip.

# Alternative safeguard: Complement of $H_m$

## Alternatively (in case of one hypothesis of interest)

Evaluate hypothesis of interest against its complement;  
that is, all other possible hypotheses.

More powerful than against the unconstrained  
if  $H_m$  has maximum fit.

Reference:

Vanbrabant, L., Van Loey, N., and Kuiper, R. M. (2020). Evaluating a Theory-Based Hypothesis Against Its Complement Using an AIC-Type Information Criterion With an Application to Facial Burn Injury. *Psychological Methods*, 25(2), 129-142.  
<https://doi.org/10.1037/met0000238>

# Example $H_1$ vs $H_c$

Palmer & Gough (2007) Data

$$H_1 : \quad \mu_1 > \mu_2 > \mu_3,$$

$$H_c : \quad \text{not } H_1.$$

## GORIC

Model	Fit	Complexity	GORIC	GORIC weights
H1	-191.89	2.81	389.41	0.79
complement	-192.34	3.69	392.05	0.21

$H_1$  is  $.79/.21 \approx 3.8$  times more supported than its complement, that is, any other hypothesis.

# Table of Contents

Confirmatory methods

Generalized Order-Restricted Information Criterion (GORIC)

GORIC weights

Failsafe: Unconstrained or Complement

**GORICA**

Multiple Studies: Evidence synthesis

End & Extra

# GORICA

## GORIC: Normal linear models

GORIC can easily be applied to normal linear models (e.g., ANOVA models or regression models).

## GORIC: Other models

In case of other models (e.g., a SEM model), more cumbersome to calculate maximized order-restricted log likelihood and thus GORIC.

## GORICA: All models

Therefore, GORICA: asymptotic expression for GORIC.  
Can be used for all types of statistical models.

Reference:

Altınışık, Y., Van Lissa, C. J., Hoijtink, H., Oldehinkel, A. J., and Kuiper, R. M. (2021). Evaluation of inequality constrained hypotheses using a generalization of the AIC. *Psychological Methods*, 26(5), 599–621.

<https://doi.org/10.1037/met0000406>

# GORICA

## Similarities with GORIC

- Form:  $GORICA_m = -2 \text{ fit} + 2 \text{ complexity}$ .
- Broad type of restrictions.

## Differences compared to GORIC

- Uses asymptotic expression of the likelihood (is a normal):  
can therefore be easily applied to all types of models.  
Disadvantage: might work less well in case of small samples.
- Does not need data set; mle's and their covariance matrix suffice.
- Can leave out nuisance parameters (i.e., not part of hypotheses).

## Note

In case of normal linear models and/or not too small samples:  
GORICA weights = GORIC weights.

# Example GORICA

Palmer & Gough (2007) Data

$$H_1 : \mu_1 > \mu_2 > \mu_3,$$

$$H_c : \text{not } H_1.$$

## GORIC

Model	Fit	Complexity	GORIC	GORIC weights
H1	-1.96	1.81	7.55	0.79
complement	-2.39	2.69	10.15	0.21

$H_1$  is  $.79/.21 \approx 3.8$  times more supported than its complement, that is, any other hypothesis.

Note: GORIC weights are the same.

## Some literature

- Logistic Regression Modeling
  - Article: <https://doi.org/10.1037/met0000406>
- GORICA on SEM
  - Article:  
<https://www.tandfonline.com/doi/full/10.1080/10705511.2020.1836967>.
  - R scripts: [https://github.com/rebeccakuiper/GORICA\\_in\\_SEM](https://github.com/rebeccakuiper/GORICA_in_SEM).
- GORICA on cross-lagged panel model (CLPM) – Article:  
<https://doi.org/10.1111/bjep.12455>.
  - R scripts: [https://github.com/rebeccakuiper/GORICA\\_in\\_SEM](https://github.com/rebeccakuiper/GORICA_in_SEM).
- GORICA on Random-Intercept CLPM (RI-CLPM)
  - Article: Chuenjai Sukpan and Rebecca M. Kuiper (submitted 2023). How to evaluate causal dominance hypotheses in lagged effects models.
  - R scripts: <https://github.com/Chuenjai/Causal-dominance>.
- GORICA on CTmeta
  - Article: <https://doi.org/10.1080/10705511.2020.1823228>.
  - R scripts: [https://github.com/rebeccakuiper/GORICA\\_on\\_CTmeta](https://github.com/rebeccakuiper/GORICA_on_CTmeta).
- GORICA on cross-lagged panel model (CLPM)
  - Article: <https://doi.org/10.3390/e24111525>.
  - R scripts: [https://github.com/rebeccakuiper/GORICA\\_on\\_MetaAn](https://github.com/rebeccakuiper/GORICA_on_MetaAn).

**Note:** On github site, go to Code (green button) and download zip.



# Table of Contents

Confirmatory methods

Generalized Order-Restricted Information Criterion (GORIC)

GORIC weights

Failsafe: Unconstrained or Complement

GORICA

Multiple Studies: Evidence synthesis

End & Extra

# Possibilities multiple studies

- Update hypotheses.

First data set (or a part of it) generates one or more hypotheses.

Other data set (or part) used to determine evidence / support.

– html tutorial: [https://github.com/rebeccakuiper/Tutorials/blob/main/Tutorial\\_GORIC\\_restriktor\\_UpdateHypo.html](https://github.com/rebeccakuiper/Tutorials/blob/main/Tutorial_GORIC_restriktor_UpdateHypo.html)

– R script tutorial: [https://github.com/rebeccakuiper/Tutorials/blob/main/Hands-on%20files/Hands-on\\_4\\_GORIC\\_UpdateHypo\\_restriktor.R](https://github.com/rebeccakuiper/Tutorials/blob/main/Hands-on%20files/Hands-on_4_GORIC_UpdateHypo_restriktor.R)

- Aggregate evidence for hypotheses.

Aggregate the support for theories (diverse designs allowed).

Bear in mind: Meta-analysis aggregates parameter estimates or effect sizes which need to be comparable (often same designs required).

– html tutorial: [https://github.com/rebeccakuiper/Tutorials/blob/main/Tutorial\\_GORIC\\_restriktor\\_AggrSupport.html](https://github.com/rebeccakuiper/Tutorials/blob/main/Tutorial_GORIC_restriktor_AggrSupport.html)

– R script tutorial: [https://github.com/rebeccakuiper/Tutorials/blob/main/Hands-on%20files/Hands-on\\_5\\_GORICA\\_CombEv\\_restriktor.R](https://github.com/rebeccakuiper/Tutorials/blob/main/Hands-on%20files/Hands-on_5_GORICA_CombEv_restriktor.R)


**Note:** On github site, go to Code (green button) and download zip.

## GORIC(A) for Multiple Studies: Aggregating support (= evidence synthesis)

### References:

- Kuiper, R.M., Buskens, V.W., Raub, W., and Hoijtink, H. (2013). Combining statistical evidence from several studies: A method using Bayesian updating and an example from research on trust problems in social and economic exchange. *Sociological Methods and Research*, 42 (1), (pp. 60-81) (22 p.).
- Caspar J. Van Lissa, Eli-Boaz Clapper, and Rebecca Kuiper (submitted 2023). Aggregating evidence from conceptual replication studies using the product Bayes factor. [10.31234/osf.io/nvqpw](https://doi.org/10.31234/osf.io/nvqpw)
- Rebecca Kuiper and Eli-Boaz Clapper (to be submitted in 2023). GORIC Evidence Aggregation: Combining Statistical Evidence for a Central Theory from Diverse Studies using an AIC-type Criterion. [10.31234/osf.io/qv76x](https://doi.org/10.31234/osf.io/qv76x)


# Current best practice



Current best practice is meta-analysis and Bayesian updating.

- Not applicable for diverse research designs.
- Not applicable for incomparable estimates.

# Need for new methodology: Evidence Synthesis



Note: All studies do investigate the same theory (using diverse designs).

# Trust Example: Meta-Analysis versus Evidence Synthesis

Study	Type of model
1	univariate regression
2	univariate regression
3	probit regression
4	three-level logistic regression

Same design? e.g., same set of predictors?

Conceptual replications!

	Meta-Analysis	Evidence Synthesis
Effect size not required		✓
Deal with diverse designs		✓
Main results	Estimate of effect size	Evidence for hypotheses
Check:		same theoretical relationships?

Reference:

Kuiper, R.M., Buskens, V.W., Raub, W., and Hoijtink, H. (2013). Combining statistical evidence from several studies: A method using Bayesian updating and an example from research on trust problems in social and economic exchange.

*Sociological Methods and Research*, 42 (1), (pp. 60-81) (22 p.).

# Trust Example: Meta-Analysis versus Evidence Synthesis

Study	Type of model
1	univariate regression
2	univariate regression
3	probit regression
4	three-level logistic regression
Same design? e.g., same set of predictors?	

Conceptual replications!

	Meta-Analysis	Evidence Synthesis
Effect size not required		✓
Deal with diverse designs		✓
Main results	Estimate of effect size	Evidence for hypotheses
Check:		same theoretical relationships?

Reference:

Kuiper, R.M., Buskens, V.W., Raub, W., and Hoijtink, H. (2013). Combining statistical evidence from several studies: A method using Bayesian updating and an example from research on trust problems in social and economic exchange.

*Sociological Methods and Research*, 42 (1), (pp. 60-81) (22 p.).

# Trust Example: Meta-Analysis versus Evidence Synthesis

Study	Type of model
1	univariate regression
2	univariate regression
3	probit regression
4	three-level logistic regression
Same design? e.g., same set of predictors?	

## Conceptual replications!

	Meta-Analysis	Evidence Synthesis
Effect size not required		✓
Deal with diverse designs		✓
Main results	Estimate of effect size	Evidence for hypotheses
Check:		same theoretical relationships?

### Reference:

Kuiper, R.M., Buskens, V.W., Raub, W., and Hoijtink, H. (2013). Combining statistical evidence from several studies: A method using Bayesian updating and an example from research on trust problems in social and economic exchange. *Sociological Methods and Research*, 42 (1), (pp. 60-81) (22 p.).



# Trust Example: Meta-Analysis versus Evidence Synthesis

Study	Type of model
1	univariate regression
2	univariate regression
3	probit regression
4	three-level logistic regression
Same design? e.g., same set of predictors?	

Conceptual replications!

	Meta-Analysis	Evidence Synthesis
Effect size not required		✓
Deal with diverse designs		✓
Main results	Estimate of effect size	Evidence for hypotheses
<b>Check:</b>		<b>same theoretical relationships?</b>

Reference:

Kuiper, R.M., Buskens, V.W., Raub, W., and Hoijtink, H. (2013). Combining statistical evidence from several studies: A method using Bayesian updating and an example from research on trust problems in social and economic exchange.

*Sociological Methods and Research*, 42 (1), (pp. 60-81) (22 p.).

## Example: 4 studies regarding one concept

Study	Type of study	Number of observations $n$	Type of model
1	survey	895 transactions	univariate regression
2	experiment	348 decisions by 40 subjects	univariate regression
3	experiment	1249 decisions by 125 subjects	probit regression
4	experiment	2160 decisions by 144 subjects	three-level logistic regression
Study	Outcome $y$ ( <b>trust</b> )		scale $y$
1	effort invested in management		ratio
2	effort invested in management		ratio
3	choice of vignettes		dummy
4	trustfulness		dummy
Study	Predictor $x_1$ ( <b>past / previous experience</b> )		scale $x_1$
1	existence relationship with supplier		dummy
2	type of relationship with supplier		interval
3	bought a car from The Autosshop before		dummy
4	number of times a trustee honored trust in the past		ratio
Study	some of the other predictors		
1	transaction characteristics, expected future transactions, network embeddedness		
2	transaction characteristics, expected future transactions, network embeddedness		
3	expected future transactions, network embeddedness		
4	future interactions, network embeddedness		

# One-Parameter Example: Hypotheses of interest

## Parameter of interest in each study

parameter corresponding to  $x_1$  = previous experience; i.e.,  $\beta_1$ .  
For simplicity, only one here, could have been more.

## Expectation in each study

$x_1$  = previous experience has a positive effect on  $y$  = trust; i.e.,  $\beta_1 > 0$ .

## Set of central theories

$H_0$  : no effect,

$H_{>}$  : positive effect,

$H_{<}$  : negative effect.

Note 1: Central hypotheses for all studies, not w.r.t. average parameter.  
In data set, possibly study-specific hypotheses.

Note 2: In practice, I would not include  $H_0$ ...

# One-Parameter Example: Hypotheses of interest

## Parameter of interest in each study

parameter corresponding to  $x_1 =$  previous experience; i.e.,  $\beta_1$ .  
For simplicity, only one here, could have been more.

## Expectation in each study

$x_1 =$  previous experience has a positive effect on  $y =$  trust; i.e.,  $\beta_1 > 0$ .

## Set of central theories

$H_0$  : no effect,

$H_{>}$  : positive effect,

$H_{<}$  : negative effect.

Note 1: Central hypotheses for all studies, not w.r.t. average parameter.  
In data set, possibly study-specific hypotheses.

Note 2: In practice, I would not include  $H_0$ ...

# One-Parameter Example: Hypotheses of interest

## Parameter of interest in each study

parameter corresponding to  $x_1 =$  previous experience; i.e.,  $\beta_1$ .  
For simplicity, only one here, could have been more.

## Expectation in each study

$x_1 =$  previous experience has a positive effect on  $y =$  trust; i.e.,  $\beta_1 > 0$ .

## Set of central theories

$H_0$  : *no effect*,

$H_{>}$  : *positive effect*,

$H_{<}$  : *negative effect*.

Note 1: Central hypotheses for all studies, not w.r.t. average parameter.  
In data set, possibly study-specific hypotheses.

Note 2: In practice, I would not include  $H_0$ ...

## Example: Trust ( $y$ ) & previous experience ( $x_1$ )

Not full data set (and probit regression), so use

- GORICA (not GORIC) using *goric* function in R package *restriktor*

Input:

- parameter estimates and their covariance matrix

$t$	$\hat{\beta}_1$	$\hat{\sigma}_{\beta_1}$
1	0.090	0.029
2	0.140	0.054
3	1.090	0.093
4	1.781	0.179

Note: Here, one parameter ( $\beta_1$ ); thus, cov. matrix  $\hat{\beta}_1 = \text{variance } \hat{\beta}_1 = \hat{\sigma}_{\beta_1}^2$  (not  $\hat{\sigma}_{\beta_1}$ )

# One-Parameter Example: results per study

using GORICA

Results per study (not aggregated yet)!

**Table:** GORICA weights ( $w_{t,m}$ ) for Hypothesis  $H_m$  in Study  $t$

$m / t$	$w_{t,m}$			
	1	2	3	4
0	0.013	0.052	0.000	0.000
>	<b>0.979</b>	<b>0.916</b>	<b>1.000</b>	<b>1.000</b>
<	0.008	0.032	0.000	0.000

Note: Weight is at max 1.

So, now on forehand already clear.... but no quantification yet.

# One-Parameter Example: Results & Conclusions

using GORICA

Table: Overall GORICA weights ( $w_{t,m}^1$ ) for Hypothesis  $H_m$  in Study  $t$

$m / t$	$w_{t,m}^1$			
	1	2	3	4
0	0.013	0.001	0.000	0.000
>	<b>0.979</b>	<b>0.999</b>	<b>1.000</b>	<b>1.000</b>
<	0.008	0.000	0.000	0.000

- $w_{4,>}^1 = 1 \Rightarrow$  full support for  $H_>$   
 $w_{4,0}^1 = w_{4,<}^1 = 0 \Rightarrow$  no support for  $H_0$  and  $H_<$
- Support for  $H_>$  ( $w_{4,1}^1$ ) is highest: favor  $H_>$  over  $H_0$  and  $H_<$ .



# One-Parameter Example: Results & Conclusions

using GORICA

Table: Overall GORICA weights ( $w_{t,m}^1$ ) for Hypothesis  $H_m$  in Study  $t$

$m / t$	$w_{t,m}^1$			
	1	2	3	4
0	0.013	0.001	0.000	0.000
>	<b>0.979</b>	<b>0.999</b>	1.000	1.000
<	0.008	0.000	0.000	0.000

- $w_{4,>}^1 = 1 \Rightarrow$  full support for  $H_>$   
 $w_{4,0}^1 = w_{4,<}^1 = 0 \Rightarrow$  no support for  $H_0$  and  $H_<$
- Support for  $H_>$  ( $w_{4,1}^1$ ) is highest: favor  $H_>$  over  $H_0$  and  $H_<$ .

# One-Parameter Example: Results & Conclusions

using GORICA

Table: Overall GORICA weights ( $w_{t,m}^1$ ) for Hypothesis  $H_m$  in Study  $t$

$m / t$	$w_{t,m}^1$			
	1	2	3	4
0	0.013	0.001	0.000	0.000
>	<b>0.979</b>	<b>0.999</b>	<b>1.000</b>	<b>1.000</b>
<	0.008	0.000	0.000	0.000

- $w_{4,>}^1 = 1 \Rightarrow$  full support for  $H_>$   
 $w_{4,0}^1 = w_{4,<}^1 = 0 \Rightarrow$  no support for  $H_0$  and  $H_<$
- Support for  $H_>$  ( $w_{4,1}^1$ ) is highest: favor  $H_>$  over  $H_0$  and  $H_<$ .

# One-Parameter Example: Results & Conclusions

using GORICA

Table: Overall GORICA weights ( $w_{t,m}^1$ ) for Hypothesis  $H_m$  in Study  $t$

$m / t$	$w_{t,m}^1$			
	1	2	3	4
0	0.013	0.001	0.000	0.000
>	<b>0.979</b>	<b>0.999</b>	<b>1.000</b>	<b>1.000</b>
<	0.008	0.000	0.000	0.000

- $w_{4,>}^1 = 1 \Rightarrow$  full support for  $H_>$   
 $w_{4,0}^1 = w_{4,<}^1 = 0 \Rightarrow$  no support for  $H_0$  and  $H_<$
- Support for  $H_>$  ( $w_{4,1}^1$ ) is highest: favor  $H_>$  over  $H_0$  and  $H_<$ .

# One-Parameter Example: Results & Conclusions

using GORICA

Table: Overall GORICA weights ( $w_{t,m}^1$ ) for Hypothesis  $H_m$  in Study  $t$

$m / t$	$w_{t,m}^1$			
	1	2	3	4
0	0.013	0.001	0.000	0.000
>	<b>0.979</b>	<b>0.999</b>	<b>1.000</b>	<b>1.000</b>
<	0.008	0.000	0.000	0.000

- $w_{4,>}^1 = 1 \quad \Rightarrow \quad \text{full support for } H_>$   
 $w_{4,0}^1 = w_{4,<}^1 = 0 \quad \Rightarrow \quad \text{no support for } H_0 \text{ and } H_<$
- Support for  $H_>$  ( $w_{4,1}^1$ ) is highest: favor  $H_>$  over  $H_0$  and  $H_<$ .

# One-Parameter Example: Results & Conclusions

using GORICA

Table: Overall GORICA weights ( $w_{t,m}^1$ ) for Hypothesis  $H_m$  in Study  $t$

$m / t$	$w_{t,m}^1$			
	1	2	3	4
0	0.013	0.001	0.000	0.000
$>$	<b>0.979</b>	<b>0.999</b>	<b>1.000</b>	<b>1.000</b>
$<$	0.008	0.000	0.000	0.000

- $w_{4,>}^1 = 1 \quad \Rightarrow \quad \text{full support for } H_>$   
 $w_{4,0}^1 = w_{4,<}^1 = 0 \quad \Rightarrow \quad \text{no support for } H_0 \text{ and } H_<$
- Support for  $H_>$  ( $w_{4,1}^1$ ) is highest: favor  $H_>$  over  $H_0$  and  $H_<$ .

## Possible type of sets of studies

- Conceptual replications of same authors, done as a robustness check.
- Searching for direct and indirect/conceptual replications in the literature.
- Using different cohorts, where one can measure the variables in their own way.
- Using different subpopulations, possibly using different operationalisations.
- ...

## EXTRA: Possible type of sets of studies (1/2)

- Conceptual replications of same authors, done as a robustness check. This was done in the Trust example of Buskens and Raub.  
Note: There, the central hypotheses regard one parameter of interest, but one can inspect multiple parameters and also compare them.
- Searching for direct and indirect/conceptual replications in the literature.  
E.g., using the central hypothesis that the absolute strength of the relationship of communication competence (C) with willingness to communicate in a second language (WTC) is greater than the absolute strength of the relation of communication anxiety (A) with WTC, which is greater than the absolute strength of the relation of motivation (M) with WTC, that is,  $|C| > |A| > |M|$ .
  - ‘Article’: Example in bachelor thesis of Martijn Sips
  - R scripts: <https://github.com/rebeccakuiper/Tutorials/tree/main/Examples%20evSyn/Example%20WtC>

**Note:** On github site, go to Code (green button) and download zip

## EXTRA: Possible type of sets of studies (2/2)

- Using different cohorts, where one can measure the variables in their own way.

So, possibly using different operationalisation of the same constructs.

– Article: Zondervan-Zwijnenburg et al. (2020).

Uses cohorts: Gen-R, NTR, and TRAILS.

- Using different subpopulations, possibly using different operationalisations.

E.g., the Municipal Health Services (Dutch acronym: GGD) studied the positive consequences of corona on loneliness (a), mental health (b), and stress (b) (conditional on sex, age, and health); they expected  $a < b < c$ .

– Article: in progress

– R scripts: <https://github.com/rebeccakuiper/Tutorials/tree/main/Examples%20evSyn/Example%20corona%20GGD>

**Note:** On github site, go to Code (green button) and download zip.



# Table of Contents

Confirmatory methods

Generalized Order-Restricted Information Criterion (GORIC)

GORIC weights

Failsafe: Unconstrained or Complement

GORICA

Multiple Studies: Evidence synthesis

End & Extra

# Software

Currently, beta versions of software:

- R function *evSyn* in R package *restriktor*
- Interactive web application (Shiny app) of *GoricEvSyn*

## Websites

<https://github.com/rebeccakuiper/Tutorials>

[www.uu.nl/staff/RMKuiper/Software](http://www.uu.nl/staff/RMKuiper/Software)

[www.uu.nl/staff/RMKuiper/Websites%20%2F%20Shiny%20apps](http://www.uu.nl/staff/RMKuiper/Websites%20%2F%20Shiny%20apps)

[informative-hypotheses.sites.uu.nl/software/goric/](http://informative-hypotheses.sites.uu.nl/software/goric/)

## Promo: Publish in special issue

### Applying GORIC(A)?

If you collected data or will collect,  
if you have one or more a-priori hypotheses,  
then you can apply the GORIC(A).

### FYI: Special issue

Journal (mdpi): Mathematics (ISSN 2227-7390)

Special Issue:

*Evaluation of Theory-Driven Hypotheses: No Hypothesis, No G(L)ORIC*

Guest Editor: Rebecca M. Kuiper

[https://www.mdpi.com/journal/mathematics/special\\_issues/97C26430R3](https://www.mdpi.com/journal/mathematics/special_issues/97C26430R3)

Contact: [r.m.kuiper@uu.nl](mailto:r.m.kuiper@uu.nl)

Please contact me ([r.m.kuiper@uu.nl](mailto:r.m.kuiper@uu.nl)), if you want to explore the possibilities (and to obtain discount).

# The End

Thanks for listening!

Are there any questions?

## Websites

<https://github.com/rebeccakuiper/Tutorials>  
[www.uu.nl/staff/RMKuiper/Software](http://www.uu.nl/staff/RMKuiper/Software)  
[www.uu.nl/staff/RMKuiper/Websites%20%2F%20Shiny%20apps](http://www.uu.nl/staff/RMKuiper/Websites%20%2F%20Shiny%20apps)  
[informative-hypotheses.sites.uu.nl/software/goric/](http://informative-hypotheses.sites.uu.nl/software/goric/)

## E-mail

[r.m.kuiper@uu.nl](mailto:r.m.kuiper@uu.nl)