

Informative hypotheses evaluation

R. M. Kuiper

Department of Methodology & Statistics
Utrecht University

Table of Contents

Example Null hypothesis (H_0) vs Informative hypothesis (H_m)

GORIC weights

Examples with R code

Failsafe hypotheses: Unconstrained & Complement

GORICA (comparable to bain)

GORIC(A) in JASP

End & Extra

ANOVA Example: Comparisons of 3 Means

Palmer and Gough (2007) examined the difference in the attribution of importance of defective education as an explanation for criminal behavior between three types of “offenders”:

(1) non-offenders, (2) property offenders, and (3) person offenders.

ANOVA

$$y_j = \mu_1 d_{1j} + \mu_2 d_{2j} + \mu_3 d_{3j} + \epsilon_j,$$

where μ_i ($i = 1, 2, 3$) is the mean in group i ,
 d_{ij} is a dummy variable denoting group membership,
 ϵ_j is an error term and $\epsilon_j \sim N(0, \sigma^2)$.

$(n_i = 20, 20, 31)$

Hypotheses of interest

Palmer and Gough (2007) expect that

- non-offenders (1) attribute more importance to defective education for explaining crime than the other two offenders;
i.e., $\mu_1 > \mu_2$ and $\mu_1 > \mu_3$,
- property offenders (2) attribute more importance to defective education than person offenders (3);
i.e., $\mu_2 > \mu_3$.

$$H_1 : \quad \mu_1 > \mu_2 > \mu_3.$$

Note that ‘<’ denotes “smaller than” and ‘>’ denotes “larger than”.

Conclusions w.r.t. hypothesis of interest $H_1 : \mu_1 > \mu_2 > \mu_3$

With post-hoc test (e.g., Bonferroni) and on basis of the sample means, we could say a bit more.

But, what if

- $\mu_1 \neq \mu_2, \mu_1 \neq \mu_3, \mu_2 = \mu_3,$
- $\bar{y}_1 > \bar{y}_2,$

which implies $\mu_1 > \mu_2 = \mu_3$.

But also not interested in this.

Furthermore, inconsistent results are possible (like in this data example):

$$\mu_1 \neq \mu_3, \mu_1 = \mu_2, \text{ and } \mu_2 = \mu_3,$$

which is logically impossible, since latter two imply $\mu_1 = \mu_3$.

Using planned contrasts would overcome this, but still = restrictions then.

NHST on Palmer & Gough (2007) Data

```
PandG_data <- read.table("Data_PalmerAndGough.txt",  
                          header=TRUE)  
PandG_data$group <- factor(PandG_data$group)  
pairwise.t.test(PandG_data$Importance, PandG_data$group,  
                p.adj = 'bonferroni')  
  
##  
## Pairwise comparisons using t tests with pooled SD  
##  
## data: PandG_data$Importance and PandG_data$group  
##  
##      1      2  
## 2 0.191 -  
## 3 0.011 1.000  
##  
## P value adjustment method: bonferroni
```


Confirmatory methods

Most researchers are able to specify “order-restricted” / “informative” / “theory-based” hypotheses, like $H_1 : \mu_1 > \mu_2 > \mu_3$.
Use prior knowledge and/or expertise in hypothesis.

Methods to evaluate theory-based hypotheses

- Hypothesis testing: $F_{\text{bar}} (\bar{F})$ test
(renders p-value and can test only one theory-based hypothesis)
- Confirmatory model selection using information criteria:
GORIC and GORICA
- (Confirmatory) Bayesian model selection (BMS)

GORIC on Palmer & Gough (2007) Data

$$H_0 : \mu_1 = \mu_2 = \mu_3,$$

$$H_1 : \mu_1 > \mu_2 > \mu_3,$$


$$H_u : \mu_1, \mu_2, \mu_3.$$

GORIC


Model	Fit	Complexity	GORIC	GORIC weights
H_0	-196.36	2.00	396.71	0.01
H_1	-191.89	2.81	389.41	0.50
H_u	-191.89	4.00	391.79	0.50

Confirmatory methods have more “power” than their exploratory counterparts (cf. one-sided testing).

Confirmation more power: 1 data set. GORIC values for 3 groups, effect size ES , and $n = 10$ observations per group



Confirmation more power: 1 data set. GORIC values for 3 groups, effect size ES , and $n = 100$ observations per group



Confirmatory methods - some of my references

- Kuiper, R. M., and Hoijtink, H. (2010). Comparisons of Means Using Exploratory and Confirmatory Approaches. *Psychological Methods*, 15(1), 69–86.
- Kuiper, R. M., Klugkist, I., and Hoijtink, H. (2010). A Fortran 90 Program for Confirmatory Analysis of Variance. *Journal of Statistical Software*, 34(8), 1–31.
- Kuiper, R.M., Hoijtink, H. and Silvapulle, M.J. (2011). An Akaike type information criterion for model selection under inequality constraints. *Biometrika*, 98, 495-501. (GORIC)
- Kuiper, R.M., Nederhof, T., and Klugkist, I. (2015). Properties of hypothesis testing techniques and (Bayesian) model selection for exploration-based and theory-based (order-restricted) hypotheses. *British Journal of Mathematical and Statistical Psychology*, 68(2), 220 – 245.
- Altınışik, Y., Van Lissa, C. J., Hoijtink, H., Oldehinkel, A. J., and Kuiper, R. M. (2021). Evaluation of inequality constrained hypotheses using a generalization of the AIC. *Psychological Methods*, 26(5), 599-621. (GORICA)

Table of Contents

Example Null hypothesis (H_0) vs Informative hypothesis (H_m)

Generalized Order-Restricted Information Criterion (GORIC)

GORIC weights

Examples with R code


Failsafe hypotheses: Unconstrained & Complement

GORICA (comparable to b_{ain})

GORIC(A) in JASP

End & Extra

Intermezzo: Balance Fit and Complexity




$$\mu_1 = \mu_2 = \mu_3 = \mu_4 = \mu_5 = \mu$$

parameters: 1 mean parameter (i.e., μ)

Fit: Bad fit

Intermezzo: Balance Fit and Complexity



$$\mu_1 = \mu_2 = \mu_3 = \mu_{\text{low}}$$

$$\mu_4 = \mu_5 = \mu_{\text{high}}$$


parameters:

2 mean parameters (i.e., μ_{low} and μ_{high})

Fit:

Better than with one parameter, even good fit.

Intermezzo: Balance Fit and Complexity




Now, we also incorporate order-restrictions (e.g., $\mu_1 < \mu_2$).

Then, helpful to look at likelihood (not scatter plot).

In presentation, I will use contour plots (in case of two means, not five).

Idea fit


order-restricted maximum likelihood (or-ml)



mle not in H_2 : Find highest likelihood in allowable (= green) space.
The resulting estimated mean $\hat{\mu} = (\hat{\mu}_1, \hat{\mu}_2)$ is referred to as or-ml.

Idea complexity

penalty (PT)



Note: sampling from null distribution: $H_0: \mu_1 = \mu_2 = 0$.

complexity $H_2 = PT_2 = 1 + 0.5 \times 1 + 0.5 \times 2 = 2.5$.

PT = number of expected distinct parameters (under H_0).

Idea complexity

loose interpretation

$$H_1 : \mu_1 > \mu_2 > \mu_3$$

contains 1 ordering of three means, 1-2-3.

Thus, not complex (i.e., parsimonious).

$$H_2 : \mu_1 > \mu_2, \mu_3$$

contains 2 orderings of three means: 1-2-3 and 1-3-2.

Thus, more complex (less parsimonious).

$$H_u : \mu_1, \mu_2, \mu_3$$

contains all six possible orderings of three means.

Thus, is most complex one (least parsimonious).

GORIC on Palmer & Gough (2007) Data

$$H_0 : \mu_1 = \mu_2 = \mu_3,$$

$$H_1 : \mu_1 > \mu_2 > \mu_3,$$

$$H_2 : \mu_1 > \mu_2 < \mu_3,$$

$$H_3 : \mu_1 < \mu_2 < \mu_3,$$

$$H_u : \mu_1, \mu_2, \mu_3.$$

GORIC

Model	Fit	Complexity	GORIC	GORIC weights
H_0	-196.36	2.00	396.71	0.00
H_1	-191.89	2.81	389.41	0.38
H_2	-192.34	3.19	391.05	0.24
H_3	-196.36	2.81	398.34	0.00
H_u	-191.89	4.00	391.79	0.38

GORIC: Lowest value is best

GORIC is like AIC expected distance from the truth (KL-distance). Hence, smallest value is best.

GORIC on Palmer & Gough (2007) Data

$$H_0 : \mu_1 = \mu_2 = \mu_3,$$

$$H_1 : \mu_1 > \mu_2 > \mu_3,$$

$$H_u : \mu_1, \mu_2, \mu_3.$$

GORIC

Model	Fit	Complexity	GORIC	GORIC weights
H_0	-196.36	2.00	396.71	0.01
H_1	-191.89	2.81	389.41	0.50
H_u	-191.89	4.00	391.79	0.50

Hands-on/Demo (1): GORIC

Start Rstudio and let's practice.

- Go to <https://github.com/rebeccakuiper/Tutorials>:
 1. Click on green button called Code.
 2. Download zip (last option in list).
 3. Unzip it on your machine (that folder is now your working directory).
- Optional: Open 'Tutorial_GORIC_restriktor_ANOVA.html'.
- Start R (optional: make project).
Optional: open 'PalmerAndGough_and_Lucas.R'.
- Install packages and load them.
- Read and inspect data. Use Data_PalmerAndGough.txt.
- Run model (lm()).
- Specify hypotheses (make up your own).
Note: Use names used in the model, or overwrite those.
- Run goric().
- Inspect and interpret output.
Note: 'GORIC weights' will be explained later on.

Interpretation: GORIC weights

GORIC values

GORIC values cannot be interpreted, only compared:
Smallest is best.

GORIC weights (w_m) and ratios ($w_m/w_{m'}$)

GORIC weight quantifies how much H_m is more supported than others in set.

Ratio of GORIC weights quantifies relative support of H_m vs $H_{m'}$.

The bigger, the better.

Reference:

Kuiper, R.M., Hoijtink, H. and Silvapulle, M.J. (2012). Generalization of the order restricted information criterion for multivariate normal linear models. *Journal of Statistical Planning and Inference*, 142, 2454-2463.

Illustration of the GORIC weights (w_m)

$$H_0 : \mu_1 = \mu_2 = \mu_3,$$

$$H_1 : \mu_1 > \mu_2 > \mu_3,$$

$$H_u : \mu_1, \mu_2, \mu_3.$$

GORIC

Model	Fit	Complexity	GORIC	GORIC weights
H_0	-196.36	2.00	396.71	0.01
H_1	-191.89	2.81	389.41	0.50
H_u	-191.89	4.00	391.79	0.50

Note on GORIC weights vs BF and PMPs

ratio GORIC weights ($w_m/w_{m'}$) \sim Bayes factor ($BF_{mm'}$).

GORIC weight (w_m) \sim posterior model probability (PMP).

1 - w_m = conditional error probability.

Like PMP, w_m depends on set of hypotheses.

Example

$$w_m = .8$$

$$w_{m'} = .2$$

H_m is $.8/.2 = 4$ times more supported than $H_{m'}$.

There is an error probability of $(1 - .8) * 100\% = 20\%$ that H_m is true.

Table of Contents

Example Null hypothesis (H_0) vs Informative hypothesis (H_m)

Generalized Order-Restricted Information Criterion (GORIC)

GORIC weights

Examples with R code

Failsafe hypotheses: Unconstrained & Complement

GORICA (comparable to bain)

GORIC(A) in JASP

End & Extra

Palmer & Gough

```
fit.PandG <- lm(Importance ~ group - 1, data = PandG_data)
```


R code: GORIC

Palmer & Gough

$$H_0 : \mu_1 = \mu_2 = \mu_3$$

$$H_1 : \mu_1 = \mu_2 > \mu_3$$

$$H_u : \quad \mu_1, \mu_2, \mu_3$$

```
# GORIC (using goric function in restriktor package)
library(restriktor)
set.seed(123) # Set seed value
goric.PandG <- goric(fit.PandG,
                     constraints = list(H0, H1))
```


GORIC on Palmer & Gough Data

$H_0 :$
 $\mu_1 = \mu_2 = \mu_3$

$H_1 :$
 $\mu_1 = \mu_2 > \mu_3$

$H_u :$
 μ_1, μ_2, μ_3

GORIC

Model	Fit	Complexity	GORIC	GORIC weights
H_0	-196.36	2.00	396.71	0.01
H_1	-191.89	2.81	389.41	0.50
H_u	-191.89	4.00	391.79	0.50

Another illustration: one-way ANOVA

Lucas: 5 groups

Lucas (2003) investigated difference between female and male leadership w.r.t. influence of the leader.

Five experimental groups:

1. a randomly selected male leader
2. a randomly selected female leader
3. male leader selected via task
4. female leader selected via task
5. female leader selected via task + institutionalized female leadership via movie

R code: Data and fit object

Lucas

```
# Data
```

```
lucas_data <- read.table("Data_Lucas.txt", header=TRUE)
```

```
lucas_data$group <- factor(lucas_data$group)
```

```
# fit object (needed as input for goric function)
```

```
fit.lucas <- lm(Influence ~ group - 1, data = lucas_data)
```

R code: Hypotheses

Lucas

$$H_0: \mu_1 = \mu_2 = \mu_3 = \mu_4 = \mu_5,$$

$$H_1: \mu_5 = \mu_3 > \{\mu_1, \mu_4\} > \mu_2$$

$$H_2: \mu_3 > \mu_1 > \mu_4 = \mu_5 > \mu_2,$$

$$H_u : \quad \mu_1, \mu_2, \mu_3, \mu_4, \mu_5.$$

Hypotheses

```
H0 <- 'group1 = group2 = group3 = group4 = group5'
```

```
H1 <- 'group5 = group3 > group1 > group2;  
      group3 > group4 > group2'
```

```
H2 <- 'group3 > group1 > group4 = group5 > group2'
```


GORIC on Lucas' Data

$$H_0 : \mu_1 = \mu_2 = \mu_3 = \mu_4 = \mu_5,$$

$$H_1 : \mu_5 = \mu_3 > \{\mu_1, \mu_4\} > \mu_2$$

$$H_2 : \mu_3 > \mu_1 > \mu_4 = \mu_5 > \mu_2,$$

$$H_u : \mu_1, \mu_2, \mu_3, \mu_4, \mu_5.$$

GORIC

Model	Fit	Complexity	GORIC	GORIC weights
H_0	-292.27	2.00	588.54	0.00
H_1	-278.05	3.20	562.49	0.49
H_2	-281.76	3.14	569.79	0.01
H_u	-278.05	6.00	568.10	0.49

GORIC on Lucas' Data: GORIC weights (w_m)

GORIC

Model	Fit	Complexity	GORIC	GORIC weights
H_0	-292.27	2.00	588.54	0.00
H_1	-278.05	3.20	562.49	0.49
H_2	-281.76	3.14	569.79	0.01
H_u	-278.05	6.00	568.10	0.49

H_1 is $.921/.024 \approx 38.6$ (∞) times more supported than H_2 (H_0 , resp.). Thus, there is quite some evidence that H_1 is the best of this set.

Note:

H_1 is $.921/.056 \approx 16.5 > 1$ times more supported than H_ν .

Hence, H_1 not weak and thus comparable to the other informative hypotheses (as explained next).

Hands-on/Demo (2): GORIC weights

Start Rstudio and let's practice.

- Optional: Open 'Tutorial_GORIC_restriktor_ANOVA.html'.
- If needed: Start R again (optional: make project) and then also load packages again.
Optional: open 'PalmerAndGough_and_Lucas.R'.
- Read and inspect data.
Use Data_Lucas.txt and/or Data_PalmerAndGough.txt.
- Run model (lm()).
- Specify hypotheses (make up your own). Note: Use names used in the model, or overwrite those.
- Run goric().
- Inspect and interpret output: Focus on GORIC weights.

Hands-on/Demo: EXTRA - make it crash :-)

Extra for restriktor from github:

```

if (!require("devtools")) install.packages("devtools")
library(devtools)
install_github("LeonardV/restriktor")
#install_github("LeonardV/restriktor", force = T)
library(restriktor) # for goric() and GoricEvSyn() function
# Difference:
# CRAN:    goric(fit.object, H1, H2)
# Github: goric(fit.object, constraints = list(H1, H2))
  
```

- Check on restrictions using brackets ().
- Check goric on lm() with continuous variable with missing data (also with auxiliary variables).
- Check evidence synthesis functions (possibly explained later).

! Possible SA (also updating tutorials, creating extra tutorials, and searching for articles to be used in evidence synthesis) !

If suggestions or errors

- Check on restrictions using brackets ().
- Check goric on `lm()` with continuous variable with missing data (also with auxiliary variables).
- Check evidence synthesis functions (possibly explained later).

If suggestions or errors

Email L.Vanbrabant@ggdwestbrabant.nl

Send: `suggestion(s)` and/or errors + code + data

If SA

Email R.M.Kuiper@uu.nl

Table of Contents

Example Null hypothesis (H_0) vs Informative hypothesis (H_m)

Generalized Order-Restricted Information Criterion (GORIC)

GORIC weights

Examples with R code

Failsafe hypotheses: Unconstrained & Complement

GORICA (comparable to bain)

GORIC(A) in JASP

End & Extra

Include “unconstrained” hypothesis

If set of hypotheses does not contain a reasonable/good one:
Select the best of set of weak hypotheses.
E.g.: $w_0 = .2$ and $w_1 = .8$.

Prevent choosing a weak hypothesis

Include unconstrained hypothesis H_u (or H_a):

$$\begin{aligned} H_0 : & \quad \mu_1 = \mu_2 = \mu_3, \\ H_1 : & \quad \mu_1 > \mu_2 > \mu_3, \\ H_u : & \quad \mu_1, \mu_2, \mu_3, \\ & \quad (\text{i.e., no restrictions}). \end{aligned}$$

H_u highest fit but also most complex, thus failsafe.

E.g.: $w_0 = .02$, $w_1 = .08$, and $w_\mu = .90$.

Use of H_u : Palmer & Gough (2007) Data

$$H_0 : \mu_1 = \mu_2 = \mu_3,$$

$$H_1 : \mu_1 > \mu_2 > \mu_3,$$

$$H_u : \mu_1, \mu_2, \mu_3.$$

GORIC

Model	Fit	Complexity	GORIC	GORIC weights
H_0	-196.36	2.00	396.71	0.01
H_1	-191.89	2.81	389.41	0.50
H_u	-191.89	4.00	391.79	0.50

If at least one informative hypothesis not weak ($w_1 > w_u$ or $w_1/w_u > 1$), then compare informative hypotheses.

Hence: H_u is only a failsafe not another hypothesis of interest.

H_1 vs H_u : Palmer & Gough (2007) Data

What if only one informative hypothesis:

$$H_1 : \quad \mu_1 > \mu_2 > \mu_3,$$

$$H_u : \quad \mu_1, \mu_2, \mu_3.$$

GORIC

Model	Fit	Complexity	GORIC	GORIC weights
H_1	-191.89	2.81	389.41	0.5
H_u	-191.89	4.00	391.79	0.5

H_1 is $.77/.23 \approx 3.3$ times more supported than H_u .

BUT: H_u includes H_1 .

So, support for H_u contains support for H_1 .

H_1 vs H_u : Palmer & Gough (2007) Data

GORIC

Model	Fit	Complexity	GORIC	GORIC weights
H_1	-191.89	2.81	389.41	0.5
H_u	-191.89	4.00	391.79	0.5

Unconstrained is always true, so will always receive support.
Support for H_u contains support for H_1 (if any).

Now: Fit H_1 and H_u the same;
thus, GORIC weights are based on only complexity.

Consequently, $.77/.23 \approx 3.3$ is an upper bound.

Alternative safeguard: Complement of H_m using GORIC

Alternatively (in case of one hypothesis of interest)

Evaluate hypothesis of interest against its complement;
that is, all other possible hypotheses.


More powerful than against the unconstrained
if H_m has maximum fit.

Reference:


Vanbrabant, L., Van Loey, N., and Kuiper, R. M. (2020). Evaluating a Theory-Based Hypothesis Against Its Complement Using an AIC-Type Information Criterion With an Application to Facial Burn Injury. *Psychological Methods*, 25(2), 129-142.
<https://doi.org/10.1037/met0000238>

Alternative safeguard: Complement of H_m using GORIC

vs complement




(a)




(b)

vs unconstrained



(c)



(d)

R code: complement

Palmer & Gough

$$H_1 : \quad \mu_1 > \mu_2 > \mu_3,$$

$$H_c : \quad \text{not } H_1.$$

GORIC

```
H1 <- 'group1 > group2 > group3'  
# GORIC (using goric function in restriktor package)  
library(restriktor)  
set.seed(123) # Set seed value  
goric.lucas_C <- goric(fit.PandG,  
                        constraints = list(H1),  
                        comparison = 'complement')
```

Palmer & Gough (2007): H_1 vs H_c

$$H_1 : \quad \mu_1 > \mu_2 > \mu_3,$$

$$H_c : \quad \text{not } H_1.$$

GORIC

Model	Fit	Complexity	GORIC	GORIC weights
H1	-191.89	2.81	389.41	0.61
complement	-192.34	3.69	392.05	0.39

H_1 is $.79/.21 \approx 3.8$ times more supported than its complement, that is, any other hypothesis.

Hands-on/Demo (3): GORIC - H_m vs complement

Start Rstudio and let's practice.

- Optional: Open 'Tutorial_GORIC_restriktor_ANOVA.html'.
- If needed: Start R again (optional: make project) and then also load packages again.
- Read and inspect data.
Use Data_Lucas.txt and/or Data_PalmerAndGough.txt.
- Run model (`lm()`).
- Specify hypotheses (make up your own). E.g., for Lucas:

$$H_1 : \quad \mu_5 = \mu_3 > \{\mu_1, \mu_4\} > \mu_2$$

Note: Use names used in the model, or overwrite those.

- Run `goric()`; now, add:
`comparison = "complement"`

- Inspect and interpret output.

GORICA

GORIC: Normal linear models

GORIC can easily be applied to normal linear models (e.g., ANOVA models or regression models).

GORIC: Other models

In case of other models (e.g., a SEM model), more cumbersome to calculate maximized order-restricted log likelihood and thus GORIC.

GORICA: All models

Therefore, GORICA: asymptotic expression for GORIC.
Can be used for all types of statistical models.

Reference:

Altınışık, Y., Van Lissa, C. J., Hoijtink, H., Oldehinkel, A. J., and Kuiper, R. M. (2021). Evaluation of inequality constrained hypotheses using a generalization of the AIC. *Psychological Methods*, 26(5), 599-621. (GORICA)

GORICA

Similarities with GORIC

- Form: $GORICA_m = -2 \text{ fit} + 2 \text{ complexity}$.
- Broad type of restrictions.

Differences compared to GORIC

- Uses asymptotic expression of the likelihood (is a normal):
can therefore be easily applied to all types of models.
Disadvantage: might work less well in case of small samples.
- Does not need data set; mle's and their covariance matrix suffice.
- Can leave out nuisance parameters (i.e., not part of hypotheses).

Note

In case of normal linear models and/or not too small samples:
GORICA weights = GORIC weights.

R code: GORICA

Palmer & Gough

$$H_1 : \quad \mu_1 > \mu_2 > \mu_3,$$

$$H_c : \quad \text{not } H_1.$$

GORICA: type = "gorica"

```
H1 <- 'group1 > group2 > group3'  
# GORICA (using goric function in restriktor package)  
library(restriktor)  
set.seed(123) # Set seed value  
gorica.lucas_C <- goric(fit.PandG,  
                        constraints = list(H1),  
                        comparison = 'complement',  
                        type = "gorica")
```


Hands-on/Demo (4): GORICA - H_m vs complement

Start Rstudio and let's practice.

- Optional: Open 'Tutorial_GORIC_restriktor_ANOVA.html'.
- If needed: Start R again (optional: make project) and then also load packages again.
- Read and inspect data.
Use Data_Lucas.txt and/or Data_PalmerAndGough.txt.
- Run model (`lm()`).
- Specify hypotheses (make up your own). E.g., for Lucas:

$$H_1 : \quad \mu_5 = \mu_3 > \{\mu_1, \mu_4\} > \mu_2$$

Note: Use names used in the model, or overwrite those.

- Run `goric()`; now, add:

```
type = "gorica"
```

- Inspect and interpret output.

Table of Contents

GORIC weights

Examples with R code

Failsafe hypotheses: Unconstrained & Complement

GORICA (comparable to bain)

GORIC(A) in JASP

End & Extra

GORIC in JASP: Palmer & Gough

- Open JASP and 'Data_PalmerAndGough_JASP.txt'. Check measurement levels.
- Go to ANOVA and denote variables of interest.
- Go to Order Restrictions tab.
- Specify hypotheses. For example,

$$H_1 : \quad \mu_1 > \mu_2 > \mu_3$$

$$JASP : \quad group1 > group2 > group3$$

- Press Ctrl+Enter. Inspect and interpret output.

More in (demo and) lab.

◀ ◻ ▶ ◀ ◻ ▶ ◀ ≡ ▶ ◀ ≡ ▶ ≡ ▶ ↺ 🔍 ↻

GORIC in JASP: Lucas

- Open JASP and 'Data_Lucas_JASP.txt'. Check measurement levels.
- Go to ANOVA and denote variables of interest.
- Go to Order Restricted Hypotheses tab.
- Specify hypotheses. For example,

$$H_1 : \quad \mu_5 = \mu_3 > \{\mu_1, \mu_4\} > \mu_2$$

$$\begin{aligned} \text{JASP :} \quad & \text{group5} = \text{group3} > \text{group1} > \text{group2} \\ & \text{group3} > \text{group4} > \text{group2} \end{aligned}$$

- Press Ctrl+Enter. Inspect and interpret output.

GORIC and GORICA in JASP

	GORIC R	GORIC JASP	GORICA R	GORICA JASP
ANOVA	✓	✓	✓	
RM-ANOVA			✓	✓
Regression	✓	Not yet	✓	
SEM		Future?	✓	Future?
Other model		Future?	✓	Future?

GORIC and GORICA in JASP

GORIC or GORICA in the following statistical models (as part of that model)

The screenshot displays the JASP software interface with three statistical models: ANOVA, ANCOVA, and Repeated Measures ANOVA. In each model's left sidebar, the 'Order Restricted Hypotheses' option is highlighted with a yellow circle. The ANOVA model is labeled 'GORIC' in blue text. The ANCOVA model is labeled 'GORIC' in blue text. The Repeated Measures ANOVA model is labeled 'GORICA' in blue text. The ANOVA model's main panel shows a table with columns: Cases, Sum of Squares, df, Mean Square, F, and p. The ANCOVA model's main panel shows a table with columns: Cases, Sum of Squares, df, Mean Square, F, and p. The Repeated Measures ANOVA model's main panel shows a table with columns: Cases, Sum of Squares, df, Mean Square, F, and p. The right sidebar of the Repeated Measures ANOVA model shows the 'Order Restricted Hypotheses' option highlighted with a red circle. The top right corner of the JASP window shows the 'Display' section with options for Descriptive statistics, Estimates of effect size, and Model.

Table of Contents

Example Null hypothesis (H_0) vs Informative hypothesis (H_m)

Generalized Order-Restricted Information Criterion (GORIC)

GORIC weights

Examples with R code

Failsafe hypotheses: Unconstrained & Complement

GORICA (comparable to b_{ain})

GORIC(A) in JASP

End & Extra

The End

Thanks for listening!

Are there any questions?

Websites

www.uu.nl/staff/RMKuiper/Software

[www.uu.nl/staff/RMKuiper/Websites%20%2F%20Shiny%20apps
informative-hypotheses.sites.uu.nl/software/goric/](http://www.uu.nl/staff/RMKuiper/Websites%20%2F%20Shiny%20apps%20informative-hypotheses.sites.uu.nl/software/goric/)

E-mail

r.m.kuiper@uu.nl

Promo: Publish in special issue

Applying GORIC(A)?

If you collected data or will collect,
if you have one or more a-priori hypotheses,
then you can apply the GORIC(A).

FYI: Special issue

Journal (mdpi): Mathematics (ISSN 2227-7390)

Special Issue:

Evaluation of Theory-Driven Hypotheses: No Hypothesis, No G(L)ORIC

Guest Editor: Rebecca M. Kuiper

https://www.mdpi.com/journal/mathematics/special_issues/97C26430R3

Contact: r.m.kuiper@uu.nl

Please contact me (r.m.kuiper@uu.nl), if you want to explore the possibilities (and to obtain discount).

What's next

Depending on time and wishes:

- Some notes
- Demo in R
- Demo in JASP
- Evidence synthesis / Support aggregation

We end with:

- Lab
- Discussion

Notes

Note on comparable estimates

Until now: comparing means.

Continuous predictors

If compare relative strength/importance of parameters (e.g., $\beta_1 > \beta_2$), then make sure comparable:
e.g., standardize continuous predictors.

Multiple outcomes

If compare parameters across outcomes,
then (also) standardize outcomes.

Note on BF as an IC

The fit in GORICA refers to the maximum log likelihood.

The fit in BF refers to the maximum likelihood.

Notably, BF can be written as an IC (and vice versa):

$$\begin{aligned} -2 \log BF_{1u} &= -2 \log \frac{f_1}{c_1} \\ &= -2 \log f_1 + 2 \log c_1 \\ &= -2 \log \text{fit } H_1 + 2 \log \text{complexity } H_1. \end{aligned}$$

Note: complexity value in BF depends on prior.

Note on GORIC weights vs BF and PMPs

$w_m/w_{m'} = \text{relative weight} \sim \text{Bayes factor } (BF_{mm'})$.

$w_m \sim \text{posterior model probability (PMP)}$.

$1 - w_m = \text{conditional error probability}$.

Note: w_m depends on set of hypotheses.

Note on conditional error probability

using PMPS and GORIC(A) weights

What if we compare many hypotheses?

H_m	weights
H_1 : Sex Match	.013
H_2 : Gender Role Match	.270
H_3 : Sex Mismatch	.003
H_4 : Gender Role Mismatch	.000
H_5 : Lets try this one too	.180
...	...
H_{12} : Don't miss something	.040
H_u :	.047

The conditional error probability for H_2 becomes $1 - .27 = .77$ (was .19).

namely, included additional hypotheses may obtain some support.
Hence, the conditional error probabilities becomes larger.

Note on conditional error probability

using PMPS and GORIC(A) weights

Hypotheses that overlap can also share support (like H_u does).

My advise:

Only use error probabilities if one hypothesis versus its complement.
(Or when you are sure that there is no overlap in hypotheses)

H_m	weights
H_2 : Gender Role Match	.82
H_c : complement H_2	.18

Extra material

- GORICA on SEM, using R scripts from https://github.com/rebeccakuiper/GORICA_in_SEM.
This is material that belongs to the 'GORICA on SEM' article on <https://www.tandfonline.com/doi/full/10.1080/10705511.2020.1836967>.
- GORICA on cross-lagged panel model (CLPM), using R scripts from https://github.com/rebeccakuiper/GORICA_in_CLPM.

Note: On github site, go to Code (green button) and download zip.

Simulation study

Properties of Methods: Simulation study

1. All methods (e.g., (G)ORIC and BMS) programmed in Fortran.
2. Generate data sets with know structure (conditions).
3. Apply methods to all data sets (within one condition).
4. Summarize results (per condition):
proportion of (correct) chosen hypotheses.

Reference:

Kuiper, R.M., Nederhof, T., and Klugkist, I. (2015). Properties of hypothesis testing techniques and (Bayesian) model selection for exploration-based and theory-based (order-restricted) hypotheses. *British Journal of Mathematical and Statistical Psychology*, 68(2), 220 – 245.

Population Values in Simulated Data Sets when $k = 3$

Values of Population Means (μ_i) and Standard Deviations (σ_i) Used to Simulate 1000 Data Sets

$k = 3$	Case Number ('ES')	μ_1	μ_2	μ_3	ES (Cohen)
	0	0.00	0.00	0.00	
	1	-0.10	0.00	0.10	.1
	2	-0.20	0.00	0.20	.2
	3	-0.30	0.00	0.30	.3
	4	-0.40	0.00	0.40	.4
	5	-0.50	0.00	0.50	.5
		σ_1	σ_2	σ_3	
	0 - 5	1.00	1.00	1.00	

NB in Case 0 $H_{30C} : \mu_1 = \mu_2 = \mu_3$ is true
 in Cases 1 to 5 $H_{31C} : \mu_1 < \mu_2 < \mu_3$ is true.

Results Exploration vs Confirmatory ($k = 3$ and $n = 50$)

$H_{30E} : \mu_1 = \mu_2 = \mu_3$	$H_{30C} : \mu_1 = \mu_2 = \mu_3$
$H_{31E} : \mu_1 = \mu_2, \mu_3$	$H_{31C} : \mu_1 < \mu_2 < \mu_3$
$H_{32E} : \mu_1, \mu_2 = \mu_3$	$H_{32C} : \mu_1 = \mu_2 < \mu_3$
$H_{33E} : \mu_1 = \mu_3, \mu_2$	$H_{33C} : \mu_1 < \mu_2 > \mu_3$
$H_{34E} : \mu_1, \mu_2, \mu_3$	$H_{34C} : \mu_1, \mu_2, \mu_3$

'ES'	Method	H_{30E}	H_{31E}	H_{32E}	H_{33E}	H_{34E}
0	PCIC: AIC	0.658	0.113	0.115	0.112	0.002
2	PCIC: AIC	0.201	0.356	0.358	0.028	0.057
5	PCIC: AIC	0.000	0.135	0.136	0.000	0.729

'ES'	Method	H_{30C}	H_{31C}	H_{32C}	H_{33C}	H_{34C}
0	ORIC	0.662	0.065	0.119	0.120	0.034
2	ORIC	0.148	0.506	0.311	0.029	0.006
5	ORIC	0.000	0.955	0.044	0.001	0.000

Results Confirmation ($k = 3$ and $n = 50$)

$H_{30C} : \mu_1 = \mu_2 = \mu_3$

$H_{31C} : \mu_1 < \mu_2 < \mu_3$

$H_{32C} : \mu_1 = \mu_2 < \mu_3$

$H_{33C} : \mu_1 < \mu_2 > \mu_3$

$H_{34C} : \mu_1, \mu_2, \mu_3$

'ES'	Method	'Prior'	H_{30C}	H_{31C}	H_{32C}	H_{33C}	H_{34C}
0	ORIC		0.662	0.065	0.119	0.120	0.034
0	BMS	3	0.865	0.012	0.075	0.036	0.012
0	BMS	2	0.774	0.022	0.124	0.058	0.022
0	BMS	1	0.656	0.047	0.147	0.120	0.033
2	ORIC		0.148	0.506	0.311	0.029	0.006
2	BMS	3	0.299	0.299	0.371	0.031	0.001
2	BMS	2	0.252	0.328	0.381	0.040	0.000
2	BMS	1	0.125	0.494	0.321	0.063	0.000
5	ORIC		0.000	0.955	0.044	0.001	0.000
5	BMS	3	0.000	0.887	0.112	0.001	0.000
5	BMS	2	0.001	0.890	0.107	0.002	0.000
5	BMS	1	0.000	0.928	0.070	0.002	0.000

Results Confirmation ($k = 3$ and $n = 50$)

$H_{30C} : \mu_1 = \mu_2 = \mu_3$

$H_{31C} : \mu_1 < \mu_2 < \mu_3$

$H_{32C} : \mu_1 = \mu_2 < \mu_3$

$H_{33C} : \mu_1 < \mu_2 > \mu_3$

$H_{34C} : \mu_1, \mu_2, \mu_3$

1 – Type I error (here):

not preferring the true null hypothesis,
given the set of hypotheses (!).

'ES'	Method	'Prior'	H_{30C}	H_{31C}	H_{32C}	H_{33C}	H_{34C}
0	ORIC		0.662	0.065	0.119	0.120	0.034
0	BMS	3	0.865	0.012	0.075	0.036	0.012
0	BMS	2	0.774	0.022	0.124	0.058	0.022
0	BMS	1	0.656	0.047	0.147	0.120	0.033
2	ORIC		0.148	0.506	0.311	0.029	0.006
2	BMS	3	0.299	0.299	0.371	0.031	0.001
2	BMS	2	0.252	0.328	0.381	0.040	0.000
2	BMS	1	0.125	0.494	0.321	0.063	0.000
5	ORIC		0.000	0.955	0.044	0.001	0.000
5	BMS	3	0.000	0.887	0.112	0.001	0.000
5	BMS	2	0.001	0.890	0.107	0.002	0.000
5	BMS	1	0.000	0.928	0.070	0.002	0.000

Results Confirmation ($k = 3$ and $n = 50$)

$H_{30C} : \mu_1 = \mu_2 = \mu_3$

$H_{31C} : \mu_1 < \mu_2 < \mu_3$

$H_{32C} : \mu_1 = \mu_2 < \mu_3$

$H_{33C} : \mu_1 < \mu_2 > \mu_3$

$H_{34C} : \mu_1, \mu_2, \mu_3$

1 – Type II error (here):
not preferring the true hypothesis,
given the set of hypotheses (!).

'ES'	Method	'Prior'	H_{30C}	H_{31C}	H_{32C}	H_{33C}	H_{34C}
0	ORIC		0.662	0.065	0.119	0.120	0.034
0	BMS	3	0.865	0.012	0.075	0.036	0.012
0	BMS	2	0.774	0.022	0.124	0.058	0.022
0	BMS	1	0.656	0.047	0.147	0.120	0.033
2	ORIC		0.148	0.506	0.311	0.029	0.006
2	BMS	3	0.299	0.299	0.371	0.031	0.001
2	BMS	2	0.252	0.328	0.381	0.040	0.000
2	BMS	1	0.125	0.494	0.321	0.063	0.000
5	ORIC		0.000	0.955	0.044	0.001	0.000
5	BMS	3	0.000	0.887	0.112	0.001	0.000
5	BMS	2	0.001	0.890	0.107	0.002	0.000
5	BMS	1	0.000	0.928	0.070	0.002	0.000

Extra: Prior choice for BMS

Prior

$$P(\boldsymbol{\mu}, \sigma^2) = P(\mu_1) \times \dots \times P(\mu_k) \times P(\sigma^2),$$

where $P(\mu_i) = P(\mu) \forall i = 1, \dots, k$.

$P(\mu)$ is a data-based normal distribution, with parameters β_0 and τ_0^2 , and $P(\sigma^2)$ an inverse chi-squared distribution (see Klugkist et al., 2005)

β_0 and τ_0^2 data-based hyperparameters

For each μ_i ($i = 1, \dots, k$) a credibility interval is computed:

estimated group mean \pm **PV** \times *the group standard error*.

The credibility interval for μ_i is represented by " $[LB_i, UB_i]$ ".

Lowest lower bound: LB_{min} .

Highest upper bound: UB_{max} .

Then, $\beta_0 = \frac{LB_{min} + UB_{max}}{2}$ and $\tau_0^2 = [\frac{UB_{max} - LB_{min}}{2}]^2$.

Three types of prior vagueness: BMS PV 1, BMS PV 2 and BMS PV 3