

Table of Contents

Example Null hypothesis (H_0) vs Informative hypothesis (H_m)

Generalized Order-Restricted Information Criterion (GORIC)

GORIC weights

Examples with R code

Failsafe/Safeguard hypotheses: Unconstrained & Complement

GORICA

GORIC(A) in JASP

End & Extra

ANOVA Example: Comparisons of 3 Means

Example Palmer and Gough (P&G)

Palmer and Gough (2007) examined the difference in the attribution of importance of defective education as an explanation for criminal behavior

between three types of “offenders”:

 $(n_i = 20, 20, 31)$

ANOVA Example: Comparisons of 3 Means

Example Palmer and Gough (P&G)

Palmer and Gough (2007) examined the difference in the attribution of importance of defective education as an explanation for criminal behavior between three types of “offenders”:
(1) non-offenders, (2) property offenders, and (3) person offenders.

ANOVA

$$y_j = \mu_1 d_{1j} + \mu_2 d_{2j} + \mu_3 d_{3j} + \epsilon_j,$$

where μ_i ($i = 1, 2, 3$) is the mean in group i ,
 d_{ij} is a dummy variable denoting group membership,
 ϵ_j is an error term and $\epsilon_j \sim N(0, \sigma^2)$.

$(n_i = 20, 20, 31)$

Hypotheses of interest

Example Palmer and Gough (P&G)

Palmer and Gough (2007) expect that

$$H_1: \mu_1 > \mu_2 > \mu_3.$$

Hypotheses of interest

Example Palmer and Gough (P&G)

Palmer and Gough (2007) expect that

- non-offenders (1) attribute more importance to defective education for explaining crime than the other two offenders (2 & 3):
i.e., $\mu_1 > \mu_2$ and $\mu_1 > \mu_3$,
- property offenders (2) attribute more importance to defective education than person offenders (3):
i.e., $\mu_2 > \mu_3$.

$$H_1: \mu_1 > \mu_2 > \mu_3.$$

Hypotheses of interest

Example Palmer and Gough (P&G)

Palmer and Gough (2007) expect that

- non-offenders (1) attribute more importance to defective education for explaining crime than the other two offenders (2 & 3):
i.e., $\mu_1 > \mu_2$ and $\mu_1 > \mu_3$,
- property offenders (2) attribute more importance to defective education than person offenders (3):
i.e., $\mu_2 > \mu_3$.

$$H_1: \mu_1 > \mu_2 > \mu_3.$$

Hypotheses of interest

Example Palmer and Gough (P&G)

Palmer and Gough (2007) expect that

- non-offenders (1) attribute more importance to defective education for explaining crime than the other two offenders (2 & 3):
i.e., $\mu_1 > \mu_2$ and $\mu_1 > \mu_3$,
- property offenders (2) attribute more importance to defective education than person offenders (3):
i.e., $\mu_2 > \mu_3$.

This leads to the theory-based hypothesis:

$$H_1 : \quad \mu_1 > \mu_2 > \mu_3.$$

Note that ‘<’ denotes “smaller than” and ‘>’ denotes “larger than”.

Conclusions w.r.t. hypothesis of interest $H_1 : \mu_1 > \mu_2 > \mu_3$

Example 3 means

With post-hoc test (e.g., Bonferroni) and on basis of the sample means, we could say a bit more.

But, what if

- $\mu_1 \neq \mu_2, \mu_1 \neq \mu_3, \mu_2 = \mu_3,$
- $\bar{y}_1 > \bar{y}_2,$

which implies $\mu_1 > \mu_2 = \mu_3$.

But also not interested in this.

Furthermore, inconsistent results are possible (like in P&G example):

$$\mu_1 = \mu_2, \mu_2 = \mu_3, \text{ and } \mu_1 \neq \mu_3,$$

which is logically impossible, since the first two imply $\mu_1 = \mu_3$.

Using planned contrasts would overcome this, but still = restrictions then.

NHST

Example Palmer and Gough

```
PandG_data <- read.table("Data_PalmerAndGough.txt",
                          header=TRUE)
PandG_data$group <- factor(PandG_data$group)
pairwise.t.test(PandG_data$Importance, PandG_data$group,
                p.adj = 'bonferroni')

##
## Pairwise comparisons using t tests with pooled SD
##
## data: PandG_data$Importance and PandG_data$group
##
##      1      2
## 2 0.191 -
## 3 0.011 1.000
##
## P value adjustment method: bonferroni
```


Exploration and Confirmation

Exploration (like in post hoc tests and possibly AIC)

Evaluate all possible pairs/subsets of means whether significant different (“,”) or not (“=”).

For example, when $k = 3$:

$$H_{0E} : \mu_1 = \mu_2 = \mu_3$$

$$H_{1E} : \mu_1 = \mu_2, \mu_3$$

$$H_{2E} : \mu_1, \mu_2 = \mu_3$$

$$H_{3E} : \mu_1 = \mu_3, \mu_2$$

$$H_a : \mu_1, \mu_2, \mu_3$$

When $k = 5$, there are even 52 hypotheses.

Confirmation

Limited set: Evaluate only prespecified hypotheses including order restrictions ($<$, $>$, but also $=$).

Exploration and Confirmation

Exploration (like in post hoc tests and possibly AIC)

Evaluate all possible pairs/subsets of means whether significant different (“,”) or not (“=”).

For example, when $k = 3$:

$$H_{0E} : \mu_1 = \mu_2 = \mu_3$$

$$H_{1E} : \mu_1 = \mu_2, \mu_3$$

$$H_{2E} : \mu_1, \mu_2 = \mu_3$$

$$H_{3E} : \mu_1 = \mu_3, \mu_2$$

$$H_a : \mu_1, \mu_2, \mu_3$$

When $k = 5$, there are even 52 hypotheses.

Confirmation

Limited set: Evaluate only prespecified hypotheses including order restrictions ($<$, $>$, but also $=$).

Confirmatory methods

Most researchers are able to specify “order-restricted” / “informative” / “theory-based” hypotheses, like $H_1 : \mu_1 > \mu_2 > \mu_3$.
Use prior knowledge and/or expertise in hypothesis.

Note: 'model' refers to hypothesis.

Confirmatory methods

Most researchers are able to specify “order-restricted” / “informative” / “theory-based” hypotheses, like $H_1 : \mu_1 > \mu_2 > \mu_3$.
Use prior knowledge and/or expertise in hypothesis.

Methods to evaluate theory-based hypotheses

- Hypothesis testing: F_{bar} ($\bar{F}$) test
(renders p-value and can test only one theory-based hypothesis)
- Confirmatory model selection using information criteria:
GORIC and GORICA
- (Confirmatory) Bayesian model selection (BMS)

Note: 'model' refers to hypothesis.

GORIC

Example Palmer and Gough, that is, 3 means

$$H_0 : \mu_1 = \mu_2 = \mu_3,$$

$$H_1 : \mu_1 > \mu_2 > \mu_3,$$


$$H_u : \mu_1, \mu_2, \mu_3.$$

GORIC – will be explained later


Model	Fit	Complexity	GORIC	GORIC weights
H_0	-196.36	2.00	396.71	0.02
H_1	-191.89	2.81	389.41	0.75
H_u	-191.89	4.00	391.79	0.23

Confirmatory methods (e.g., GORIC) have more “power” than their exploratory counterparts (e.g., AIC; cf. one- vs two-sided testing).

Confirmation more power: 1 data set. GORIC values for 3 groups, effect size ES , and $n = 10$ observations per group



Confirmation more power: 1 data set. GORIC values for 3 groups, effect size ES , and $n = 100$ observations per group



ANOVA Example: Comparisons of 3 Means

Simple, hypothetical example

Examine the difference in happiness

between three types of “treatments”:

(A) treatment A, (B) treatment B, and (C) no treatment / control.

Possible (sets of) theory-based hypothesis/-es:

$$H_1 : \quad \mu_A > \mu_B > \mu_C.$$

$$H_1 : \quad \mu_A > \mu_B > \mu_C,$$

$$H_2 : \quad \mu_B > \mu_A > \mu_C.$$

$$H_1 : \quad \{\mu_A, \mu_B\} > \mu_C.$$

$$H_1 : \quad \mu_A - \mu_B > 0.1 \quad \& \quad \mu_B - \mu_C > 0.2.$$

Your hypothesis of interest

If/When you have your own data

- What is your research question?
- What is your theory / expectation?
- What is your statistical hypothesis?
- Is there a competing statistical hypothesis?

Notably, think outside of the H_0 -box...

Table of Contents

Generalized Order-Restricted Information Criterion (GORIC)

GORIC weights

Examples with R code

GORICA


GORIC(A) in JASP

End & Extra


Intermezzo: Balance Fit and Complexity (1/6)

Example 5 means

Data for 5 groups – Oversimplified representation




Data for 5 groups – Better representation (scatter of data points)



Intermezzo: Balance Fit and Complexity (2/6)

Example 5 means




$$\mu_1 = \mu_2 = \mu_3 = \mu_4 = \mu_5 = \mu$$

parameters: 1 mean parameter (i.e., μ)
Fit: Bad fit

Intermezzo: Balance Fit and Complexity (3/6)

Example 5 means



$$\mu_1 = \mu_2 = \mu_3 = \mu_{\text{low}}$$

$$\mu_4 = \mu_5 = \mu_{\text{high}}$$

parameters:


2 mean parameters (i.e., μ_{low} and μ_{high})

Fit:

Better than with one parameter, even good fit.

Intermezzo: Balance Fit and Complexity (4/6)

Example 5 means




parameters: 5 mean parameter (complex)

Fit: Better than with two parameters, even best possible fit.

Thus: Best fit (= highest likelihood), but also most complex (= highest penalty).

Intermezzo: Balance Fit and Complexity (5/6)

Example 5 means



$$\mu_1 = \mu_2 = \mu_3 = \mu_4 = \mu_5 = \mu$$

$$\mu_1 = \mu_2 = \mu_3 = \mu_{\text{low}} \text{ \& }$$


$$\mu_4 = \mu_5 = \mu_{\text{high}}$$

$$\mu_1, \mu_2, \mu_3, \mu_4, \mu_5$$

In this example: **Two means** may be best trade-off between fit & complexity.

Intermezzo: Balance Fit and Complexity (6b/6)

Order-restrictions: From “Example 5 means” to ‘Example 2 means’





In theory-based hypotheses, we also incorporate **order-restrictions** (e.g., $\mu_1 < \mu_2$). Then, helpful to look at likelihood using **contour plots** (not scatter plot, as done here).

For ease, I will next use **2 means** (instead of 5).

Intermezzo: Contour plot (1/2)

Example 2 means





Intermezzo: Contour plot (2/2)



Example 2 means

Multivariate Gaussian (Normal) examples



$$\mu = \begin{bmatrix} 0 \\ 0 \end{bmatrix} \quad \Sigma = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$$



$$\mu = \begin{bmatrix} 0 \\ 0 \end{bmatrix} \quad \Sigma = \begin{bmatrix} 1 & 0.5 \\ 0.5 & 1 \end{bmatrix}$$




$$\mu = \begin{bmatrix} 0 \\ 0 \end{bmatrix} \quad \Sigma = \begin{bmatrix} 1 & 0.8 \\ 0.8 & 1 \end{bmatrix}$$



Andreas

Note: if in agreement, or-mle = mle

Example 2 means with or-mle = mle



mle in H_3 : highest likelihood is in allowable (= green) space.
Hence, or-mle = mle.

Idea fit

Example Palmer and Gough, that is, 3 means

$$H_0 : \mu_1 = \mu_2 = \mu_3,$$

$$H_1 : \mu_1 > \mu_2 > \mu_3,$$

$$H_2 : \mu_1 > \mu_2 < \mu_3,$$

$$H_3 : \mu_1 < \mu_2 < \mu_3,$$

$$H_u : \mu_1, \mu_2, \mu_3.$$


```
## group1 group2 group3
## 11.95 9.75 8.77
```

GORIC

Model	Fit	Complexity	GORIC	GORIC weights
H_0	-196.36	2.00	396.71	0.01
H_1	-191.89	2.81	389.41	0.56
H_2	-192.34	3.19	391.05	0.25
H_3	-196.36	2.81	398.34	0.01
H_u	-191.89	4.00	391.79	0.17

Idea complexity: penalty (PT)

Example 2 means




Note: sampling from distribution with $\mu_1 = \mu_2 = 0$.

Complexity H_2 is $PT_2 = 1 + 0.5 \times 1 + 0.5 \times 2 = 2.50$.

PT = number of expected distinct/free parameters.

Idea complexity: penalty (PT)

Example 2 means: Contour distribution




Sampling from distribution with $\mu_1 = \mu_2 = 0$. (Note: not likelihood!)

Contour denotes distribution for a variance value.

Choice variance does not matter, see next slide.

Idea complexity: penalty (PT)


Example 2 means: PT value independent from choice variance



For both variance values, that is, for both contours:
 Complexity H_2 is $PT_2 = 1 + 0.5 \times 1 + 0.5 \times 2 = 2.50$.
 PT = number of expected distinct/free parameters.

Extra: Levels and level probabilities

Example 2 means



Level (l): number of distinct/free parameters.

Level probability (LP_l): probability that there are l levels.

$$PT_4 = 1 + 0.25 \times 0 + (0.25 + 0.25) \times 1 + 0.25 \times 2 = 2.00.$$

Idea complexity

Example Palmer and Gough, that is, 3 means

$$H_0 : \mu_1 = \mu_2 = \mu_3,$$

$$H_1 : \mu_1 > \mu_2 > \mu_3,$$

$$H_2 : \mu_1 > \mu_2 < \mu_3,$$

$$H_3 : \mu_1 < \mu_2 < \mu_3,$$

$$H_u : \mu_1, \mu_2, \mu_3.$$

GORIC

Model	Fit	Complexity	GORIC	GORIC weights
H_0	-196.36	2.00	396.71	0.01
H_1	-191.89	2.81	389.41	0.56
H_2	-192.34	3.19	391.05	0.25
H_3	-196.36	2.81	398.34	0.01
H_u	-191.89	4.00	391.79	0.17

GORIC: Lowest value is best

GORIC is like AIC expected distance from the truth (KL-distance).
One wants smallest distance.

Or: One wants the smallest 'misfit + complexity'.

Note: $\text{GORIC} = -2 \text{ fit} + 2 \text{ complexity} = 2 \text{ misfit} + 2 \text{ complexity}$

Hence, smallest value is best.

GORIC

Example Palmer and Gough

- $H_0 :$ $\mu_1 = \mu_2 = \mu_3$ (include only when of interest)
 $H_1 :$ $\mu_1 > \mu_2 > \mu_3$
 $H_2 :$ $\mu_2 > \mu_1 > \mu_3$ (if competing theory/hypothesis)
 $H_u :$ μ_1, μ_2, μ_3 .

GORIC

Model	Fit	Complexity	GORIC
H_0	-196.36	2.00	396.71
H_1	-191.89	2.81	389.41
H_2	-193.70	2.81	393.03
H_u	-191.89	4.00	391.79

Hands-on/Demo (1a): GORIC

- Go to <https://github.com/rebeccakuiper/Tutorials>:
 - Click on green button called Code.
 - Download zip (last option in list).
 - Unzip it on your machine (that folder is now your working dir.).
- Start Rstudio. Optional: make project.
- Open 'Tutorial_GORIC_restriktor_ANOVA.html', 'PalmerAndGough_and_Lucas.R', and/or '**Hands-on_1_GORIC_Unc_ANOVA_restriktor.R**' ('Hands-on files').
- Install packages and load them.
- Read and inspect data.
Use **Data_PalmerAndGough.txt** and/or Data_Lucas.txt.
- Run model (`lm()`).
- Specify hypotheses (make up your own).
Note: Use names used in the model.
- Run `goric()`.
- Inspect and interpret output.
Note: 'GORIC weights' will be explained next.



Table of Contents

Interpretation: GORIC weights

GORIC values

GORIC values cannot be interpreted, only compared:
Smallest is best.

GORIC weights (w_m)

w_m quantifies relative support for H_m versus others in the set.
Values between 0 and 1, and they sum to 1.
The bigger, the better.

GORIC weight ratios ($w_m/w_{m'}$)

$w_m/w_{m'}$ quantifies relative support of H_m vs $H_{m'}$.
Values between 0 and infinity:
* If < 1 , lack of support for H_m (nl. $w_{m'}/w_m$, support for $H_{m'}$).
* If > 1 , support for H_m .

Illustration of the GORIC weights (w_m)

Example Palmer and Gough

$H_0 : \mu_1 = \mu_2 = \mu_3$ (include only when of interest)

$$H_1 : \mu_1 > \mu_2 > \mu_3$$
$$H_2: \mu_2 > \mu_1 > \mu_3 \text{ (if competing theory/hypothesis)}$$
$$H_\mu : \quad \mu_1, \mu_2, \mu_3.$$

GORIC

Model	Fit	Complexity	GORIC	GORIC weights
H_1	-191.89	2.81	389.41	0.68
H_2	-193.70	2.81	393.03	0.11
H_u	-191.89	4.00	391.79	0.21

H_1 is $0.68 / 0.11 \approx 6.11$ times more supported than H_2 .

- If needed: Start Rstudio again (optional: make project) and then also load packages again.
- Read and inspect data.
- Run model (`lm()`).
- Specify hypotheses (make up your own).
Note: Use names used in the model.
- Run `goric()`.
- Inspect and interpret output: Focus on GORIC weights.

GORIC output and interpretation

Palmer & Gough

$$H_1 : \mu_1 > \mu_2 > \mu_3$$

$$H_2 : \mu_2 > \mu_1 > \mu_3$$

$$H_u : \mu_1, \mu_2, \mu_3$$

GORIC

Model	Fit	Complexity	GORIC	GORIC weights
H_1	-191.89	2.81	389.41	0.68
H_2	-193.70	2.81	393.03	0.11
H_u	-191.89	4.00	391.79	0.21

H_1 and H_2 are not weak (nl, better than H_u - will be explained in a bit).

H_1 is $0.68 / 0.11 \approx 6.11$ times more supported than H_2 .

Another illustration: one-way ANOVA

Lucas: 5 groups

Lucas (2003) investigated difference between female and male leadership w.r.t. influence of the leader.

Five experimental groups:

1. a randomly selected male leader
2. a randomly selected female leader
3. male leader selected via task
4. female leader selected via task
5. female leader selected via task + institutionalized female leadership via movie

60/122

Use of H_u

Palmer & Gough

$$H_1 : \mu_1 > \mu_2 > \mu_3,$$

$$H_2 : \mu_2 > \mu_1 > \mu_3$$

$$H_u : \mu_1, \mu_2, \mu_3.$$

GORIC

Model	Fit	Complexity	GORIC	GORIC weights
H_1	-191.89	2.81	389.41	0.68
H_2	-193.70	2.81	393.03	0.11
H_u	-191.89	4.00	391.79	0.21

If at least one informative hypothesis not weak (i.e., $w_m > w_u$ or $w_m/w_u > 1$), then compare informative hypotheses.

Hence: H_u is only a failsafe, not another hypothesis of interest.

Use of H_u

Palmer & Gough

$$H_1 : \mu_1 > \mu_2 > \mu_3,$$

$$H_2 : \mu_2 > \mu_1 > \mu_3$$

$$H_u : \mu_1, \mu_2, \mu_3.$$

GORIC

Model	Fit	Complexity	GORIC	GORIC weights
H_1	-191.89	2.81	389.41	0.68
H_2	-193.70	2.81	393.03	0.11
H_u	-191.89	4.00	391.79	0.21

If at least one informative hypothesis not weak (i.e., $w_m > w_u$ or $w_m/w_u > 1$), then compare informative hypotheses.

Hence: H_u is only a failsafe, not another hypothesis of interest.

Use of H_u

Palmer & Gough

$$H_1 : \mu_1 > \mu_2 > \mu_3,$$

$$H_2 : \mu_2 > \mu_1 > \mu_3$$

$$H_u : \mu_1, \mu_2, \mu_3.$$

GORIC

Model	Fit	Complexity	GORIC	GORIC weights	without Hu
H_1	-191.89	2.81	389.41	0.68	0.86
H_2	-193.70	2.81	393.03	0.11	0.14
H_u	-191.89	4.00	391.79	0.21	NA

H_1 is not weak and is $0.68 / 0.11 = 0.86 / 0.14 \approx 6.11$ times more supported than competing hypothesis H_2 .

H_1 vs H_u

Palmer & Gough

What if only one informative hypothesis:

$$H_1 : \mu_1 > \mu_2 > \mu_3,$$

$$H_u : \mu_1, \mu_2, \mu_3.$$

GORIC

Model	Fit	Complexity	GORIC	GORIC weights
H_1	-191.89	2.81	389.41	0.77
H_u	-191.89	4.00	391.79	0.23

H_1 is $0.77 / 0.23 \approx 3.27$ times more supported than H_u .

BUT: H_u includes H_1 .

So, support for H_u contains support for H_1 .

H_1 vs H_u

Palmer & Gough

What if only one informative hypothesis:

$$H_1 : \mu_1 > \mu_2 > \mu_3,$$

$$H_u : \mu_1, \mu_2, \mu_3.$$

GORIC

Model	Fit	Complexity	GORIC	GORIC weights
H_1	-191.89	2.81	389.41	0.77
H_u	-191.89	4.00	391.79	0.23

H_1 is $0.77 / 0.23 \approx 3.27$ times more supported than H_u .

BUT: H_u includes H_1 .

So, support for H_u contains support for H_1 .

H_1 vs H_u

Palmer & Gough

What if only one informative hypothesis:

$$H_1 : \mu_1 > \mu_2 > \mu_3,$$

$$H_u : \mu_1, \mu_2, \mu_3.$$

GORIC

Model	Fit	Complexity	GORIC	GORIC weights
H_1	-191.89	2.81	389.41	0.77
H_u	-191.89	4.00	391.79	0.23

H_1 is $0.77 / 0.23 \approx 3.27$ times more supported than H_u .


BUT: H_u includes H_1 .

So, support for H_u contains support for H_1 .


Alternative failsafe: Complement of H_m

Based on simulation

vs complement




(a)




(b)

vs unconstrained



(c)



(d)

Note on use of safeguard hypotheses (2/2)

My advise and default in goric function:

- If multiple informative, theory-based hypotheses:
Goal: Comparison of informative, theory-based hypotheses.
Safeguard: Unconstrained.
- If one informative, theory-based hypothesis (H_m):
Goal: Compare H_m with all other possibilities / theories.
Safeguard: Its complement.

Note on use of safeguard hypotheses (2/2)

My advise and default in goric function:

- If multiple informative, theory-based hypotheses:
Goal: Comparison of informative, theory-based hypotheses.
Safeguard: Unconstrained.
- If one informative, theory-based hypothesis (H_m):
Goal: Compare H_m with all other possibilities / theories.
Safeguard: Its complement.

Extra: complement & 'Heq' (1/4)

Palmer & Gough

H_1 and H_c do not really overlap, but they do share a border. The border is defined by boundary hypotheses, that is, all possible hypotheses where ≥ 1 inequalities ('<', '>') are replaced by '='.

$$\begin{aligned} H_{eq} : & \quad \mu_1 = \mu_2, \mu_3, \\ H_1 : & \quad \mu_1 > \mu_2, \mu_3, \\ H_c : & \quad \text{not } H_1, \end{aligned}$$

then one can use 'Heq = TRUE',
as demonstrated on the next slide.

Extra: complement & 'Heq' (3/4)

Palmer & Gough

R output (partly)

```
## restriktor (0.7): generalized order-restricted information criterion:
##
## Results:
##      model      loglik  penalty   goric  loglik.weights  penalty.weights
## 1      Heq   -193.703    3.000  393.406         0.123         0.452
## 2      H1   -191.893    3.500  390.786         0.753         0.274
## 3 complement -193.703    3.500  394.406         0.123         0.274
##      goric.weights  goric.weights_without_heq
## 1             0.188
## 2             0.698
## 3             0.114
##
## Conclusion:
## - The order-restricted hypothesis H1 is the best in the set, as it has the h
## * 'H1' is 6.109 times more supported than 'complement'.
```


- If needed: Start Rstudio again (optional: make project) and then also load packages again.
- Optional: Open 'Tutorial_GORIC_restriktor_ANOVA.html', 'PalmerAndGough_and_Lucas.R', and/or '**Hands-on_2_GORIC_Compl_ANOVA_restriktor.R**'.
- Read and inspect data.
Use Data_PalmerAndGough.txt and/or **Data_Lucas.txt**.
- Run model (lm()).
- Specify hypotheses (make up your own). E.g., for Lucas:

Note: Use names used in the model.

- ◀ ◻ ▶ ◀ ◻ ▶ ◀ ≡ ▶ ◀ ≡ ▶ ≡ ↺ 🔍 ↻ 79/127

Table of Contents

Generalized Order-Restricted Information Criterion (GORIC)

GORIC weights

Examples with R code

GORICA

GORIC(A) in JASP

End & Extra

GORICA

GORIC: Normal linear models

GORIC can easily be applied to normal linear models (e.g., ANOVA models or regression models).

GORIC: Other statistical models

In case of other statistical models (e.g., a SEM model), more cumbersome to calculate maximized order-restricted log likelihood and thus GORIC.

GORICA: All statistical models

Therefore, GORICA: asymptotic expression for GORIC.
Can be used for all types of statistical models.

Reference:

Altınışık, Y., Van Lissa, C. J., Hoijtink, H., Oldehinkel, A. J., and Kuiper, R. M. (2021). Evaluation of inequality constrained hypotheses using a generalization of the AIC. *Psychological Methods*, 26(5), 599–621.

<https://doi.org/10.1037/met0000406>

GORICA

GORIC: Normal linear models

GORIC can easily be applied to normal linear models (e.g., ANOVA models or regression models).

GORIC: Other statistical models

In case of other statistical models (e.g., a SEM model), more cumbersome to calculate maximized order-restricted log likelihood and thus GORIC.

GORICA: All statistical models

Therefore, GORICA: asymptotic expression for GORIC.
Can be used for all types of statistical models.

Reference:

Altınışık, Y., Van Lissa, C. J., Hoijtink, H., Oldehinkel, A. J., and Kuiper, R. M. (2021). Evaluation of inequality constrained hypotheses using a generalization of the AIC. *Psychological Methods*, 26(5), 599–621.

<https://doi.org/10.1037/met0000406>

GORICA

GORIC: Normal linear models

GORIC can easily be applied to normal linear models (e.g., ANOVA models or regression models).

GORIC: Other statistical models

In case of other statistical models (e.g., a SEM model), more cumbersome to calculate maximized order-restricted log likelihood and thus GORIC.

GORICA: All statistical models

Therefore, GORICA: asymptotic expression for GORIC.
Can be used for all types of statistical models.

Reference:

Altınışık, Y., Van Lissa, C. J., Hoijsink, H., Oldehinkel, A. J., and Kuiper, R. M. (2021). Evaluation of inequality constrained hypotheses using a generalization of the AIC. *Psychological Methods*, 26(5), 599–621.

<https://doi.org/10.1037/met0000406>

GORICA

Similarities with GORIC

- Form: $GORICA_m = -2 \text{ fit} + 2 \text{ complexity}$.
- Broad type of restrictions.

Differences compared to GORIC

- Uses asymptotic expression of the likelihood (is a normal):
can therefore be easily applied to all types of statistical models.
Disadvantage: might work less well in case of small samples.
- Does not need data set; mle's and their covariance matrix suffice.
- Can leave out nuisance parameters (i.e., not part of hypotheses).

Note

In case of normal linear models or, otherwise, for not too small samples:
GORICA weights = GORIC weights.

GORICA

Similarities with GORIC

- Form: $GORICA_m = -2 \text{ fit} + 2 \text{ complexity}$.
- Broad type of restrictions.

Differences compared to GORIC

- Uses asymptotic expression of the likelihood (is a normal):
can therefore be easily applied to all types of statistical models.
Disadvantage: might work less well in case of small samples.
- Does not need data set; mle's and their covariance matrix suffice.
- Can leave out nuisance parameters (i.e., not part of hypotheses).

Note

In case of normal linear models or, otherwise, for not too small samples:
GORICA weights = GORIC weights.

GORICA

Similarities with GORIC

- Form: $GORICA_m = -2 \text{ fit} + 2 \text{ complexity}$.
- Broad type of restrictions.

Differences compared to GORIC

- Uses asymptotic expression of the likelihood (is a normal):
can therefore be easily applied to all types of statistical models.
Disadvantage: might work less well in case of small samples.
- Does not need data set; mle's and their covariance matrix suffice.
- Can leave out nuisance parameters (i.e., not part of hypotheses).

Note

In case of normal linear models or, otherwise, for not too small samples:
GORICA weights = GORIC weights.

R code: GORICA

Palmer & Gough

$$H_1 : \mu_1 > \mu_2 > \mu_3,$$

$$H_c : \text{not } H_1.$$

GORICA: type = "gorica" (not default for lm objects)

```

H1 <- 'group1 > group2 > group3'
# vs its complement (default in case of one hypothesis)
#
# GORICA (using goric function in restriktor package)
library(restriktor)
set.seed(123) # Set seed value
gorica.PandG_C <- goric(fit.PandG,
                        hypotheses = list(H1),
                        type = "gorica") # needed if lm object
    
```

GORICA

Palmer & Gough

$$H_1 : \mu_1 > \mu_2 > \mu_3,$$

$$H_c : \text{not } H_1.$$

GORIC

Model	Fit	Complexity	GORICA	GORICA weights
H1	-1.90	1.81	7.42	0.79
complement	-2.34	2.69	10.06	0.21

H_1 is $0.79 / 0.21 \approx 3.74$ times more supported than its complement, that is, any other hypothesis.

Note: GORIC(A) weights are (approx.) the same (ratio may differ a bit).

GORICA

Palmer & Gough

$$H_1 : \mu_1 > \mu_2 > \mu_3,$$

$$H_c : \text{not } H_1.$$

GORIC

Model	Fit	Complexity	GORICA	GORICA weights
H1	-1.90	1.81	7.42	0.79
complement	-2.34	2.69	10.06	0.21

H_1 is $0.79 / 0.21 \approx 3.74$ times more supported than its complement, that is, any other hypothesis.

Note: GORIC(A) weights are (approx.) the same (ratio may differ a bit).

GORIC in JASP: screenshot

Palmer & Gough

Descriptives

T-Tests

ANOVA

Mixed Models

Regression

Frequencies

Factor

Bain

SEM

R (Beta)

Order Restricted Hypotheses

Enter each restriction of one hypothesis on a new line, e.g.,
factorLow == factorMid
factorMid < factorHigh
where 'factor' is the factor (or covariate) name and 'Low'/'Mid'/'High' are the factor level names.
Click on the 'plus' icon to add more hypotheses.
Click the information icon for more examples.

Syntax settings

☐ Include intercept
☐ Show available coefficients

Set for all models

☐ Model summary
☐ Marginal means
☐ Informed hypothesis tests

Model 1

group1 > group2 > group3

Summary for Model 1

Marginal means for Model 1

Informed hypothesis tests for Model 1

Results

ANOVA

ANOVA - Importance

Cases	Sum of Squares	df	Mean Square	F	p
group	123.960	2	61.980	4.554	0.014
Residuals	925.418	68	13.609		

Note. Type III Sum of Squares

Order Restricted Hypotheses

Model Comparison

Model Comparison Table

Model	Log-likelihood	Penalty	GORIC	Weight	Weights ratio
Model 1	-191.893	2.814	389.415	0.789	3.731
Complement	-192.338	3.686	392.048	0.211	1.000

Note. Weights ratios indicate the relative weight for each model against the "complement" model. GORIC = Generalized Order-Restricted Information Criterion (Kuiper, Hoijtink, & Silvapulle, 2011).

89/122

GORIC and GORICA in JASP

GORIC or GORICA in the following statistical models (as part of that model)

The image displays three JASP model configuration windows, each with a list of options on the left and a summary table on the right. The windows are labeled GORIC, GORIC, and GORICA.

GORIC (Top): The left panel shows 'Order Restricted Hypotheses' circled in blue. The right panel shows the 'ANOVA' model summary table.

GORIC (Middle): The left panel shows 'Order Restricted Hypotheses' circled in blue. The right panel shows the 'ANOVA' model summary table.

GORICA (Bottom): The left panel shows 'Order Restricted Hypotheses' circled in blue. The right panel shows the 'Repeated Measures ANOVA' model summary table.

ANCova (Right): The right panel shows the 'ANCova' model summary table, with 'ANCova' circled in blue.

Table of Contents

The End

GORIC(A)

Thanks for listening!

Are there any questions?

Websites

<https://github.com/rebeccakuiper/Tutorials>

www.uu.nl/staff/RMKuiper/Extra1

www.uu.nl/staff/RMKuiper/Extra2

informative-hypotheses.sites.uu.nl/software/goric/

E-mail

r.m.kuiper@uu.nl

Extra material (1/3)

- GORICA on logistic Regression Model
 - Article: doi.org/10.1037/met0000406
- GORICA on SEM
 - Article: www.tandfonline.com/doi/full/10.1080/10705511.2020.1836967.
 - R scripts: github.com/rebeccakuiper/GORICA_in_SEM.
- GORICA on cross-lagged panel model (CLPM)
 - Article: doi.org/10.1111/bjep.12455.
 - R scripts: github.com/rebeccakuiper/GORICA_in_CLPM.
- GORICA on Random-Intercept CLPM (RI-CLPM)
 - Article: Sukpan, C. and Kuiper, R.M. (2023). How to evaluate causal dominance hypotheses in lagged effects models.
 - R scripts: github.com/rebeccakuiper/GORICA%20in%20RI-CLPM.
- GORICA for EffectLiteR
 - R scripts: <https://github.com/rebeccakuiper/Tutorials/tree/main/GORICA%20for%20EffectLiteR>.

Note: On github site, go to Code (green button) and download zip.

Extra material (2/3)

- **GORICA on CTmeta**
 - Article: doi.org/10.1080/10705511.2020.1823228.
 - R scripts: github.com/rebeccakuiper/GORICA_on_CTmeta.
- **GORICA on Meta-analysis**
 - Article: doi.org/10.3390/e24111525.
 - R scripts: github.com/rebeccakuiper/GORICA_on_MetaAn.

Note: On github site, go to Code (green button) and download zip.

Extra material (3/3)

Possibilities multiple studies

- Update GORIC(A) values & weights.
More data collected: (re-)calculate.
- Update hypotheses.
If no expectations (or no support), use exploration and confirmation:
First data set (or a part of it) generates one or more hypotheses.
Other data set (or part) used to determine evidence / support.
Download 'Tutorial_GORIC_restriktor_UpdateHypo.html' and/or
'Hands-on_4_GORIC_UpdateHypo_restriktor.R' from
<https://github.com/rebeccakuiper/Tutorials>.
- Aggregate evidence for hypotheses.
Aggregate the support for theories (diverse designs allowed).
Bear in mind: Meta-analysis aggregates parameter estimates or
effect sizes which need to be comparable (often same designs
required).
Download 'Tutorial_GORIC_restriktor_evSyn.html' and/or
'Hands-on_4_GORIC_evSyn_restriktor.R' from
<https://github.com/rebeccakuiper/Tutorials>.

Note on type of hypotheses: Interaction

ANOVA: possible interaction

What is the relation between “knowledge of numbers after watching Sesame Street for a year” and sex & setting.


Dependent variable: Knowledge of numbers.

Factors: sex (boy, girl) and setting (watching at home, watching at school).

Gr: 1 = boyhome, 2 = boyschool, 3 = girlhome, 4 = girlschool.

Informative Hypotheses

Example 2: ANOVA Interaction Effect



Interaction bottom right:

$$H_m: \quad gr2 - gr1 > gr4 - gr3 \quad \& \quad gr2 > gr1 \quad \& \quad gr2 > gr4$$

Note on type of hypotheses: Magnitude-hypotheses

Hypotheses can regard height of effect size.

In the case of Cohen's d , one could evaluate:

$$H_1 : d < 0,$$

$$H_2 : d > 0,$$

$$H_3 : d > 0.2,$$

$$H_4 : d > 0.5,$$

$$H_5 : d > 0.8.$$

$$H_1 : d < 0,$$

$$H_2 : 0 < d < 0.2,$$

$$H_3 : 0.2 < d < 0.5,$$

$$H_4 : 0.5 < d < 0.8,$$

$$H_5 : d > 0.8.$$

Now, overlapping hypotheses.

Now, range restrictions
(complexity as if equalities).

Or better: One of these versus its complement.

Or: incorporate such cut-off values when inspecting ordering of parameters (say, means; e.g., $\mu_1 - \mu_2 > 0.2 \times s.d.$).

Note on conditional error probability (1/3)

using PMPs and GORIC(A) weights

H_m	weights
H_1 : Sex Match	.04
H_2 : Gender Role Match	.81
H_3 : Sex Mismatch	.01
H_4 : Gender Role Mismatch	.00
H_u :	.14

The conditional error probability for preferred hypotheses H_2 is $1 - .81 = .19$.

Note on conditional error probability (2/3)

using PMPs and GORIC(A) weights

What if we compare many hypotheses?

H_m	weights
H_1 : Sex Match	.013
H_2 : Gender Role Match	.270
H_3 : Sex Mismatch	.003
H_4 : Gender Role Mismatch	.000
H_5 : Lets try this one too	.180
...	...
H_{12} : Don't miss something	.040
H_u :	.047

The conditional error probability for H_2 becomes $1 - .27 = .77$ (was .19).

Namely, included additional hypotheses may obtain some support. Hence, the conditional error probabilities becomes larger.

Note on conditional error probability (3/3)

using PMPS and GORIC(A) weights

What does conditional error probability reflect when hypotheses overlap? Notably, when hypotheses overlap they can also share/divide support (like H_u does with any hypothesis).

My advise:

Only use error probabilities if one hypothesis versus its complement.
(Or when you are sure that there is no overlap in hypotheses;
and they preferably do cover all possibilities)

H_m	weights
H_2 : Gender Role Match	.82
H_c : complement H_2	.18

Simulation study

Properties of Methods: Simulation study

1. All methods (e.g., (G)ORIC and BMS) programmed in Fortran.
2. Generate data sets with know structure (conditions).
3. Apply methods to all data sets (within one condition).
4. Summarize results (per condition):
proportion of (correct) chosen hypotheses.

Reference:

Kuiper, R.M., Nederhof, T., and Klugkist, I. (2015). Properties of hypothesis testing techniques and (Bayesian) model selection for exploration-based and theory-based (order-restricted) hypotheses. *British Journal of Mathematical and Statistical Psychology*, 68(2), 220 – 245.

Results Confirmation ($k = 3$ and $n = 50$)

$$H_{30C} : \mu_1 = \mu_2 = \mu_3$$

$$H_{31C} : \mu_1 < \mu_2 < \mu_3$$

$$H_{32C} : \mu_1 = \mu_2 < \mu_3$$

$$H_{33C} : \mu_1 < \mu_2 > \mu_3$$

$$H_{34C} : \mu_1, \mu_2, \mu_3$$

1 – Type I error (here):

preferring the true null hypothesis,
given the set of hypotheses (!).

'ES'	Method	'Prior'	H_{30C}	H_{31C}	H_{32C}	H_{33C}	H_{34C}
0	ORIC		0.662	0.065	0.119	0.120	0.034
0	BMS	3	0.865	0.012	0.075	0.036	0.012
0	BMS	2	0.774	0.022	0.124	0.058	0.022
0	BMS	1	0.656	0.047	0.147	0.120	0.033
2	ORIC		0.148	0.506	0.311	0.029	0.006
2	BMS	3	0.299	0.299	0.371	0.031	0.001
2	BMS	2	0.252	0.328	0.381	0.040	0.000
2	BMS	1	0.125	0.494	0.321	0.063	0.000
5	ORIC		0.000	0.955	0.044	0.001	0.000
5	BMS	3	0.000	0.887	0.112	0.001	0.000
5	BMS	2	0.001	0.890	0.107	0.002	0.000
5	BMS	1	0.000	0.928	0.070	0.002	0.000

Results Confirmation ($k = 3$ and $n = 50$)

$$H_{30C} : \mu_1 = \mu_2 = \mu_3$$

$$H_{31C} : \mu_1 < \mu_2 < \mu_3$$

$$H_{32C} : \mu_1 = \mu_2 < \mu_3$$

$$H_{33C} : \mu_1 < \mu_2 > \mu_3$$

$$H_{34C} : \mu_1, \mu_2, \mu_3$$

1 – Type II error (here):
preferring the true hypothesis,
given the set of hypotheses (!).

'ES'	Method	'Prior'	H_{30C}	H_{31C}	H_{32C}	H_{33C}	H_{34C}
0	ORIC		0.662	0.065	0.119	0.120	0.034
0	BMS	3	0.865	0.012	0.075	0.036	0.012
0	BMS	2	0.774	0.022	0.124	0.058	0.022
0	BMS	1	0.656	0.047	0.147	0.120	0.033
2	ORIC		0.148	0.506	0.311	0.029	0.006
2	BMS	3	0.299	0.299	0.371	0.031	0.001
2	BMS	2	0.252	0.328	0.381	0.040	0.000
2	BMS	1	0.125	0.494	0.321	0.063	0.000
5	ORIC		0.000	0.955	0.044	0.001	0.000
5	BMS	3	0.000	0.887	0.112	0.001	0.000
5	BMS	2	0.001	0.890	0.107	0.002	0.000
5	BMS	1	0.000	0.928	0.070	0.002	0.000

Extra: Prior choice for BMS

Prior

$$P(\mu, \sigma^2) = P(\mu_1) \times \dots \times P(\mu_k) \times P(\sigma^2),$$

where $P(\mu_i) = P(\mu) \forall i = 1, \dots, k$.

$P(\mu)$ is a data-based normal distribution, with parameters β_0 and τ_0^2 , and $P(\sigma^2)$ an inverse chi-squared distribution (see Klugkist et al., 2005)

β_0 and τ_0^2 data-based hyperparameters

For each μ_i ($i = 1, \dots, k$) a credibility interval is computed:

estimated group mean $\pm$ $PV \times$ *the group standard error*.

The credibility interval for μ_i is represented by " $[LB_i, UB_i]$ ".

Lowest lower bound: LB_{min} .

Highest upper bound: UB_{max} .

Then, $\beta_0 = \frac{LB_{min} + UB_{max}}{2}$ and $\tau_0^2 = \left[\frac{UB_{max} - LB_{min}}{2} \right]^2$.

Three types of prior vagueness: BMS PV 1, BMS PV 2 and BMS PV 3