1. Choose the **hypothesis** with the **highest GORIC(A) weight**, this is the preferred/best hypothesis (called **H_H**).

2. Check: Is there one or more hypotheses (H.) for which the ratio of loglik.weights for H_H vs H. is (approximately) 1 or even smaller than 1?

No, ratio > 1

The ratio of loglik.weights for H_H vs all others is > 1.

i.e., H_H has the highest fit.

There is **support for the preferred Hypothesis H_H**.
See Section 4 of the guidelines.

One can check the ratio of GORIC(A) weights of H_H with other hypotheses and check the case-specific benchmarks (Section 6).

Notes:

In case of overlapping hypotheses, this indicates support for the non-overlapping part. See Section 5.1.1.2 of the guidelines.

In case the failsafe (i.e., the complement or the unconstrained) is the best, one can take on an additional exploratory approach to create one or more new hypotheses for future research.

Yes, ratio = 1 or ≈ 1

The ratio of loglik.weights for H_H vs another is (≈) 1.

i.e., H_H has together with at least one other hypothesis the highest fit (or close to the highest fit).

There is **support for the overlap** in (or boundary of) H_H with the hypotheses with the same/similar fit.
See Section 5.1 of the guidelines.

Interpretating the ratio of GORIC(A) weights is not meaningful.

Note: As a basis for future research, one could specify the overlap (or (the preferred) boundary) and evaluate that against its complement.

Yes, ratio < 1

The ratio of loglik.weights for H_H vs another is < 1.

i.e., at least one other hypothesis has the highest fit.

There is **support for the preferred Hypothesis H_H, but not convincing.** See Section 5.3 of the guidelines.

One can check the ratio of GORIC(A) weights of H_H with other hypotheses; but its support is not convincing.

Notes:

Hypothesis H_H is preferred: It has the best balance between fit and complexity. This is possibly because of a low complexity, since another hypothesis has the highest fit. One can inspect this further with the order-restricted estimates (or-mle's): They show which restrictions are active

The sample size is probably too small (to detect another hypothesis, possibly the fail-safe). Future research (with a higher sample size) can give more insight. One could additionally use an exploratory approach to create one or more new or competing hypotheses for future research.

Yes, both < 1 and (≈) 1

At least one other hypothesis has the highest fit.

&

H_H has (approximately) the same fit as at least one other hypothesis.

There is **support for the overlap** in (or boundary of) H_H with the hypotheses with the same/similar fit, but not convincing.
See Sections 5.1 and 5.3 of the guidelines.

Interpretating the ratio of GORIC(A) weights is not meaningful.

Note: **The sample size is probably too small.** Future research (with a higher sample size) can give more insight in whether the overlap of hypotheses indeed still seems to be the best.

Note: If H_H is true in the population and no other hypothesis contains the truth, its GORIC(A) weight will go to 1 when sample size or effect size increases.
Stated otherwise, in such a case, the GORIC(A) weights reflect the uncertainty of a hypothesis being the best.