

Pre-Survey

Please complete the survey below.

Thank you!

Pre-Session Attitudes


- | | |
|--|---|
| 1) On a scale of 1-10, how confident are you that you can reliably facilitate delivery standard of care resuscitation for an in-hospital cardiac arrest? (meaning, almost every time you lead a code, patients receive the standard of care) 1 is not confident at all, 5 is neutral, 10 is very confident. | ☐ 1
☐ 2
☐ 3
☐ 4
☐ 5
☐ 6
☐ 7
☐ 8
☐ 9
☐ 10 |
| <hr/> | |
| 2) On a scale of 1-10, how confident are you that you can reliably facilitate delivery standard of care resuscitation for rapid response/MET? (meaning, almost every time you lead a rapid response/MET, patients receive the standard of care) 1 is not confident at all, 5 is neutral, 10 is very confident. | ☐ 1
☐ 2
☐ 3
☐ 4
☐ 5
☐ 6
☐ 7
☐ 8
☐ 9
☐ 10 |
| <hr/> | |
| 3) On a scale of 1-10, how nervous are you about the process of leading a code blue team? 1 is not nervous at all, 5 is neutral, 10 is very nervous. | ☐ 1
☐ 2
☐ 3
☐ 4
☐ 5
☐ 6
☐ 7
☐ 8
☐ 9
☐ 10 |
| <hr/> | |
| 4) On a scale of 1-10, how nervous are you about the process of leading a rapid response team? 1 is not nervous at all, 5 is neutral, 10 is very nervous. | ☐ 1
☐ 2
☐ 3
☐ 4
☐ 5
☐ 6
☐ 7
☐ 8
☐ 9
☐ 10 |

Pre-session experience:

- 5) Roughly, how many code blues have you been to as an intern (best guess)? _____
- 6) Roughly, how many rapid responses/METs have you been to as an intern (best guess)? _____
- 7) Have you led either a rapid response or a code blue as an intern? How many? _____
- 8) How many times have you led a simulated/scenario code blue or rapid response in training? _____

Knowledge Assessment

- 9) You are on the code team. You're paged to a code on a patient on the surgical floor. The attending surgeon is present. You...
- ☐ defer management to the primary team
☐ ask if anyone is running the code
☐ begin running the code
☐ Leave



- 10) 2 minute pulse check shows the above image. What is the next step?
- ☐ Cardiovert
☐ Defibrillate
☐ Transcutaneously pace
☐ Resume compressions
- 11) A cardiac arrest occurring in which of the following patients would warrant calling for additional help beyond those that usually respond at the U of U ?
- ☐ a code blue in a 30 year old with myocarditis
☐ a respiratory arrest on the floor that has been paged out as a code blue
☐ a rapid response on a surgical patient.
☐ a rapid response to rehab
- 12) Bringing family in to observe an attempt at resuscitation has been shown to: [select all that apply]
- ☐ increase ROSC rate
☐ decrease ROSC rate
☐ increase PTSD in the family
☐ decrease PTSD in the family
- 13) In cardiac arrest scenarios, what interventions have a known, well-established mortality benefit for in hospital cardiac arrest? (Select all that apply)
- ☐ Minimally interrupted CPR
☐ Sodium bicarb in acidemic patients
☐ LUCAS device compressions for patients undergoing more than 30 minutes of CPR
☐ Defibrillation of Shockable Rhythms
☐ ECMO (E-CPR)
- 14) Which of the following is likely to cause a narrow QRS complex PEA arrest?
- ☐ Hyperkalemia
☐ Amitriptyline Overdose
☐ Pulmonary Embolism
☐ Ostial LAD Obstruction

- 15) A rapid response is called on a patient for agitation. They appear intoxicated, perhaps from methamphetamine use. They are tachycardic and mildly hyperthermic, but vitals are otherwise stable. At the last rapid, 2mg of lorazepam, 2.5 mg of haloperidol, and 25 mg of benadryl were used. This was effective, but took 25 minutes before he was safely calmed, and was short-lived. The nurse has been unable to leave the room all morning. What should you do?
- ☐ 1. Advocate for moving the patient to the ICU for staffing reasons.
 - ☐ 2. Increase the strength of sedative by switching to phenobarbital
 - ☐ 3. Defer management to the primary team
 - ☐ 4. Place the patient in restraints so that the nurse can go care for their other patients.