

Chronic Respiratory Disease

Association of obstructive sleep apnea with all-cause and cardiovascular mortality: A population-based study

Journal:	<i>Chronic Respiratory Disease</i>
Manuscript ID	CRD-22-0131
Manuscript Type:	Original Paper
Date Submitted by the Author:	03-Aug-2022
Complete List of Authors:	Gao, Jia; Jiangsu Province Geriatric Hospital Shi, Licheng; Jiangsu Province Geriatric Hospital Sun, Min; Jiangsu Province Geriatric Hospital Guo, Hongmei; Jiangsu Province Geriatric Hospital Gu, Jianhua; Jiangsu Province Geriatric Hospital Zhang, Dandan; Jiangsu Province Geriatric Hospital Wang, Wenjun; Jiangsu Province Geriatric Hospital Zhu, Xuanfeng; Jiangsu Province Geriatric Hospital, Liu, Jiannan; Jiangsu Province Geriatric Hospital
Keywords:	obstructive sleep apnea, cardiometabolic diseases, hypertension, diabetes, cardiovascular diseases
Abstract:	<p>Objectives: Obstructive sleep apnea (OSA) is one of the leading respiratory disorders, increasing the risk of cardiometabolic diseases. In the study, we investigated the association between OSA and the risk of cardiometabolic diseases and all-cause and cardiovascular mortality in adults.</p> <p>Methods: We made analysis based on National Health and Nutrition Examination Survey (NHANES) 2005-2008. The diagnosis of OSA was obtained from self-reported interviews. The baseline covariates were compared between participants with and without OSA status. Multivariable logistic regression was performed to explore the association between OSA and cardiometabolic diseases while multivariable Cox regression for all-cause and cardiovascular mortality.</p> <p>Results: OSA status was positively associated with higher risks of cardiometabolic diseases, including hypertension (odds ratio [OR] 1.28, 95% confidence interval [CI] 1.14-1.45; $P < 0.001$), diabetes (OR 1.46, 95%CI 1.22-1.76; $P < 0.001$) and cardiovascular diseases (OR 1.29; 95%CI 1.08-1.54; $P = 0.006$) after adjusting for numerous covariates. However, no associations of OSA with all-cause or cardiovascular mortality were observed.</p> <p>Conclusions: OSA was associated with a higher risk of hypertension, diabetes and cardiovascular diseases, while had no significant association with all-cause and cardiovascular mortality.</p>

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60



Association of obstructive sleep apnea with all-cause and cardiovascular mortality: A population-based study

Gao, Jia MD^{a*}; Shi, Licheng MD^{a*}; Sun, Min MD^{b*}; Guo, Hongmei MD^{b*}; Gu, Jianhua MD^a; Zhang, Dandan MD^a; Wang, Wenjun MD^a; Zhu, Xuanfeng MD^{a#}; Liu, Jiannan MD^{a#}

a. Department of Respiratory Medicine, Geriatric Hospital of Nanjing Medical University; Jiangsu Province Official Hospital, Nanjing, China.

b. Hypertension Research Institute of Geriatric Hospital of Nanjing Medical University; Jiangsu Province Official Hospital, Nanjing, China.

*Contributed equally

#Corresponding to

Xuanfeng Zhu, Department of Respiratory Medicine, Geriatric Hospital of Nanjing Medical University; Jiangsu Province Official Hospital. 65 Jiangsu Road, Nanjing, 210009, China. Email: zhuxuanfeng@snapmail.cc

Jiannan Liu, Department of Respiratory Medicine, Geriatric Hospital of Nanjing Medical University; Jiangsu Province Official Hospital. 65 Jiangsu Road, Nanjing, 210009, China. Email: liujiannan@lista.cc

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

23 **Abstract**

24 **Objectives:** Obstructive sleep apnea (OSA) is one of the leading respiratory disorders, increasing
25 the risk of cardiometabolic diseases. In the study, we investigated the association between OSA
26 and the risk of cardiometabolic diseases and all-cause and cardiovascular mortality in adults.

27 **Methods:** We made analysis based on National Health and Nutrition Examination Survey
28 (NHANES) 2005-2008. The diagnosis of OSA was obtained from self-reported interviews. The
29 baseline covariates were compared between participants with and without OSA status.
30 Multivariable logistic regression was performed to explore the association between OSA and
31 cardiometabolic diseases while multivariable Cox regression for all-cause and cardiovascular
32 mortality.

33 **Results:** OSA status was positively associated with higher risks of cardiometabolic diseases,
34 including hypertension (odds ratio [OR] 1.28, 95% confidence interval [CI] 1.14-1.45; $P < 0.001$),
35 diabetes (OR 1.46, 95%CI 1.22-1.76; $P < 0.001$) and cardiovascular diseases (OR 1.29; 95%CI
36 1.08-1.54; $P = 0.006$) after adjusting for numerous covariates. However, no associations of OSA
37 with all-cause or cardiovascular mortality were observed.

38 **Conclusions:** OSA was associated with a higher risk of hypertension, diabetes and cardiovascular
39 diseases, while had no significant association with all-cause and cardiovascular mortality.

40
41 **Keywords:** obstructive sleep apnea; cardiometabolic diseases; hypertension; diabetes;
42 cardiovascular diseases

1 Introduction

Obstructive sleep apnea (OSA) is one of the sleep-related breathing disorders and repetitive episodes of upper airway obstruction results in reduced airflow, nocturnal hypoxemia and hypercapnia, arousals, sympathetic stimulation, and exaggerated intrathoracic pressure swings[1]. The apnea-hypopnea index (AHI) is the most commonly used metric to determine the diagnosis and severity classification of OSA, as well as the associations between OSA and comorbidities.

Several studies have emphasized that OSA was an independent risk factor for coronary heart disease[2, 3], arrhythmia[4, 5], stroke[6, 7], and hypertension[8]. Metabolic disorders, including diabetes and impaired lipid metabolism, were also associated with OSA[9, 10]. Besides, OSA is a condition with potential for negative feedback in which it worsens conditions that may in turn worsen the OSA (eg, OSA → hypertension → worsened OSA). The underlying mechanisms are thought to include increased sympathetic nervous system activity, oxidative stress, and systemic inflammation. While OSA treatment with positive airway pressure is associated with a reduction in cardiovascular disease risk[11]. Previous studies found that OSA was reported to be associated with all-cause and cardiovascular mortality in patients with chronic obstructive pulmonary disease[12], acute coronary syndrome[13], and chronic kidney disease[14]. However, data on the link between OSA and mortality in general population was limited.

In the present study, we aimed to examine the association between OSA with all-cause and cardiovascular mortality in a nationally representative sample of the US population. It will improve our insights into the prevention and management of OSA.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

67 **2 Methods**

68 **2.1 Study population**

69 We made a secondary analysis based on the National Health and Nutrition Examination Survey
70 (NHANES) 2005-2008, a nationwide multistage-sampling survey. Firstly, participants with
71 missing data on OSA (n = 8310) and cardiometabolic diseases (n =1715) were excluded.
72 Participants with cancer and pregnancy (n = 1279) and unavailable mortality status (n =11) were
73 also excluded because pregnancy and cancer could have cofounding effect the sleep status and
74 mortality respectively. A total of 9076 respondents were analyzed in this study (**Figure 1**). The
75 study was approved by the Review Board of National Center of Health Statistics and the written
76 consent was obtained from the participants.

77 **2.2 Exposure and outcomes**

78 The presence of OSA was determined through questionnaire that was administered by trained
79 interviewers. Subjects who answered yes to the question “Have you ever been told by a doctor or
80 other health professional that you have a sleep disorder?” and reported sleep apnea to “What was
81 the sleep disorder?” were defined as having OSA[15].
82 Cardiovascular diseases consisting of congestive heart failure, coronary heart disease, angina
83 pectoris, heart attack, and stroke were obtained from the self-reports. The questions included “Has
84 a doctor or other health professional ever told you that you have congestive heart failure /coronary
85 heart disease/angina pectoris/heart attack/stroke?”. Hypertension was diagnosed as the previous
86 hypertension, or systolic blood pressure ≥ 140 mmHg, or diastolic blood pressure ≥ 90 mmHg, or
87 taking antihypertensive drugs. Diabetes was diagnosed as previous diabetes or fasting glucose >7
88 mmol/L or HBA1c $>6.5\%$ or use of hypoglycemic drugs.

1
2
3
4 89 Mortality status was determined from the National Death Index by 31 December 2015. The ICD-
5
6 90 10 codes for cardiovascular diseases included I00-I09, I11, I13, I20-I51, I60-I69 and I70-78.
7
8

9 91 **2.3 Survey and measurement**

10
11 92 A standard questionnaire was used to collect demographic information, including age, sex, race,
12
13 93 educational level, poverty income ratio (PIR), body mass index (BMI), drinking and smoking
14
15 94 status, history of diseases (hypertension, diabetes and cardiovascular disease) and history of
16
17 95 medication (antihypertensive drug, hypoglycemic drug and lipid-lowering drug). Smoking was
18
19 96 categorized into current, former and never. Race/ethnicity was divided into four groups: non-
20
21 97 Hispanic white or black, Mexican American and Others. Education level included four categories:
22
23 98 less than or equivalent high school, college or above. Height and weight were measured by
24
25 99 physical examinations, and BMI was calculated as body weight divided height squared. Serum
26
27 100 levels of cholesterol, triglycerides, high density lipoprotein cholesterol (HDL-C) and low density
28
29 101 lipoprotein cholesterol (LDL-C) were measured using by an enzymatic assay. Estimated
30
31 102 glomerular filtration rate (eGFR) was calculated based on CKD-EPI equation.
32
33
34
35
36
37
38
39

40 103 **2.4 Statistical analysis:**

41
42 104 Numeric parameters were expressed as the mean \pm standard deviation and categorical variables
43
44 105 were presented as numbers (percentages). Differences between groups was compared using
45
46 106 analysis of variance for quantitative values and Chi-squared test for qualitative parameters.
47
48 107 Multivariable logistic regression analysis was used to explore the cross-sectional relationship
49
50 108 between OSA with cardiometabolic diseases. Multivariable Cox regression was performed to
51
52 109 explore the prognostic effect of OSA in all-cause and cardiovascular mortality. Model 1 was
53
54 110 unadjusted; Model 2 was adjusted for gender, and age. Model 3 was adjusted for Model 2
55
56
57
58
59
60

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

111 covariates plus race, education, PIR, BMI, drinking, smoking, and activity. Model 4 was adjusted
112 for Model 3 covariates plus medications use and cholesterol, triglycerides, HDL, LDL, and eGFR.
113 All statistical analysis was performed using IBM SPSS 25.0. P value < 0.05 was deemed as
114 statistically significant.
115
116
117
118
119
120
121
122
123
124
125
126
127
128
129
130
131
132

3 Results

9076 individuals were included in our study. Compared to participants without OSA, population with OSA tend to be young, female and non-Hispanic white, and had more percentage of diabetes, hypertension, and cardiovascular disease. Cholesterol, LDL-C and eGFR were lower among participants with OSA (**Table 1**).

To investigate the association between OSA and cardiometabolic diseases (hypertension, diabetes and cardiovascular diseases), we established multiple logistic regression models shown in **Table 2**. OSA was positively associated with higher odds of hypertension (OR = 1.44; 95%CI, 1.29, 1.60; $P < 0.001$), diabetes (OR = 1.86; 95%CI, 1.65, 2.09; $P < 0.001$) and cardiovascular diseases (OR = 1.54; 95%CI, 1.32, 1.80; $P < 0.001$) in Model 1. This association was not altered after adjusting for gender, age, race, education, PIR, BMI, drinking, smoking, activity, hypertension, diabetes, CVD, antihypertensive drugs, hypoglycemic drugs, lipid-lowering drugs, cholesterol, triglycerides, HDL, LDL, and eGFR in Model 4. Furthermore, we investigated the association between OSA with all-cause and cardiovascular mortality. Multivariable Cox regression analysis found that OSA had no significant association with all-cause and cardiovascular mortality in all models (**Table 3**).

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

155

156

157

158

159

160

161

162

163

164

165

166

167

168

169

170

171

172

173

174

175

176

4 Discussion

Our study concluded that OSA was associated with the presence of cardiometabolic diseases, including diabetes, hypertension and cardiovascular diseases, while had no significant relationship with all-cause and cardiovascular mortality.

In according to our results that OSA increased the risk of cardiovascular diseases. Punjabi et al. showed that severe hypopneas were independently associated with cardiovascular diseases[16]. Sarah et al conducted a longitudinal study showing that hypertension was related to OSA during rapid eye movement sleep during community-dwelling men[17]. OSA and diabetes also has been known to be closely related to each other[10].The prevalence of OSA in people with type 2 diabetes mellitus was higher than the general population[18]. The underlying mechanism might be that OSA induced intermittent hypoxia and arousals, which may result in decreased insulin sensitivity, sympathetic excitation, and systemic inflammation that eventually led to diabetes[19-21]. Our study was in consistent with previous results and found OSA was associated with the presence of cardiometabolic diseases. OSA and cardiovascular disease may share multiple common risk factors, including obesity, male sex, and older age. Besides, the mechanisms underlying this association are thought to include increased sympathetic nervous system activity, oxidative stress, and systemic inflammation, with these effects mediated in part by elevated blood pressure and impaired glucose metabolism.

Recent study showed that hypoxemic burden due to sleep apnea was strongly predictive of cardiovascular diseases related mortality[22]. Lavie et al. conducted a large prospective cohort study and concluded that severe OSA increased the risk of mortality in middle-aged male

177 patients[23]. Marin et al. also concluded that those with severe OSA were at higher risk for both
178 fatal and non-fatal cardiovascular events[24]. However, we found OSA had no significant
179 relationship with all-cause and cardiovascular mortality. The difference could be due to the fact
180 that OSA diagnosis was based on patient self-reports rather than objective tests. The underlying
181 reasons need further investigations.

182 Our study has some strengths. Firstly, our association was robust after adjusting so many
183 covariates, which was not included in previous studies. Besides, our sample was relatively large.
184 Some limitations also existed in our study. First, some variables were based on questionnaires.
185 Secondly, OSA was not diagnosed by polysomnography and therefore the prevalence of OSA
186 could be overestimated.

187 **5 Conclusion**

188 In conclusion, we found that OSA was associated with the presence of cardiometabolic diseases,
189 including diabetes, hypertension and cardiovascular diseases. However, it could not predict the
190 mortality in general population.

191

192 **List of abbreviations**

193 OSA, Obstructive sleep apnea;

194 NHANES, National Health and Nutrition Examination Survey;

195 OR, odds ratio;

196 CI, confidence interval;

197 AHI, apnea-hypopnea index;

198 NCHS, National Center of Health Statistics;

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

199 PIR, poverty income ratio;

200 BMI, body mass index;

201 HDL-C, high density lipoprotein cholesterol;

202 LDL-C, low density lipoprotein cholesterol;

203 eGFR, estimated glomerular filtration rate;

204

205 **Declarations**

206 **Ethics approval and consent to participate**

207 The study was approved by the Review Board of National Center of Health Statistics and the

208 written consent was obtained from the participants.

209 **Consent for publication**

210 Not applicable.

211 **Availability of data and materials**

212 All data could be obtained upon request from Xuanfeng Zhu.

213 **Competing interests**

214 The authors have nothing to disclose.

215 **Funding**

216 None.

217 **Acknowledgements**

218 None.

219 **Author contribution**

220 ZXF and LJN designed this study; GJ, SLC and SM performed the statistical analysis; GHM and

GJH wrote the manuscript; ZDD and WWJ prepared the tables and figures.

References

1. Won CHJ, Qin L, Selim B, Yaggi HK. Varying Hypopnea Definitions Affect Obstructive Sleep Apnea Severity Classification and Association With Cardiovascular Disease. *J Clin Sleep Med*. 2018 Dec 15;14(12):1987-94. PMID: 30518445. doi: 10.5664/jcsm.7520.
2. Hla KM, Young T, Hagen EW, Stein JH, Finn LA, Nieto FJ, et al. Coronary heart disease incidence in sleep disordered breathing: the Wisconsin Sleep Cohort Study. *Sleep*. 2015 May 1;38(5):677-84. PMID: 25515104. doi: 10.5665/sleep.4654.
3. Cepeda-Valery B, Acharjee S, Romero-Corral A, Pressman GS, Gami AS. Obstructive sleep apnea and acute coronary syndromes: etiology, risk, and management. *Curr Cardiol Rep*. 2014;16(10):535. PMID: 25135347. doi: 10.1007/s11886-014-0535-y.
4. Selim BJ, Koo BB, Qin L, Jeon S, Won C, Redeker NS, et al. The Association between Nocturnal Cardiac Arrhythmias and Sleep-Disordered Breathing: The DREAM Study. *J Clin Sleep Med*. 2016 Jun 15;12(6):829-37. PMID: 26951420. doi: 10.5664/jcsm.5880.
5. Marti-Almor J, Jimenez-Lopez J, Casteigt B, Conejos J, Valles E, Farre N, et al. Obstructive Sleep Apnea Syndrome as a Trigger of Cardiac Arrhythmias. *Curr Cardiol Rep*. 2021 Feb 20;23(3):20. PMID: 33611699. doi: 10.1007/s11886-021-01445-y.
6. Culebras A, Anwar S. Sleep Apnea Is a Risk Factor for Stroke and Vascular Dementia. *Curr Neurol Neurosci Rep*. 2018 Jun 25;18(8):53. PMID: 29938308. doi: 10.1007/s11910-018-0855-1.
7. Catalan-Serra P, Campos-Rodriguez F, Reyes-Nunez N, Selma-Ferrer MJ, Navarro-Soriano C, Ballester-Canelles M, et al. Increased Incidence of Stroke, but Not Coronary Heart Disease, in Elderly Patients With Sleep Apnea. *Stroke*. 2019 Feb;50(2):491-4. PMID: 30580706. doi: 10.1161/STROKEAHA.118.023353.
8. Van Ryswyk E, Mukherjee S, Chai-Coetzer CL, Vakulin A, McEvoy RD. Sleep Disorders, Including Sleep Apnea and Hypertension. *Am J Hypertens*. 2018 Jul 16;31(8):857-64. PMID: 29788034. doi: 10.1093/ajh/hpy082.
9. Tamura A, Kawano Y, Watanabe T, Kadota J. Relationship between the severity of obstructive sleep apnea and impaired glucose metabolism in patients with obstructive sleep apnea. *Respir Med*. 2008 Oct;102(10):1412-6. PMID: 18606532. doi: 10.1016/j.rmed.2008.04.020.
10. Song SO, He K, Narla RR, Kang HG, Ryu HU, Boyko EJ. Metabolic Consequences of Obstructive Sleep Apnea Especially Pertaining to Diabetes Mellitus and Insulin Sensitivity. *Diabetes Metab J*. 2019 Apr;43(2):144-55. PMID: 30993938. doi: 10.4093/dmj.2018.0256.
11. Fava C, Dorigoni S, Dalle Vedove F, Danese E, Montagnana M, Guidi GC, et al. Effect of CPAP on blood pressure in patients with OSA/hypopnea a systematic review and meta-analysis. *Chest*. 2014 Apr;145(4):762-71. PMID: 24077181. doi: 10.1378/chest.13-1115.
12. Kendzerska T, Leung RS, Aaron SD, Ayas N, Sandoz JS, Gershon AS. Cardiovascular Outcomes and All-Cause Mortality in Patients with Obstructive Sleep Apnea and Chronic Obstructive Pulmonary Disease (Overlap Syndrome). *Ann Am Thorac Soc*. 2019 Jan;16(1):71-81. PMID: 30372124. doi: 10.1513/AnnalsATS.201802-136OC.
13. Mazaki T, Kasai T, Yokoi H, Kuramitsu S, Yamaji K, Morinaga T, et al. Impact of Sleep-Disordered Breathing on Long-Term Outcomes in Patients With Acute Coronary Syndrome Who Have Undergone Primary Percutaneous Coronary Intervention. *J Am Heart Assoc*. 2016 Jun 15;5(6). PMID: 27307401.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

doi: 10.1161/JAHA.116.003270.

14. Tang SC, Lam B, Yao TJ, Leung WS, Chu CM, Ho YW, et al. Sleep apnea is a novel risk predictor of cardiovascular morbidity and death in patients receiving peritoneal dialysis. *Kidney Int.* 2010 Jun;77(11):1031-8. PMID: 20237456. doi: 10.1038/ki.2010.76.

15. Du W, Liu J, Zhou J, Ye D, OuYang Y, Deng Q. Obstructive sleep apnea, COPD, the overlap syndrome, and mortality: results from the 2005-2008 National Health and Nutrition Examination Survey. *Int J Chron Obstruct Pulmon Dis.* 2018;13:665-74. PMID: 29520134. doi: 10.2147/COPD.S148735.

16. Punjabi NM, Newman AB, Young TB, Resnick HE, Sanders MH. Sleep-disordered breathing and cardiovascular disease: an outcome-based definition of hypopneas. *Am J Respir Crit Care Med.* 2008 May 15;177(10):1150-5. PMID: 18276938. doi: 10.1164/rccm.200712-1884OC.

17. Appleton SL, Vakulin A, Martin SA, Lang CJ, Wittert GA, Taylor AW, et al. Hypertension Is Associated With Undiagnosed OSA During Rapid Eye Movement Sleep. *Chest.* 2016 Sep;150(3):495-505. PMID: 27001264. doi: 10.1016/j.chest.2016.03.010.

18. Elmasry A, Lindberg E, Berne C, Janson C, Gislason T, Awad Tageldin M, et al. Sleep-disordered breathing and glucose metabolism in hypertensive men: a population-based study. *J Intern Med.* 2001 Feb;249(2):153-61. PMID: 11240844. doi: 10.1046/j.1365-2796.2001.00787.x.

19. Narkiewicz K, van de Borne PJ, Montano N, Dyken ME, Phillips BG, Somers VK. Contribution of tonic chemoreflex activation to sympathetic activity and blood pressure in patients with obstructive sleep apnea. *Circulation.* 1998 Mar 17;97(10):943-5. PMID: 9529260. doi: 10.1161/01.cir.97.10.943.

20. Deibert DC, DeFronzo RA. Epinephrine-induced insulin resistance in man. *J Clin Invest.* 1980 Mar;65(3):717-21. PMID: 6243677. doi: 10.1172/JCI109718.

21. Lee EJ, Heo W, Kim JY, Kim H, Kang MJ, Kim BR, et al. Alteration of Inflammatory Mediators in the Upper and Lower Airways under Chronic Intermittent Hypoxia: Preliminary Animal Study. *Mediators Inflamm.* 2017;2017:4327237. PMID: 29038619. doi: 10.1155/2017/4327237.

22. Azarbarzin A, Sands SA, Stone KL, Taranto-Montemurro L, Messineo L, Terrill PI, et al. The hypoxic burden of sleep apnoea predicts cardiovascular disease-related mortality: the Osteoporotic Fractures in Men Study and the Sleep Heart Health Study. *Eur Heart J.* 2019 Apr 7;40(14):1149-57. PMID: 30376054. doi: 10.1093/eurheartj/ehy624.

23. Lavie P, Lavie L, Herer P. All-cause mortality in males with sleep apnoea syndrome: declining mortality rates with age. *Eur Respir J.* 2005 Mar;25(3):514-20. PMID: 15738297. doi: 10.1183/09031936.05.00051504.

24. Marin JM, Carrizo SJ, Vicente E, Agusti AG. Long-term cardiovascular outcomes in men with obstructive sleep apnoea-hypopnoea with or without treatment with continuous positive airway pressure: an observational study. *Lancet.* 2005 Mar 19-25;365(9464):1046-53. PMID: 15781100. doi: 10.1016/S0140-6736(05)71141-7.

Table 1. Baseline variables of the study participants according to OSA status.

Variables	OSA (n=4648)	No OSA (n=4428)	P
Male	1915 (43.2)	2725 (58.6)	<0.001
Age, years	56.84 (9.09)	59.99 (6.26)	<0.001
Race (%)			
Non-Hispanic white	2042 (46.1)	2130 (45.8)	0.416
Non-Hispanic black	1009 (22.8)	1028 (22.1)	
Mexican American	857 (19.4)	892 (19.2)	
Others	520 (11.7)	598 (12.9)	
Education (%)			
Less than high school	1293 (29.2)	1398 (30.1)	0.002
High school or equivalent	1018 (23.0)	1188 (25.6)	
College or above	2111 (47.7)	2057 (44.3)	
PIR (%)			
<1	849 (20.8)	780 (17.9)	0.003
1~3	1714 (42.0)	1850 (42.6)	
>3	1521 (37.2)	1716 (39.5)	
BMI, kg/m ²	27.14 (5.73)	30.55 (7.25)	<0.001
Drinking (%)	578 (48.4)	632 (57.0)	<0.001
Smoking (%)			
Current	773 (22.2)	987 (28.8)	<0.001
Past	168 (4.8)	182 (5.3)	
Never	2543 (73.0)	2257 (65.9)	
Activity (%)			
Vigorous	1166 (44.2)	1182 (43.3)	0.52
Moderate	1089 (41.3)	1168 (42.8)	
Inactive	381 (14.5)	378 (13.9)	
Past history (%)			
HBP	658 (16.6)	955 (22.3)	<0.001
DM	497 (11.2)	884 (19.0)	<0.001
CVD	288 (6.5)	450 (9.7)	<0.001
Prior medication (%)			
Antihypertensive drug	986 (85.2)	1536 (86.5)	0.343
Hypoglycemic drug	292 (53.1)	507 (56.1)	0.29
Lipid-lowering drug	541 (80.6)	849 (81.1)	0.861
Cholesterol, mg/dL	196.95 (40.72)	199.09 (42.78)	0.021
Triglycerides, mg/dL	144.45 (122.20)	173.82 (148.55)	<0.001
HDL, mg/dL	55.11 (16.11)	50.33 (15.56)	<0.001
LDL, mg/dL	113.00 (34.81)	116.14 (35.90)	0.006
eGFR, ml/min per 1.73 m ²	91.05 (25.24)	89.49 (24.42)	0.005

Data are presented as mean (SD) or n (%). PIR, poverty income ratio; BMI, body mass index; HBP, high blood pressure; DM, diabetes mellites; CVD, cardiovascular diseases; HDL, high density lipoprotein; LDL, low density lipoprotein; eGFR, estimated glomerular filtration rate.

Table 2. Association of OSA with cardiometabolic diseases.

Cardiometabolic diseases	OR	95% CI	P
Hypertension			
Model 1	1.44	[1.29, 1.60]	<0.001
Model 2	1.37	[1.22, 1.54]	<0.001
Model 3	1.30	[1.15, 1.46]	<0.001
Model 4	1.28	[1.14, 1.45]	<0.001
Diabetes			
Model 1	1.86	[1.65, 2.09]	<0.001
Model 2	1.85	[1.63, 2.09]	<0.001
Model 3	1.37	[1.20, 1.57]	<0.001
Model 4	1.46	[1.22, 1.76]	<0.001
Cardiovascular diseases			
Model 1	1.54	[1.32, 1.80]	<0.001
Model 2	1.51	[1.28, 1.79]	<0.001
Model 3	1.36	[1.14, 1.62]	0.001
Model 4	1.29	[1.08, 1.54]	0.006

Model 1: unadjusted.
Model 2: adjusted for gender, and age.
Model 3: adjusted for gender, age, race, education, PIR, BMI, drinking, smoking, and activity.
Model 4: adjusted for gender, age, race, education, PIR, BMI, drinking, smoking, activity, antihypertensive drugs, hypoglycemic drugs, lipid-lowering drugs, cholesterol, triglycerides, HDL, LDL, and eGFR.

Table 3. Association of OSA with all-cause and cardiovascular mortality.

	HR	95% CI	P
All-cause mortality			
Model 1	0.93	[0.82, 1.04]	0.206
Model 2	0.97	[0.86, 1.10]	0.634
Model 3	0.99	[0.87, 1.12]	0.886
Model 4	0.96	[0.84, 1.09]	0.507
Cardiovascular mortality			
Model 1	0.96	[0.72, 1.28]	0.787
Model 2	0.94	[0.70, 1.25]	0.658
Model 3	0.94	[0.70, 1.27]	0.685
Model 4	0.91	[0.67, 1.23]	0.532

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

351 **Figure 1. The flow chart of participant selection.**

For Peer Review




Figure 1. The flow chart of participant selection.