Table of Contents

1. PRISMA checklist
2. Characteristics of Included Studies
3. QUADAS 2 tool for ROB analysis
4. Search strategy

Supplementary Table 1: PRISMA checklist for systematic reviews

	#	Checklist item	Reported on page #
TITLE			
Title	1	Identify the report as a systematic review, meta-analysis, or both.	1
ABSTRACT			
Structured summary	2	Provide a structured summary including, as applicable: background; objectives; data sources; study eligibility criteria, participants, and interventions; study appraisal and synthesis methods; results; limitations; conclusions and implications of key findings; systematic review registration number.	4
INTRODUCTION			
Rationale	3	Describe the rationale for the review in the context of what is already known.	5
Objectives	4	Provide an explicit statement of questions being addressed with reference to participants, interventions, comparisons, outcomes, and study design (PICOS).	5
METHODS			
Protocol and registration	5	Indicate if a review protocol exists, if and where it can be accessed (e.g., Web address), and, if available, provide registration information including registration number.	5
Eligibility criteria	6	Specify study characteristics (e.g., PICOS, length of follow-up) and report characteristics (e.g., years considered, language, publication status) used as criteria for eligibility, giving rationale.	6
Information sources	7	Describe all information sources (e.g., databases with dates of coverage, contact with study authors to identify additional studies) in the search and date last searched.	5
Search	8	Present full electronic search strategy for at least one database, including any limits used, such that it could be repeated.	5
Study selection	9	State the process for selecting studies (i.e., screening, eligibility, included in systematic review, and, if applicable, included in the meta-analysis).	5
Data collection process	10	Describe method of data extraction from reports (e.g., piloted forms, independently, in duplicate) and any processes for obtaining and confirming data from investigators.	6
Data items	11	List and define all variables for which data were sought (e.g., PICOS, funding sources) and any assumptions and simplifications made.	6

Risk of bias in individual studies	12	Describe methods used for assessing risk of bias of individual studies (including specification of whether this was done at the study or outcome level), and how this information is to be used in any data synthesis.	6
Summary measures	13	State the principal summary measures (e.g., risk ratio, difference in means).	6
Synthesis of results	14	Describe the methods of handling data and combining results of studies, if done, including measures of consistency (e.g., I^2) for each meta-analysis.	6
Section/topic	**#**	**Checklist item**	**Reported on page #**
Risk of bias across studies	15	Specify any assessment of risk of bias that may affect the cumulative evidence (e.g., publication bias, selective reporting within studies).	6
Additional analyses	16	Describe methods of additional analyses (e.g., sensitivity or subgroup analyses, meta-regression), if done, indicating which were pre-specified.	6
RESULTS			
Study selection	17	Give numbers of studies screened, assessed for eligibility, and included in the review, with reasons for exclusions at each stage, ideally with a flow diagram.	7
Study characteristics	18	For each study, present characteristics for which data were extracted (e.g., study size, PICOS, follow-up period) and provide the citations.	7
Risk of bias within studies	19	Present data on risk of bias of each study and, if available, any outcome level assessment (see item 12).	Appendix
Results of individual studies	20	For all outcomes considered (benefits or harms), present, for each study: (a) simple summary data for each intervention group (b) effect estimates and confidence intervals, ideally with a forest plot.	7 - 8
Synthesis of results	21	Present results of each meta-analysis done, including confidence intervals and measures of consistency.	7 - 8
Risk of bias across studies	22	Present results of any assessment of risk of bias across studies (see Item 15).	Appendix
Additional analysis	23	Give results of additional analyses, if done (e.g., sensitivity or subgroup analyses, meta-regression [see Item 16]).	8
DISCUSSION			
Summary of evidence	24	Summarize the main findings including the strength of evidence for each main outcome; consider their relevance to key groups (e.g., healthcare providers, users, and policy makers).	7 - 8
Limitations	25	Discuss limitations at study and outcome level (e.g., risk of bias), and at review-level (e.g., incomplete retrieval of identified research, reporting bias).	9
Conclusions	26	Provide a general interpretation of the results in the context of other evidence, and implications for future research.	9

FUNDING			
Funding	27	Describe sources of funding for the systematic review and other support (e.g., supply of data); role of funders for the systematic review.	1

From: Moher D, Liberati A, Tetzlaff J, Altman DG, The PRISMA Group (2009). Preferred Reporting Items for Systematic Reviews and Meta-Analyses: The PRISMA Statement. PLoS Med 6(7): e1000097. doi:10.1371/journal.pmed1000097

Supplementary Table 2: Characteristics of Included Studies

Studies that compared SF ratio directly to PF ratio

Study and country	Patient population	Number of measurements	SF/PF correlation							
					PF 300		PF 200		PF 100	
			Equation	Spearman rho	SF	Sn, Sp, AUC	SF	Sn, Sp, AUC	SF	Sn, Sp, AUC
Wongsrichanalai 2019; Thailand	Medical ICU	735	PF = 18.28 + 0.81 SF	0.781	261	NR, NR, NR	180	NR, NR, NR	99	NR, NR, NR
Solomon 2010; Romania	ARDS	61	S/F = 111 + 0,67 x P/F.	NR	312	NR, NR, NR	245	NR, NR, NR	NR	NR, NR, NR
Sim 2013; UK	Necrotizing enterocolitis (bell stage II/III) in PICU	945	Quantile logistic regression	NR; Kendall's tau b 0.57	377	NR, NR, NR	270	NR, NR, NR	165	NR, NR, NR
Schmidt 2015; USA	ICU	7544	Log(PF ratio) = 1.07*Log(SF ratio) − 0.15	0.72	295	99.0; 9.9; 0.73	195	NR, NR, 0.84	102	NR, NR, 0.95
Sdiri 2023; Tunisia	ICU	206	Linear regression	0.583	NR	NR, NR, NR	NR	NR, NR, NR	NR	NR, NR, NR
Schmidt 2014; USA	ICU	25733	Log(PF ratio)=0.388+0.834*Log(SF ratio)	0.7	321	99.7; 4.0, NR	198	88.2; 56.6, NR	NR	NR, NR, NR
Sathiyakumar 2011; USA and Brazil	ICU	1710	Generalized estimation equation regression	0.59	NR	NR, NR, NR	NR	NR, NR, NR	NR	NR, NR, NR
Santana 2013; Brazil	ICU	118	Linear/Pearson coefficient	0.645	NR	NR, NR, NR	NR	NR, NR, NR	NR	NR, NR, NR
Rice 2007; USA	ARDS Network Trial	2673	S/F = 64 + 0.84 * (P/F)	0.89	315	91.0; 56.0, 0.88	235	85.0; 85.0, 0.93	NR	NR, NR, NR
Ray 2017; UK	PICU	1703	ln(PF) = {1.086 × ln(SF)} + 0.103	NR	NR	NR, NR, NR	NR	61.0, 73.0, NR	NR	NR, NR, NR
Philippe 2018; France	ICU	421	Rice and Ellis equation	0.84 [rice], 0.85 [ellis]	NR	NR, NR, NR	NR	NR, NR, NR	NR	NR, NR, NR
Pandharipande 2009; USA	Mechanically ventilated	4728	Log(PF) = 0.48 + 0.78 x Log(SF)	PEEP <8 0.58; PEEP 8 – 12 0.61; PEEP > 12 0.59	357	NR, NR, NR	214	NR, NR, NR	89	NR, NR, NR
Ortiz 2022; Colombia	ICU with invasive ventilatory support	664	Non-linear – Ellis equation Linear – Rice equation Log linear – Pandharipande equation	0.77	NR	NR, NR, NR	NR	NR, NR, NR	NR	NR, NR, NR
Neto 2013; Brazil and Netherlands	ICU	260	S/F ratio = 132.27 + 0.30 × (P/F)	0.487	241	NR, NR, 0.74	NR	NR, NR, NR	154	NR, NR, 0.926
Muniraman 2022; USA	Respiratory failure	1019	1/PF = – 0.0004304 + 2.0897987/SF	0.9	NR	NR, NR, NR	NR	NR, NR, NR	NR	NR, NR, 0.92

Lohano 2021; Pakistan	PICU	120	Linear/Pearson coefficient	0.688	NR	NR, NR, NR	NR	NR, NR, NR	NR	NR, NR, NR
Lobete Prieto 2011; Spain	PICU	40	P/F=37.277+(1.072×S/F) — (1.567×PtcCO2)	0.469	426	NR, NR, NR	295	NR, NR, NR	NR	NR, NR, NR
Lobete 2013; Spain	PICU	3248	1/SF = 0.00164 + 0.521/PF	0.843	296	90; 85; 0.95	236	86.0, 90.0, 0.952	146	54; 99; 0.978
Ahmed 2021; Pakistan	Patients in PICU on mechanical ventilator with a stay of at least 48 hours	30	Linear/Pearson coefficient	0.603	NR	NR, NR, NR	NR	NR, NR, NR	NR	NR, NR, NR
Aslanidis 2013, Greece	Mixed ICU population	290	Linear/Pearson coefficient	0.759	NR	NR, NR, NR	NR	NR, NR, NR	NR	NR, NR, NR
Babu 2021; India	Surgical intensive therapy unit	300	Linear/Pearson coefficient	0.66	323	72; 73; 0.80	285	73; 77; 0.84	NR	NR, NR, NR
Bashar 2018; USA	Cardiac surgery ICU	533	SF = 71.149 + 0.8 PF	0.857	311	90; 80; 0.92	NR	NR, NR, NR	NR	NR, NR, NR
Bhiri 2022; Tunisia	Covid-19 patients on mechanical ventilation	172	SF = 44.21 + 0.76 PF	NR	272.21	45; 90; NR	196	49; 84; NR	120	90; 76; NR
Bilan 2015; Iran	PICU	70	SF = 57 + 0.61 PF	NR	235	57; 100, 0.89	181	71; 82; 0.86	NR	NR, NR, NR
Cambiaso-Daniel 2017; USA	Burned children with smoke inhalation injury	23072	Peripheral capillary oxygenation/Fio2 ratios were divided into four subgroups based on peripheral capillary oxygenation values The r2 was 0.66 (23,072) for less than or equal to 100%, 0.87 (18,932) for less than or equal to 98%, 0.89 (7,056) for less than or equal to 95%, and 0.93 (4,229) for less than or equal to 92%. In the subgroup of patients who developed acute respiratory distress syndrome, r2 was 0.65 (8,357) for less than or equal to 100%, 0.89 (7,578) for less than or equal to 98%, 0.89 (4,115) for less than or equal to 95%, and 0.91 (2,288) less than or equal to 92%.	NR	NR	NR, NR, NR	NR	NR, NR, NR	NR	NR, NR, NR
Chen 2015; USA	Medical ICU	238 diagnosed by P/F; 124 diagnosed by S/F	NR	NR	NR	NR, NR, NR	NR	NR, NR, NR	NR	NR, NR, NR
Cinesi-Gomez 2017, Spain	Hospital emergency department	163	PF = 41.9 + 0.50 SF	0.745	NR	NR, NR, NR	NR	NR, NR, NR	NR	NR, NR, NR

Fukuda 2017; Japan	ICU	94	SF = 60.9 + 0.75 PF	0.87	NR	NR, NR, NR	NR	NR, NR, NR	NR	NR, NR, NR
Guirgis 2016; NR	ED	2796	NR	NR	NR	NR, NR, NR	NR	NR, NR, NR	NR	NR, NR, NR
James 2018; NR	PICU	250	NR	NR	NR	NR, NR, NR	180	90; 59; 0.8	NR	NR, NR, NR
Jusi 2012; Philippines	ICU	106	SF = 29.6 + 1.09 PF	0.804	357	100; 98.2; NR	248	100; 96; NR	NR	NR, NR, NR
Khemani 2009; USA	PICU	3143	Derivation: SF = 76 + 0.62 x PF Validation: SF = 111 + 0.51 x PF	0.61	263	86; 47, 0.792	213	68; 84; 0.85	NR	NR, NR, NR
Khemani 2015; USA	Children on mechanical ventilator	1833	NR	NR	264	NR, NR, NR	221	NR, NR, NR	150	NR, NR, NR
Kibe 2018; NR	NICU	348	Linear/Pearson coefficient	0.78 Preterm infants: 0.83 Term infants: 0.75	NR	NR, NR, NR	NR	NR, NR, NR	NR	NR, NR, NR
Kumar 2022; India	ICU	249	SF = 0.80 PF + 59.8	0.86	299	NR, NR, NR	219	NR, NR, NR	NR	NR, NR, NR
Lee 2020; NR	ICU	141535	Gradient boosting decision tree algorithm variant called Light GBM	AUC 0.782	NR	NR, NR, NR	NR	NR, NR, NR	NR	NR, NR, NR
Xia 2024, China	ICU	341	PF = -21.39 + 0.9162 SF	0.873	352	63; 93; NR	241	89; 62; NR	132	61; 99; NR
Singla 2024, India	Mechanically ventilated children in PICU	1237	NR	0.68	271	NR, NR, 0.770	235	NR, NR, 0.901	200	NR, NR, 0.889
Shah 2024, India	PICU	800	NR	NR	NR	NR, NR, NR	NR	NR, NR, NR	NR	NR
Jain 2024, India	Mechanically ventilated patients	1055	Hypoxemic (PaO2 < 60): PF = 9.19 + 0.68SF – $3.23e^{-4}SF^2$ Non-hypoxemic (PaO2 >60): PF = 35.24 +SF – $3.08e^{-4}SF^2$	0.92 (hypoxemic); -0.67 (non-hypoxemic)	NR	NR, NR, NR	NR	NR, NR, NR	NR	NR, NR, NR
Gandhi 2023, India	COVID patients admitted to ICU and requiring NIV or IMV	100	PF = - 35.211 + 1.1597 SF	NR	280	NR, NR, NR	203	NR, NR, NR	NR	NR, NR, NR
Beauchamp 2024, Canada	PICU	17239	Sauthier [52] non-linear equation	NR	NR	NR, NR, NR	NR	NR, NR, NR	NR	NR, NR, 0.9
Aiur 2023, India	Acute hypoxic respiratory failure patients in ICU	250	NR	0.387	321	70; 98; 0.877	252	69; 95; 0.831	NR	NR, NR, NR

Studies that imputed PF ratio from SF ratio

Study and country	Patient population	Number of measurements	SF/PF correlation							
					PF 300		PF 200		PF 100	
			Equation	Spearman rho	SF	Sn, Sp, AUC	SF	Sn, Sp, AUC	SF	Sn, Sp, AUC
Tabary 2022, USA	ICU patients with pseudomonas pneumonia	133 individuals; ABG collected 30 minutes after SpO2 collected	Neural net algorithm; support vector regression algorithm	For imputing PaO2 from SpO2 – 0.35; 0.35	N/A	N/A	N/A	N/A	N/A	N/A
Brown 2016, USA	Mechanically ventilated patients	1034 ABGs collected prospectively	Non-linear – Ellis equation Linear – Rice equation Log linear – Pandharipande equation	N/A	N/A	Non-linear 0.90, 0.67, 0.87 linear/log-linear NR, NR, 0.80	N/A	N/A	N/A	Non-linear 0.87, 0.91, 0.95 Linear/log-linear NR, NR, 0.92

The SF/PF correlation tab shows the spearman rho or correlation coefficient, along with the discriminating capacity of the SF cutoffs at determining the corresponding PF cutoffs of 100, 200 and 300.
ICU = intensive care unit, PICU = pediatrics intensive care unit, Sn = sensitivity, Sp = specificity, AUC = area under the curve, NR = not recorded, ARDS = acute respiratory distress syndrome, NIV = non-invasive ventilation, IMV = invasive mechanical ventilation

Supplementary Table 3: QUADAS 2 tool

Study and country	Risk of bias				Applicability			Overall Rating
	Patient Selection	Index Test	Reference Standard	Flow and Timing	Patient Selection	Index Test	Reference Standard	
Wongsrichanalai 2019; Thailand	Unclear	Low	Low	Unclear	Low	Low	Low	Intermediate
Solomon 2010; Romania	Unclear	Low	Low	Low	Low	Low	Low	Low
Sim 2013; UK	Unclear	Low	Low	Unclear	Unclear	Low	Low	Intermediate
Schmidt 2015; USA	Low	Low	Low	Low	Low	Low	Low	Low
Sdiri 2023; Tunisia	Unclear	Low	Low	Unclear	Low	Low	Low	Intermediate
Schmidt 2014; USA	Low	Low	Low	Low	Low	Low	Low	Low
Sathiyakumar 2011; USA and Brazil	Low	Low	Low	Low	Low	Low	Low	Low
Santana 2013; Brazil	Unclear	Low	Low	Unclear	Low	Low	Low	Intermediate
Rice 2007; USA	Low	Low	Low	Low	Low	Low	Low	Low
Ray 2017; UK	Unclear	Low	Low	Low	Low	Low	Low	Low
Philippe 2018; France	Low	Low	Low	Low	Low	Low	Low	Low
Pandharipande 2009; USA	Low	Low	Low	Low	Low	Low	Low	Low
Ortiz 2022; Colombia	Low	Low	Low	Unclear	Low	Low	Low	Low
Neto 2013; Brazil and Netherlands	Low	Low	Low	Low	Low	Low	Low	Low
Muniraman 2022; USA	Low	Low	Low	Unclear	Low	Low	Low	Low
Lohano 2021; Pakistan	Unclear	Low	Low	Unclear	Low	Low	Low	Low
Lobete Prieto 2011; Spain	Low	Low	Low	Low	Low	Low	Low	Low
Lobete 2013; Spain	Low	Low	Low	Low	Low	Low	Low	Low
Ahmed 2021; Pakistan	Unclear	Low	Low	Low	Low	Low	Low	Low
Aslanidis 2013, Greece	Unclear	Low	Low	Unclear	Low	Low	Low	Intermediate
Babu 2021; India	Low	Low	Low	Unclear	Low	Low	Low	Low
Bashar 2018; USA	Low	Low	Low	Low	Low	Low	Low	Low
Bhiri 2022; Tunisia	Low	Low	Low	Low	Low	Low	Low	Low

Bilan 2015; Iran	Unclear	Low	Low	Low	Low	Low	Low	Low
Cambiaso-Daniel 2017; USA	Unclear	Low	Low	Low	High	Low	Low	High
Chen 2015; USA	Low	Low	Low	Unclear	Low	Low	Low	Intermediate
Cinesi-Gomez 2017, Spain	Unclear	Low	Low	Low	Low	Low	Low	Intermediate
Fukuda 2017; Japan	Low	Low	Low	Unclear	Low	Low	Low	Intermediate
Guirgis 2016; NR	Unclear	Low	Low	Unclear	Low	Low	Low	Intermediate
James 2018; NR	Low	Low	Low	Low	Low	Low	Low	Low
Jusi 2012; Philippines	Unclear	Low	Low	Unclear	Low	Low	Low	Intermediate
Khemani 2009; USA	Low	Low	Low	Unclear	Low	Low	Low	Low
Khemani 2015; USA	Low	Low	Low	Unclear	Low	Low	Low	Low
Kibe 2018; NR	Unclear	Low	Low	Unclear	Low	Low	Low	Intermediate
Kumar 2022; India	Unclear	Low	Low	Low	Low	Low	Low	Low
Lee 2020; NR	Unclear	Low	Low	Unclear	Low	Low	Low	Intermediate
Xia 2024, China	Low	Low	Low	Low	Low	Low	Low	Low
Singla 2024, India	Unclear	Low	Low	Low	Low	Low	Low	Low
Shah 2024, India	Low	Low	Low	Low	Low	Low	Low	Low
Jain 2024, India	Low	Low	Low	Unclear	Low	Low	Low	Low
Gandhi 2023, India	Unclear	Low	Low	Low	Low	Low	Low	Low
Beauchamp 2024, Canada	Unclear	Low	Low	Low	Low	Low	Low	Low
Aiur 2023, India	Low	Low	Low	Unclear	Low	Low	Low	Low
Tabary 2022, USA	Low	Low	Low	Low	Low	Low	Low	Low
Brown 2016, USA	Low	Low	Low	Low	Low	Low	Low	Low

SF/PF Ratios – SR – Literature Search (update)

Research Question(s):

1. How well does the SF ratio correlate with the PF ratio in critically ill patients?

Population	Any critically ill human patients
Intervention	SF ratio or (Pulse oximetric saturation OR peripheral oxygen saturation OR SPO2)/(Fraction of inspired oxygen OR FiO2) OR PF ratio or (PaO2 OR partial pressure of oxygen in arterial blood)/Fraction of inspired oxygen OR FiO2)
Comparison	
Outcomes	Any measure of correlation or comparison
Setting	Any type of observational study, studies examining diagnostic test characteristics, cross sectional studies that specifically compare SF and PF ratio and to other measures of hypoxia or patient important outcomes (mortality, ICU length of stay, mechanical ventilation, NIV, HFNC). We will exclude case series, case reports, review articles and editorials. We will exclude animal studies and those done in healthy volunteers.

Seed items: (19029434 OR 19242333 OR 25859313).ui.
Search by: Kaitryn Campbell
Requestor: Dipayan Chaudhuri / Bram Rochwerg
Date(s): 2024 Nov 5 (update, 2023 Oct-current)
(20231* OR 2024* OR 2025*).ed,dt. use medall OR (20231* OR 2024* OR 2025*).dc,dd. use oemezd
Limits: NOT Animal only (Ovid databases only)
Databases: Ovid Medline, Embase; The Cochrane Library; Web of Science
Filters: None
Search Output: RIS (for Covidence)

Ovid

Database(s): **Embase** 1974 to 2024 November 04**, Ovid MEDLINE(R) ALL** 1946 to November 04, 2024
Search Strategy:

#	Searches	Results
1	Spo2 Fio2 ratio.mp.	427
2	(SF ratio? or "(SF) ratio?" or "[SF] ratio?" or Sfratio? or "(SF)ratio?" or "[SF]ratio?").mp.	324
3	((SPO2* or "Spo(2)*" or "Spo[2]*" or SPO-2* or "Spo-(2)*" or "Spo-[2]*") adj5 (ratio or ratios)).mp.	984
4	exp Oximetry/	54526
5	(pulse* adj8 saturat*).ti,kf,kw.	1343
6	(pulse* adj3 saturat*).ab.	7659
7	Pulse Oximetry/ and saturat*.mp.	15912
8	(4 and (or/5-6)) or 7	16212
9	Oxygen/	441530
10	Oxygen Consumption/	217929
11	Arterial Oxygen Saturation/	13022
12	Arterial Oxygen Tension/	14438
13	Blood Oxygen Tension/	10321
14	Oxygen Tension/	22679
15	(peripheral* adj8 saturat*).ti,kf,kw.	437
16	(peripheral* adj3 saturat*).ab.	5636
17	Oxygen Saturation/ and peripheral*.mp.	6474
18	((or/9-14) and (or/15-16)) or 17	7718

19	(((peripheral* or pulse*) adj (oximetr* or oxygen* or O2) adj8 saturat*) or ((peripheral* or pulse*) adj8 (oximetr* or oxygen* or O2) adj saturat*)).mp.	14172
20	8 or 18 or 19	29329
21	(ratio or ratios).mp.	4267917
22	20 and 21	3095
23	1 or 2 or 3 or 22 [SF ratio Concept]	4077
24	(PF ratio? or "(PF) ratio?" or "[PF] ratio?" or PFratio? or "(PF)ratio?" or "[PF]ratio?").mp.	692
25	((PaO2* or "Pao(2)*" or "Pao[2]*" or PaO-2* or "Pao-(2)*" or "Pao-[2]*") adj5 (ratio or ratios)).mp.	8710
26	(partial pressur* and ((arterial* or arteries or artery) adj (blood* or circulati*))).mp.	7141
27	(or/9-14) and 26	3868
28	(partial pressur* adj2 (oxygen* or O2) adj5 (arterial* or arteries or artery) adj (blood* or circulati*)).mp.	1045
29	(or/27-28) and 21	881
30	24 or 25 or 29 [PF ratio Concept]	9835
31	("fraction of inspired oxygen*" or FiO2* or "Fio(2)*" or "Fio[2]*" or "FiO-2*" or "Fio-(2)*" or "Fio-[2]*").mp. [FiO2 Concept]	39493
32	23 and 30 and 31 [All Concepts, combined]	332
33	(exp animals/ or exp animal experimentation/ or exp animal experiment/ or exp models animal/ or nonhuman/ or exp vertebrate/ or exp vertebrates/) not (exp humans/ or exp human experimentation/ or exp human experiment/) [Animal NOT Human]	12785874
34	32 not 33 [Animal Only studies removed]	309
35	remove duplicates from 34	245
36	(20231* or 2024* or 2025*).ed,dt. use medall or (20231* or 2024* or 2025*).dc,dd. use oemezd	3625645
37	35 and 36	39 [35 after duplicates removed]

The Cochrane Library (CENTRAL)

Line	Terms	Results
#1	("SF ratio" OR "SF ratios" OR "(SF) ratio" OR "(SF) ratios" OR "[SF] ratio" OR "[SF] ratios" OR Sfratio? OR "(SF)ratio" OR "(SF)ratios" OR "[SF]ratio" OR "[SF]ratios") OR ((SPO2* OR "Spo(2)" OR "Spo[2]" OR SPO 2* OR "Spo (2)" OR "Spo [2]") NEAR/5 (ratio OR ratios))	0
#2	MeSH descriptor: [Oximetry] explode all trees	1401
#3	(pulse* NEAR/8 saturat*):ti OR (pulse* NEAR/8 saturat*):kw OR (pulse* NEAR/3 saturat*):ab	2122
#4	MeSH descriptor: [Oxygenators] in all MeSH products	136
#5	MeSH descriptor: [Oxygen Consumption] this term only	8047
#6	(peripheral* NEAR/8 saturat*):ti OR (peripheral* NEAR/8 saturat*):kw OR (peripheral* NEAR/3 saturat*):ab	1669
#7	(((peripheral* OR pulse*) NEXT (oximetr* OR oxygen* OR O2) NEAR/8 saturat*) OR ((peripheral* OR pulse*) NEAR/8 (oximetr* OR oxygen* OR O2) NEXT saturat*)):ti,ab,kw	8561
#8	(ratio OR ratios)	184101
#9	(#2 AND #3) OR ((#4 OR #5) AND #6) OR #7	8571
#10	#8 AND #9	893
#11	("PF ratio" OR "PF ratios" OR "(PF) ratio" OR "(PF) ratios" OR "[PF] ratio" OR "[PF] ratios" OR PFratio? OR "(PF)ratio" OR "(PF)ratios" OR "[PF]ratio" OR "[PF]ratios") OR ((PaO2* OR "Pao(2)" OR "Pao[2]" OR PaO 2* OR "Pao (2)" OR "Pao [2]") NEAR/5 (ratio OR ratios))	0
#12	(partial pressur* AND ((arterial* OR arteries OR artery) NEXT (blood* OR circulati*))):ti,ab,kw	931
#13	(partial pressur* NEAR/2 (oxygen* OR O2) ADJ5 ((arterial* OR arteries OR artery) NEXT (blood* OR circulati*)))	13
#14	((#4 OR #5) AND #12) OR #13	50
#15	#8 AND #14	20
#16	("fraction of inspired oxygen" OR FiO2* OR "Fio(2)" OR "Fio[2]" OR "FiO-2" OR "Fio-(2)" OR "Fio[2]")	4922
#17	#10 AND #15 AND #16 [with Publication Year from 2023 to 2024, with Cochrane Library publication date in The last year, in Trials]	0

Web of Science Core Collection

#	Searches	Results
1	ALL=(SF ratio? or "(SF) ratio?" or "[SF] ratio?" OR Sfratio? or "(SF)ratio?" or "[SF]ratio?")	8,069
2	TS=((SPO2* OR "Spo(2)*" OR "Spo[2]*" OR SPO-2* OR "Spo-(2)*" OR "Spo-[2]*") NEAR/5 (ratio OR ratios))	403
3	TI=((SPO2* OR "Spo(2)*" OR "Spo[2]*" OR SPO-2* OR "Spo-(2)*" OR "Spo-[2]*") NEAR/5 (ratio OR ratios))	45
4	AB=((SPO2* OR "Spo(2)*" OR "Spo[2]*" OR SPO-2* OR "Spo-(2)*" OR "Spo-[2]*") NEAR/5 (ratio OR ratios))	361
5	AK=((SPO2* OR "Spo(2)*" OR "Spo[2]*" OR SPO-2* OR "Spo-(2)*" OR "Spo-[2]*") NEAR/5 (ratio OR ratios))	12
6	#5 OR #4 OR #3 OR #2 OR #1	8,461
7	TS=(((peripheral* OR pulse*) NEAR/1 (oximetr* OR oxygen* OR O2) NEAR/8 saturat*) OR ((peripheral* OR pulse*) NEAR/8 (oximetr* OR oxygen* OR O2) NEAR/1 saturat*))	5,921
8	TI=(((peripheral* OR pulse*) NEAR/1 (oximetr* OR oxygen* OR O2) NEAR/8 saturat*) OR ((peripheral* OR pulse*) NEAR/8 (oximetr* OR oxygen* OR O2) NEAR/1 saturat*))	471
9	AB=(((peripheral* OR pulse*) NEAR/1 (oximetr* OR oxygen* OR O2) NEAR/8 saturat*) OR ((peripheral* OR pulse*) NEAR/8 (oximetr* OR oxygen* OR O2) NEAR/1 saturat*))	5,639
10	AK=(((peripheral* OR pulse*) NEAR/1 (oximetr* OR oxygen* OR O2) NEAR/8 saturat*) OR ((peripheral* OR pulse*) NEAR/8 (oximetr* OR oxygen* OR O2) NEAR/1 saturat*))	119
11	#7 OR #8 OR #9 OR #10	5,921
12	ALL=(ratio OR ratios)	3,468,800
13	#11 AND #12	683
14	#6 OR #13	9,055
15	ALL=(PF ratio? OR "(PF) ratio?" OR "[PF] ratio?" OR PFratio? OR "(PF)ratio?" OR "[PF]ratio?")	4,641
16	TS=((PaO2* OR "Pao(2)*" OR "Pao[2]*" OR PaO-2* OR "Pao-(2)*" OR "Pao-[2]*") NEAR/5 (ratio OR ratios))	2,954
17	TI=((PaO2* OR "Pao(2)*" OR "Pao[2]*" OR PaO-2* OR "Pao-(2)*" OR "Pao-[2]*") NEAR/5 (ratio OR ratios))	126
18	AB=((PaO2* OR "Pao(2)*" OR "Pao[2]*" OR PaO-2* OR "Pao-(2)*" OR "Pao-[2]*") NEAR/5 (ratio OR ratios))	2,859
19	AK=((PaO2* OR "Pao(2)*" OR "Pao[2]*" OR PaO-2* OR "Pao-(2)*" OR "Pao-[2]*") NEAR/5 (ratio OR ratios))	63
20	#15 OR #16 OR #17 OR #18 OR #19	7,570
21	TS=(partial pressur* NEAR/2 (oxygen* OR O2) NEAR/5 (arterial* OR arteries OR artery) NEAR/1 (blood* OR circulati*))	1,619
22	TI=(partial pressur* NEAR/2 (oxygen* OR O2) NEAR/5 (arterial* OR arteries OR artery) NEAR/1 (blood* OR circulati*))	13
23	AB=(partial pressur* NEAR/2 (oxygen* OR O2) NEAR/5 (arterial* OR arteries OR artery) NEAR/1 (blood* OR circulati*))	1,303
24	AK=(partial pressur* NEAR/2 (oxygen* OR O2) NEAR/5 (arterial* OR arteries OR artery) NEAR/1 (blood* OR circulati*))	4
25	#21 OR #22 OR #23 OR #24	1,619
26	#25 AND #12	344
27	#20 OR #26	7,850
28	ALL=("fraction of inspired oxygen*" OR FiO2* OR "Fio(2)*" OR "Fio[2]*" OR "FiO-2*" OR "Fio-(2)*" OR "Fio-[2]*")	17,545
29	#28 AND #27 AND #14 | Timespan: 2023-10-01 to 2024-09-05 (Index Date)	16 [3 after duplicates removed]