

# CSE 397 / EM 397 - Stabilized and Variational Multiscale Methods in CFD

## Homework #4

Thomas J.R. Hughes\*, Frimpong A. Baidoo† and Geonyeong Lee‡

*Oden Institute for Computational Engineering and Sciences, University of Texas at Austin*

Spring 2024

The compressible Navier-Stokes equations can be written as

$$\mathbf{U}_{,t} + \mathbf{F}_{i,i} = \mathbf{F}_{i,i}^{\text{visc}} + \mathbf{F}_{i,i}^{\text{heat}} + \mathcal{F}$$

where

$$\mathbf{U} = \rho \begin{pmatrix} 1 \\ u_1 \\ u_2 \\ u_3 \\ e \end{pmatrix}, \quad \mathbf{F}_i = u_i \mathbf{U} + p \begin{pmatrix} 0 \\ \delta_{1i} \\ \delta_{2i} \\ \delta_{3i} \\ u_i \end{pmatrix}$$
$$\mathbf{F}_i^{\text{visc}} = \begin{pmatrix} 0 \\ \tau_{1i} \\ \tau_{2i} \\ \tau_{3i} \\ \tau_{ij}u_j \end{pmatrix}, \quad \mathbf{F}_i^{\text{heat}} = \begin{pmatrix} 0 \\ 0 \\ 0 \\ 0 \\ -q_i \end{pmatrix}, \quad \mathcal{F} = \begin{pmatrix} 0 \\ \rho b_1 \\ \rho b_2 \\ \rho b_3 \\ \rho(b_i u_i + r) \end{pmatrix}$$

For perfect gases, the constitutive relations are

$$\begin{aligned} e &= \iota + \frac{1}{2}|\mathbf{u}|^2, \\ \iota &= c_v \theta, \\ p &= (\gamma - 1)\rho\iota, \\ \gamma &= c_p/c_v \\ \tau_{ij} &= 2\mu\epsilon_{ij} + \lambda\epsilon_{kk}\delta_{ij}, \\ \epsilon_{ij} &= \frac{1}{2}(u_{i,j} + u_{j,i}) \\ q_i &= -\kappa\theta_{,i}, \\ s &= \ln\left(\frac{p}{p_0}\left(\frac{\rho}{\rho_0}\right)^{-\gamma}\right). \end{aligned}$$

---

\*hughes@oden.utexas.edu

†fabaidoo@utexas.edu

‡geon@utexas.edu

**Exercise 4.1**

The entropy variables  $\mathbf{V}$  are defined as

$$\mathbf{V}^T := \frac{\partial H}{\partial \mathbf{U}},$$

where  $H := -\rho s$ . Show that the entropy variables can be written as

$$\mathbf{V} = \frac{1}{\iota} \begin{pmatrix} \nu - \frac{1}{2}|\mathbf{u}|^2 \\ u_1 \\ u_2 \\ u_3 \\ -1 \end{pmatrix},$$

where  $\nu := \iota + p/\rho - \iota s$  is the electrochemical potential. Note that there are many different but equivalent ways to express  $V_1$ .

**Exercise 4.2**

Show that

$$\begin{aligned} \mathbf{V} \cdot \mathbf{F}_{i,i} &= (Hu_i)_{,i}, \\ \mathbf{V} \cdot \mathbf{F}_i^{\text{visc}} &= 0, \\ \mathbf{V} \cdot \mathbf{F}_i^{\text{heat}} &= \frac{q_i}{c_v \theta}, \\ \mathbf{V} \cdot \mathcal{F}_i &= -\frac{\rho r}{c_v \theta}. \end{aligned}$$

**Exercise 4.3**

Show that

$$\begin{aligned} \mathbb{D} &:= \mathbf{V}_{,i} \cdot (\mathbf{F}_i^{\text{visc}} + \mathbf{F}_i^{\text{heat}}) \\ &= a \epsilon_{ij}^{\text{dev}}(\mathbf{u}) \epsilon_{ij}^{\text{dev}}(\mathbf{u}) + b (\nabla \cdot \mathbf{u})^2 + c |\nabla \theta|^2, \end{aligned}$$

where

$$\epsilon_{ij}^{\text{dev}}(\mathbf{u}) := \frac{1}{2}(u_{i,j} + u_{j,i}) - \frac{1}{3}u_{k,k}\delta_{ij}.$$

Express  $a$ ,  $b$ , and  $c$  in terms of the constitutive coefficients  $\mu$ ,  $\lambda$ , and  $\kappa$ . Determine the sign of  $\mathbb{D}$ .

**Exercise 4.4**

If we define  $\mathbf{R}$  as the residual of the compressible Navier-Stokes equations:

$$\mathbf{R} := \mathbf{U}_t + \mathbf{F}_{i,i} - \mathbf{F}_{i,i}^{\text{visc}} - \mathbf{F}_{i,i}^{\text{heat}} - \mathcal{F},$$

show that

$$0 = \mathbf{V} \cdot \mathbf{R} = H_{,t} + (Hu_i)_{,i} - \left( \frac{q_i}{\theta} \right)_{,i} + \frac{\rho r}{\theta} + c_v \mathbb{D}.$$

The entropy inequality follows from this result.