```
In [1]: import sympy
        def rat(expr):
            for i in expr.atoms(sympy.Float):
                r = sympy.Rational(str(i)).limit_denominator(1000)
                expr = expr.subs(i, r)
            return expr
In [2]: from IPython.display import display, Markdown
        import sympy
        from sympy import *
        import numpy as np
        x, xi, u = symbols(r'x \xi u', real=True)
        d1, d2, d3 = symbols(r'd_1 d_2 d_3', real=True)
        Xi = [ -1/sqrt(3), 1/sqrt(3) ]
        W = [1, 1]
        kappa = Function(r"K")(xi)
        dkappa_du = Function(r"K_{,u}")(xi)
        Na = [
        1/2*xi*(xi-1),
        1-xi*xi,
        1/2*xi*(xi+1)
        display( Markdown( f"$N_1 = {sympy.latex(Na[0])}$" ))
        display( Markdown( f"$N_2 = {sympy.latex(Na[1])}$" ))
        display( Markdown(f"$N_3 = {sympy.latex(Na[2])}$"))
      N_1 = 0.5\xi(\xi - 1)
      N_2=1-\xi^2
      N_3 = 0.5\xi(\xi+1)
In [3]: dN1_xi = diff(Na[0], xi)
        dN2_xi = diff(Na[1], xi)
        dN3_xi = diff(Na[2], xi)
        display( Markdown( r"$dN_1/d\xi = "+ f"{sympy.latex(dN1_xi)}$" ))
        display( Markdown( r"$dN_2/d\xi = "+ f"{sympy.latex(dN2_xi)}$" ))
        display( Markdown( r"$dN_3/d\xi = "+ f"{sympy.latex(dN3_xi)}$" ))
      dN_1/d\xi = 1.0\xi - 0.5
      dN_2/d\xi = -2\xi
      dN_3/d\xi = 1.0\xi + 0.5
In [4]: X = Na[1] * 0.5 + Na[2]
        X = simplify(X)
        dx dxi = diff(X,xi)
        dxi_dx = 1/dx_dxi
        display( Markdown( r"x(xi) = "+ f"{sympy.latex(X)}"))
```

```
display( Markdown( r"dx/dxi = "+ f"{sympy.latex(dx_dxi)}" ))
         display( Markdown( r"$d\xi/dx = "+ f"{sympy.latex(dxi_dx)}$" ))
       x(\xi) = 0.5\xi + 0.5
       dx/d\xi = 0.5
       d\xi/dx = 2.0
In [5]: dN1_x = dN1_xi * dxi_dx
         dN2_x = dN2_xi * dxi_dx
         dN3_x = dN3_xi * dxi_dx
         display( Markdown( r"$dN_1/dx = "+ f"{sympy.latex(dN1_x)}$" ))
         display( Markdown( r"$dN_2/dx = "+ f"{sympy.latex(dN2_x)}$" ))
         display( Markdown( r"$dN_3/dx = "+ f"{sympy.latex(dN3_x)}$" ))
       dN_1/dx = 2.0\xi - 1.0
       dN_2/dx = -4.0\xi
       dN_3/dx = 2.0\xi + 1.0
In [11]: def build_N_x( xi_ ) :
             # Derivatives in X space
             N1_x = dN1_xi.subs(xi, xi_) * dxi_dx
             N2_x = dN2_xi.subs(xi, xi_) * dxi_dx
             N3_x = dN3_xi.subs(xi, xi_) * dxi_dx
             return N1_x, N2_x, N3_x
         db = [d1, d2, d3]
         ne = [0, 0, 0]
         f = symbols(r'f_1 f_2 f_3', real=True)
         h = symbols(r'h', real=True)
         fe = [0, 0, 0]
         dna_ddb = zeros(3, 3)
         fcol = []
         for xi_, W_ in zip( Xi, W ) :
             Na_x = build_N_x(xi_)
             x_{=} = X.subs(xi, xi_{=})
             kappa_ = kappa.subs( xi, xi_ )
             dkappa_du_ = dkappa_du.subs( xi, xi_ )
             fcol.append(kappa_)
             fcol.append(dkappa_du_)
             Na_{-} = [0, 0, 0]
             for a in range(3) : Na_[a] = simplify(Na[a].subs(xi, xi_))
             q = 0
             for b in range(3) : q += db[b] * Na_x[b]
             for a in range(3) :
                 ne[a] += W_* dx_dxi * Na_x_[a] * q * kappa_
                 fe[a] += W_ * dx_dxi * Na_[a] * f[a]
                 fe[a] = simplify(fe[a])
             for a in range(3) :
                 for b in range(3) :
                     dna_ddb[a,b] += W_ * dx_dxi * Na_x_[a] * Na_[b] * q * dkappa_du_
                     dna_db[a,b] += W_* dx_dxi * kappa_* Na_x[a] * Na_x[b]
```

```
fe[0] += W_ * h * Na_[0]
                                                                                                        fe[0] = simplify(fe[0])
                                                                          for a in range(3) :
                                                                                                        ne[a] = cancel(ne[a])
                                                                                                        for c in fcol : ne[a] = collect(ne[a],c)
                                                                                                        for d in db : ne[a] = collect( ne[a], d )
                                                                                                        ne[a] = simplify(ne[a],full=True)
                                                                          for a in range(3) :
                                                                                                        fe[a] = nsimplify(fe[a])
                                                                          for a in range(3) :
                                                                                                        for b in range(3) :
                                                                                                                                       dna_ddb[a,b] = cancel(dna_ddb[a,b])
                                                                                                                                       for c in fcol : dna_ddb[a,b] = collect(dna_ddb[a,b],c)
                                                                                                                                       for d in db : dna_ddb[a,b] = collect( dna_ddb[a,b], d )
                                                                                                                                       dna ddb[a,b] = simplify(dna ddb[a,b])
In [12]: for a in range(3):
                                                                                                        display( Markdown( f"$n_{a+1}^e(d^e) = {sympy.latex(rat(ne[a]))}$" ))
                                                     n_1^e(d^e) = \left(d_1\left(rac{7}{6} - rac{2\sqrt{3}}{3}
ight) - d_2\left(rac{4}{3} - rac{2\sqrt{3}}{3}
ight) + rac{d_3}{6}
ight)K\left(rac{\sqrt{3}}{3}
ight) + \left(d_1\left(rac{2\sqrt{3}}{3} + rac{7}{6}
ight) - d_2\left(rac{2\sqrt{3}}{3} + rac{4}{3}
ight) + rac{d_3}{6}
ight)K\left(-rac{\sqrt{3}}{3}
ight)
                                                     n_2^e(d^e) = -\left(d_1\left(rac{4}{3} - rac{2\sqrt{3}}{3}
ight) - rac{8d_2}{3} + d_3\left(rac{2\sqrt{3}}{3} + rac{4}{3}
ight)
ight)K\!\left(rac{\sqrt{3}}{3}
ight) - \left(d_1\left(rac{2\sqrt{3}}{3} + rac{4}{3}
ight) - rac{8d_2}{3} + d_3\left(rac{4}{3} - rac{2\sqrt{3}}{3}
ight)
ight)K\!\left(-rac{\sqrt{3}}{3}
ight)
                                                     n_3^e(d^e) = \left(rac{d_1}{6} - d_2\left(rac{4}{3} - rac{2\sqrt{3}}{3}
ight) + d_3\left(rac{7}{6} - rac{2\sqrt{3}}{3}
ight)
ight)K\left(-rac{\sqrt{3}}{3}
ight) + \left(rac{d_1}{6} - d_2\left(rac{2\sqrt{3}}{3} + rac{4}{3}
ight) + d_3\left(rac{2\sqrt{3}}{3} + rac{7}{6}
ight)
ight)K\left(rac{\sqrt{3}}{3}
ight)
In [13]: for a in range(3) :
                                                                                                        display( Markdown( f"$f {a+1}^e = {sympy.latex(rat(fe[a]))}$" ))
                                                     f_1^e = \frac{f_1}{6} + \frac{h}{2}
                                                     f_2^e = \frac{2f_2}{2}
                                                     f_{2}^{e} = \frac{f_{3}}{c}
In [15]: for a in range(3):
                                                                                                        for b in range(3) :
                                                                                                                                       display( Markdown( r"\frac{n''}{r^2} and n''+f''(a+1)''+r''^e} = "+ f"\frac{n''}{r^2} = "+ f"\frac{n''}{r^2} (rat(dna_ddb[a,b]))}$" ))
                                                     \frac{\partial n_1^e}{\partial d_1^e} = \left(d_1\left(\frac{19}{36} - \frac{11\sqrt{3}}{36}\right) - d_2\left(\frac{5}{9} - \frac{\sqrt{3}}{3}\right) + \frac{d_3\left(1-\sqrt{3}\right)}{36}\right)K_{,u}\left(\frac{\sqrt{3}}{3}\right) + \left(d_1\left(\frac{19}{36} + \frac{11\sqrt{3}}{36}\right) - d_2\left(\frac{5}{9} + \frac{\sqrt{3}}{3}\right) + \frac{d_3\left(1+\sqrt{3}\right)}{36}\right)K_{,u}\left(-\frac{\sqrt{3}}{3}\right) + \left(\frac{2\sqrt{3}}{3} + \frac{7}{6}\right)K\left(-\frac{\sqrt{3}}{3}\right) + \left(\frac{7}{6} - \frac{2\sqrt{3}}{3}\right)K\left(\frac{\sqrt{3}}{3}\right) + \left(\frac{1}{2} + \frac{11\sqrt{3}}{3}\right) +
                                                     \frac{\partial n_1^e}{\partial d_1^e} = \left(d_1\left(\frac{7}{9} - \frac{4\sqrt{3}}{9}\right) - d_2\left(\frac{8}{9} - \frac{4\sqrt{3}}{9}\right) + \frac{d_3}{9}\right)K_{,u}\left(\frac{\sqrt{3}}{3}\right) + \left(d_1\left(\frac{4\sqrt{3}}{9} + \frac{7}{9}\right) - d_2\left(\frac{4\sqrt{3}}{9} + \frac{8}{9}\right) + \frac{d_3}{9}\right)K_{,u}\left(-\frac{\sqrt{3}}{3}\right) - \left(\frac{2\sqrt{3}}{3} + \frac{4}{3}\right)K\left(-\frac{\sqrt{3}}{3}\right) - \left(\frac{4}{3} - \frac{2\sqrt{3}}{3}\right)K\left(\frac{\sqrt{3}}{3}\right) + \left(\frac{4\sqrt{3}}{9} + \frac{7}{9}\right) - d_2\left(\frac{4\sqrt{3}}{9} + \frac{8}{9}\right) + \frac{d_3}{9}\right)K_{,u}\left(-\frac{\sqrt{3}}{3}\right) - \left(\frac{2\sqrt{3}}{3} + \frac{4}{3}\right)K\left(-\frac{\sqrt{3}}{3}\right) - \left(\frac{4\sqrt{3}}{3} - \frac{2\sqrt{3}}{3}\right)K\left(\frac{\sqrt{3}}{3}\right) + \left(\frac{4\sqrt{3}}{9} + \frac{4\sqrt{3}}{9}\right)K_{,u}\left(-\frac{\sqrt{3}}{3}\right) - \left(\frac{4\sqrt{3}}{3} - \frac{2\sqrt{3}}{3}\right)K\left(-\frac{\sqrt{3}}{3}\right) + \left(\frac{\sqrt{3}}{3} - \frac{2\sqrt{3}}{3}\right)K\left(-\frac{\sqrt{3}}{3}\right)K\left(-\frac{\sqrt{3}}{3}\right) + \left(\frac{\sqrt{3}}{3} - \frac{2\sqrt{3}}{3}\right)K\left(-\frac{\sqrt{3}}{3}\right)K\left(-\frac{\sqrt{3}}{3}\right) + \left(\frac{\sqrt{3}}{3} - \frac{2\sqrt{3}}{3}\right)K\left(-\frac{\sqrt{3}}{3}\right)K\left(-\frac{\sqrt{3}}{3}\right)K\left(-\frac{\sqrt{3}}{3}\right)K\left(-\frac{\sqrt{3}}{3}\right)K\left(-\frac{\sqrt{3}}{3}\right)K\left(-\frac{\sqrt{3}}{3}\right)K\left(-\frac{\sqrt{3}}{3}\right)K\left(-\frac{\sqrt{3}}{3}\right)K\left(-\frac{\sqrt{3}}{3}\right)K\left(-\frac{\sqrt{3}}{3}\right)K\left(-\frac{\sqrt{3}}{3}\right)K\left(-\frac{\sqrt{3}}{3}\right)K\left(-\frac{\sqrt{3}}{3}\right)K\left(-\frac{\sqrt{3}}{3}\right)K\left(-\frac{\sqrt{3}}{3}\right)K\left(-\frac{\sqrt{3}}{3}\right)K\left(-\frac{\sqrt{3}}{3}\right)K\left(-\frac{\sqrt{3}}{3}\right)K\left(-\frac{\sqrt{3}}{3}\right)K\left(-\frac{\sqrt{3}}{3}\right)K\left(-\frac{\sqrt{3}}{3}\right)K\left(-\frac{\sqrt{3}}{3}\right)K\left(-\frac{\sqrt{3}}{3}\right)K\left(-\frac{\sqrt{3}}{3}\right)K\left(-\frac{\sqrt{3}}{3}\right)K\left(-\frac{\sqrt{3}}{3}\right)K\left(-\frac{\sqrt{3}}{3}\right)K\left(-\frac{\sqrt{3}}{3}
                                                     \frac{\partial n_1^e}{\partial d_3^e} = \left(-d_1\left(\frac{5}{36} - \frac{\sqrt{3}}{12}\right) + d_2\left(\frac{1}{9} - \frac{\sqrt{3}}{9}\right) + \frac{d_3\left(1 + \sqrt{3}\right)}{36}\right)K_{,u}\left(\frac{\sqrt{3}}{3}\right) + \left(-d_1\left(\frac{5}{36} + \frac{\sqrt{3}}{12}\right) + d_2\left(\frac{1}{9} + \frac{\sqrt{3}}{9}\right) + \frac{d_3\left(1 - \sqrt{3}\right)}{36}\right)K_{,u}\left(-\frac{\sqrt{3}}{3}\right) + \frac{K\left(-\frac{\sqrt{3}}{3}\right)}{6} + \frac{K\left(\frac{\sqrt{3}}{3}\right)}{6} + \frac{K\left(\frac{\sqrt{3}}{3}\right)}{6
                                                     \frac{\partial n_2^e}{\partial d_1^e} = \left(-d_1\left(\frac{5}{9} - \frac{\sqrt{3}}{3}\right) + \frac{4d_2\left(1 - \sqrt{3}\right)}{9} + d_3\left(\frac{1}{9} + \frac{\sqrt{3}}{9}\right)\right)K_{,u}\left(\frac{\sqrt{3}}{3}\right) + \left(-d_1\left(\frac{5}{9} + \frac{\sqrt{3}}{3}\right) + \frac{4d_2\left(1 + \sqrt{3}\right)}{9} + d_3\left(\frac{1}{9} - \frac{\sqrt{3}}{9}\right)\right)K_{,u}\left(-\frac{\sqrt{3}}{3}\right) - \left(\frac{2\sqrt{3}}{3} + \frac{4}{3}\right)K\left(-\frac{\sqrt{3}}{3}\right) - \left(\frac{4}{3} - \frac{2\sqrt{3}}{3}\right)K\left(\frac{\sqrt{3}}{3}\right) + \frac{4d_2\left(1 - \sqrt{3}\right)}{9} + d_3\left(\frac{1}{9} - \frac{\sqrt{3}}{9}\right)K_{,u}\left(-\frac{\sqrt{3}}{3}\right) - \left(\frac{2\sqrt{3}}{3} + \frac{4}{3}\right)K\left(-\frac{\sqrt{3}}{3}\right) - \left(\frac{4}{3} - \frac{2\sqrt{3}}{3}\right)K\left(\frac{\sqrt{3}}{3}\right)
```

$$\frac{\partial n_2^e}{\partial d_2^e} = -\left(d_1\left(\frac{8}{9} - \frac{4\sqrt{3}}{9}\right) - \frac{16d_2}{9} + d_3\left(\frac{4\sqrt{3}}{9} + \frac{8}{9}\right)\right)K_{,u}\left(\frac{\sqrt{3}}{3}\right) - \left(d_1\left(\frac{4\sqrt{3}}{9} + \frac{8}{9}\right) - \frac{16d_2}{9} + d_3\left(\frac{8}{9} - \frac{4\sqrt{3}}{9}\right)\right)K_{,u}\left(-\frac{\sqrt{3}}{3}\right) + \frac{8K\left(-\frac{\sqrt{3}}{3}\right)}{3} + \frac{8K\left(\frac{\sqrt{3}}{3}\right)}{3} \\ \frac{\partial n_2^e}{\partial d_3^e} = \left(d_1\left(\frac{1}{9} - \frac{\sqrt{3}}{9}\right) + \frac{4d_2\left(1 + \sqrt{3}\right)}{9} - d_3\left(\frac{5}{9} + \frac{\sqrt{3}}{3}\right)\right)K_{,u}\left(\frac{\sqrt{3}}{3}\right) + \left(d_1\left(\frac{1}{9} + \frac{\sqrt{3}}{9}\right) + \frac{4d_2\left(1 - \sqrt{3}\right)}{9} - d_3\left(\frac{5}{9} - \frac{\sqrt{3}}{3}\right)\right)K_{,u}\left(-\frac{\sqrt{3}}{3}\right) - \left(\frac{4}{3} - \frac{2\sqrt{3}}{3}\right)K\left(-\frac{\sqrt{3}}{3}\right) - \left(\frac{2\sqrt{3}}{3} + \frac{4}{3}\right)K\left(\frac{\sqrt{3}}{3}\right) + \frac{4d_2\left(1 - \sqrt{3}\right)}{9} - d_3\left(\frac{5}{9} - \frac{\sqrt{3}}{3}\right)K\left(-\frac{\sqrt{3}}{3}\right) - \left(\frac{2\sqrt{3}}{3} + \frac{4}{3}\right)K\left(\frac{\sqrt{3}}{3}\right) + \frac{4d_2\left(1 - \sqrt{3}\right)}{9} - d_3\left(\frac{5}{9} - \frac{\sqrt{3}}{3}\right)K\left(-\frac{\sqrt{3}}{3}\right) - \left(\frac{2\sqrt{3}}{3} + \frac{4}{3}\right)K\left(\frac{\sqrt{3}}{3}\right) + \frac{4d_2\left(1 - \sqrt{3}\right)}{9} - d_3\left(\frac{5}{9} - \frac{\sqrt{3}}{3}\right)K\left(-\frac{\sqrt{3}}{3}\right) - \left(\frac{2\sqrt{3}}{3} + \frac{4}{3}\right)K\left(\frac{\sqrt{3}}{3}\right) + \frac{4d_2\left(1 - \sqrt{3}\right)}{9} - d_3\left(\frac{5}{9} - \frac{\sqrt{3}}{3}\right)K\left(-\frac{\sqrt{3}}{3}\right) - \left(\frac{2\sqrt{3}}{3} + \frac{4}{3}\right)K\left(\frac{\sqrt{3}}{3}\right) + \frac{4d_2\left(1 - \sqrt{3}\right)}{9} - d_3\left(\frac{5}{9} - \frac{\sqrt{3}}{3}\right)K\left(-\frac{\sqrt{3}}{3}\right) + \frac{4d_2\left(1 - \sqrt{3}\right)}{9} - d_3\left(\frac{5}{9} - \frac{\sqrt{3}}{3}\right)K\left(-\frac{\sqrt{3}}{3}\right)K\left(-\frac{\sqrt{3}}{3}\right) + \frac{4d_2\left(1 - \sqrt{3}\right)}{9} - \frac{4d_2\left(1 - \sqrt{3}\right)}{9} - \frac{4d_2\left(1 - \sqrt{3}\right)}{9} - \frac{4d_2\left(1 - \sqrt{3}\right)}{9} - \frac{4d_2\left(1 - \sqrt{3}$$

$$\frac{\partial n_3^e}{\partial d_1^e} = \left(\frac{d_1\left(1-\sqrt{3}\right)}{36} + d_2\left(\frac{1}{9} + \frac{\sqrt{3}}{9}\right) - d_3\left(\frac{5}{36} + \frac{\sqrt{3}}{12}\right)\right) K_{,u}\left(\frac{\sqrt{3}}{3}\right) + \left(\frac{d_1\left(1+\sqrt{3}\right)}{36} + d_2\left(\frac{1}{9} - \frac{\sqrt{3}}{9}\right) - d_3\left(\frac{5}{36} - \frac{\sqrt{3}}{12}\right)\right) K_{,u}\left(-\frac{\sqrt{3}}{3}\right) + \frac{K\left(-\frac{\sqrt{3}}{3}\right)}{6} + \frac{K\left(\frac{\sqrt{3}}{3}\right)}{6} +$$

In [ ]: