

Sensory and Motor Systems.

Basics of Life Science

Thursdays 9-10:30

Ray Luo


ray LC

rluo@aoni.waseda.jp

WASEDA UNIVERSITY | SILS



Sensory reception and transmission transduces signals to nervous system.




ray LC

rluo@aoni.waseda.jp

WASEDA UNIVERSITY | SILS



Sensory perception depends on pathway and source of sensation.



Reference URL

ray LC

rluo@aoni.waseda.jp

WASEDA UNIVERSITY | SILS



Sensory receptor types depend on the modality of the stimuli.

- Mechanoreceptor – pressure, touch, motion.
- Chemoreceptor – osmolarity, glucose, oxygen.
- Electromagnetic receptor – mag field, electric.
- Thermoreceptor – temperature, TRP capsaicin.
- Nociceptor – pain, amplified by prostaglandin.



Chemical pheromone receptors on silkworm antennae


ray LC

rluo@aoni.waseda.jp

WASEDA UNIVERSITY | SILS



Statoliths inside statocyst mechanoreceptors sense gravity, equilibrium.




ray LC

rluo@aoni.waseda.jp

WASEDA UNIVERSITY | SILS



The Senses:
Hearing.




ray LC

rluo@aoni.waseda.jp

WASEDA UNIVERSITY | SILS



Sound is transduced by the eardrum to middle ear (hammer, anvil, stirrups).




© 2014 Pearson Education, Inc.

ray LC

rluo@aoni.waseda.jp

WASEDA UNIVERSITY | SILS

Stirrups transmit vibrations to oval window which opens to the cochlea.



Fluid pressure in vestibular canal causes vibrations.


© 2014 Pearson Education, Inc.

ray LC

rluo@aoni.waseda.jp

WASEDA UNIVERSITY | SILS

Organ of Corti on floor of the cochlear duct transduces sound to perception.





ray LC

rluo@aoni.waseda.jp

WASEDA UNIVERSITY | SILS



Hair cells on tectorial membrane are bent when basilar membrane vibrates.



Mainly up and down motion of basilar


ray LC

rluo@aoni.waseda.jp

WASEDA UNIVERSITY | SILS



Basilar membrane vibrations cause hair cells to fire and refract -> sense.



Mechanical distortion -> hair cell channels open or close

ray LC



rluo@aoni.waseda.jp

WASEDA UNIVERSITY | SILS



Volume - amplitude of hair cell deflect;
pitch - location of deflect on cochlea.

Basilar membrane unrolled




ray LC

rluo@aoni.waseda.jp

WASEDA UNIVERSITY | SILS



Pressure waves travel: oval window through cochlea to round window.




ray LC

rluo@aoni.waseda.jp

WASEDA UNIVERSITY | SILS

Utricle and saccule of semicircular canals sense equilibrium in humans.



ray LC

rluo@aoni.waseda.jp

WASEDA UNIVERSITY | SILS

Review: Hearing.

- Which sensory cells are common to the senses of hearing and equilibrium in humans?
 - A) otolithocytes
 - B) vestibular cells
 - C) ocelli
 - D) tectorial cells
 - E) hair cells
- The function of the basilar membrane is to
 - A) transmit vibrations from the tympanic membrane to the oval window.
 - B) vibrate up and down in response to the fluid pressure waves in the vestibular canal.
 - C) vibrate in response to moving air reaching the outer ear.
 - D) create pressure waves in the perilymph (fluid inside the cochlea).



ray LC

rluo@aoni.waseda.jp

WASEDA UNIVERSITY | SILS



The Senses: Vision.



ray LC

rluo@aoni.waseda.jp

WASEDA UNIVERSITY | SILS




Insects: motion-sensitive compound eyes with many ommatidia detectors.



(a) Fly eyes

2 mm

Makes it VERY hard to sneak up on insects!



(b) Ommatidia

Ommatidium

ray LC

rluo@aoni.waseda.jp

WASEDA UNIVERSITY | SILS




The human eye focuses by changing the shape of its single lens.



Pupil: light entry

Glaucoma:
failing to drain
aqueous humor



© Pearson Education, Inc.
© 2014 Pearson Education, Inc.

ray LC

rluo@aoni.waseda.jp


WASEDA UNIVERSITY | SILS



Two types of photoreceptors sensitive to light and colors are on the retina.

Rod: low acuity, high sensitivity, peripheral, BW

Cone (three types): high acuity, low sensitivity, central (foveal), color




ray LC

rluo@aoni.waseda.jp

WASEDA UNIVERSITY | SILS

Light → cis-to-trans retinal conversion → hydrolyze cGMP → close Na channel




Saturation in bright light → temporary blindness

ray LC

rluo@aoni.waseda.jp

WASEDA UNIVERSITY | SILS

Bipolar and ganglion cells integrate photoreceptor info at the retina.




ray LC

rluo@aoni.waseda.jp

WASEDA UNIVERSITY | SILS

Lateral inhibition by horizontal and amacrine cells enhance contrast.




ray LC

rluo@aoni.waseda.jp

WASEDA UNIVERSITY | SILS

Ganglion cell axons exit the optic disk and form the optic nerve to the LGN.




ray LC

rluo@aoni.waseda.jp

WASEDA UNIVERSITY | SILS

Defects in red or green photopsins of cones on X-chrom leads to color blind.




ray LC

rluo@aoni.waseda.jp

WASEDA UNIVERSITY | SILS

Day time vision relies on high acuity cones in the fovea; night time on rods.



ray LC

rluo@aoni.waseda.jp

WASEDA UNIVERSITY | SILS

Lens is thickened in order to see near objects; flattened for distant.


(a) Near vision (accommodation)

Ciliary muscles contract, pulling border of choroid toward lens.

Suspensory ligaments relax.

Lens becomes thicker and rounder, focusing on nearby objects.

© 2014 Pearson Education, Inc.




(b) Distance vision

Ciliary muscles relax, and border of choroid moves away from lens.

Suspensory ligaments pull against lens.

Lens becomes flatter, focusing on distant objects.

© 2014 Pearson Education, Inc.




ray LC

rluo@aoni.waseda.jp

WASEDA UNIVERSITY | SILS

Visual cortex is organized into columns specific for orientations of vis scene.



ray LC

rhuo@aoni.waseda.jp

WASEDA UNIVERSITY | SILS



Review: Vision.


- In vertebrate eyes, the conversion of light energy to chemical energy occurs most directly as the result of changes to
 - A) phosphodiesterase.
 - B) Cyclic GMP (cGMP).
 - C) opsin.
 - D) retinal.
 - E) phosphodiesterase.
- Rod photoreceptors exposed to light will
 - A) depolarize due to the opening of sodium channels.
 - B) depolarize due to the opening of potassium channels.
 - C) hyperpolarize due to the opening of sodium channels.
 - D) hyperpolarize due to the closing of sodium channels.
 - E) hyperpolarize due to the opening of potassium channels.

ray LC



rhuo@aoni.waseda.jp

WASEDA UNIVERSITY | SILS






Taste buds contain sensory cells for each taste (sweet umami bitter GPCR)



Odor sensors are neurons that go directly to the olfactory bulb.




© 2014 Pearson Education, Inc.

ray LC

rluo@aoni.waseda.jp

WASEDA UNIVERSITY | SILS

Odors bind to one of thousands of receptors -> cAMP -> Na Ca enter.



Nature Reviews | Neuroscience

ray LC

rluo@aoni.waseda.jp

WASEDA UNIVERSITY | SILS

Effectors: Muscular and Skeletal systems.




ray LC

rluo@aoni.waseda.jp

WASEDA UNIVERSITY | SILS



Muscle is made up of sarcomeres (M line of myosin, Z line of actin).




ray LC

rluo@aoni.waseda.jp

WASEDA UNIVERSITY | SILS



Myosin and actin stay the same length during muscle contraction.



© 2014 Pearson Education, Inc.


ray LC

rluo@aoni.waseda.jp

WASEDA UNIVERSITY | SILS



ATP hydrolysis and release triggers myosin release and bind to actin.




ray LC

rluo@aoni.waseda.jp

WASEDA UNIVERSITY | SILS



ATP hydrolysis leads to myosin head cross linked to actin, then ADP release.



ray LC

rluo@aoni.waseda.jp

WASEDA UNIVERSITY | SILS

Regulation of muscle contraction at the level of the sarcomere.


- Creatine phosphate and glycogen breakdown to glucose provide phosphates to ATP.
- Intense activity -> lactic acid fermentation.
- Tropomyosin and troponin bound to actin.
- Calcium binds to troponin and moves tropomyosin away from myosin binding sites.
- Motor neuron degeneration => ALS; antibodies to mus AChR => myasthenia gravis.

ray LC

rluo@aoni.waseda.jp

WASEDA UNIVERSITY | SILS

Calcium exposes myosin binding sites, allowing muscle contraction.




ray LC

rluo@aoni.waseda.jp

WASEDA UNIVERSITY | SILS

Action potential at the terminal of motor neuron \rightarrow acetylcholine release.


© Pearson Education, Inc.
© 2014 Pearson Education, Inc.

ray LC

rluo@aoni.waseda.jp

WASEDA UNIVERSITY | SILS

Action potential at the fiber along T tubules -> calcium release from SR.




ray LC

rluo@aoni.waseda.jp

WASEDA UNIVERSITY | SILS

Motor unit is the motor neuron and all fibers it synapses on (one cell a fiber).




ray LC

rluo@aoni.waseda.jp

WASEDA UNIVERSITY | SILS

Actual movement requires flexing and relaxing antagonistic muscle pairs.




ray LC

rloo@aoni.waseda.jp

WASEDA UNIVERSITY | SILS



Locomotion example: tube feet of echinoderms (starfish).



ray LC

rloo@aoni.waseda.jp

WASEDA UNIVERSITY | SILS

