

99. Ranking Protein Docking Results by knowledge-based Potentials

Vera Grimm ¹ Dietmar Schomburg ²

Keywords: unbound docking, knowledge-based potentials

Abstract

Two knowledge-based potentials have been developed to discriminate native protein-protein complex structures from non-native ones. One potential is based on the *inverse* Boltzmann equation [1] while the other one represents an empirical approach. Both potentials include separate weighting factors for functional groups and h-bonds to stress the importance of these type of interactions. Furthermore a solvent-dependent part is introduced, as well as a trapeze function for smoothing the distribution functions. The generated distribution functions for both potentials are critically examined. To compare the performance of both potentials several unbound protein-protein docking studies are carried out. Besides criteria such as geometric, hydrophobic and electrostatic complementarity an appropriate “energy”-function can be used to rank complex structures [2]. The dockings are performed with CKORDO, a modified and improved reimplement of KORDO [3] based on the Fourier Correlation Technique [4]. The results indicate that the empirical potential in combination with the introduced weighting of functional groups shows the highest discriminatory power.

References

- [1] Sippl, M. J. 1990. Calculation of conformational ensembles from potentials of mean force. An approach to the knowledge-based prediction of local structures in globular proteins. *Journal of Molecular Biology*, 213:859–883.
- [2] Ben-Naim, A. and Ting, K. L. and Jernigan, R. L. 1990. Solvent effect on binding thermodynamics of biopolymers. *Biopolymers*, 2:901-919.
- [3] Meyer, M. and Wilson, P. and Schomburg, D. 1996. Hydrogen bonding and molecular surface shape complementarity as a basis for protein docking. *Journal of Molecular Biology*, 264:199–210.
- [4] Katchalski-Katzir, E. and Shariv, I. and Eisenstein, M. and Friesem, A. A. and Aflalo, C. and Vakser, I. A. 1992. Molecular surface recognition: determination of geometric fit between proteins and their ligands by correlation techniques. *Proceedings of the National Academy of Sciences USA*, 89:2195-2199.

¹Department of Biochemistry, University of Cologne, Germany, vera.grimm@uni-koeln.de

²Department of Biochemistry, University of Cologne, Germany, d.schomburg@uni-koeln.de