

Subject Index to Volume 145 (2009)

Adaptation

Women experience greater heat pain adaptation and habituation than men, **145**, 350

Alcoholic neuropathy

Tocotrienol ameliorates behavioral and biochemical alterations in the rat model of alcoholic neuropathy, **145**, 129

Allodynia

In vivo silencing of the Cav3.2 T-type calcium channels in sensory neurons alleviates hyperalgesia in rats with streptozocin-induced diabetic neuropathy, **145**, 184

Allodynia

Prefrontal cortex and spinal cord mediated anti-neuropathy and analgesia induced by sarcosine, a glycine-T1 transporter inhibitor, **145**, 176

Allodynia

Sigma-1 receptors regulate activity-induced spinal sensitization and neuropathic pain after peripheral nerve injury, **145**, 294

Allodynia

Tocotrienol ameliorates behavioral and biochemical alterations in the rat model of alcoholic neuropathy, **145**, 129

Analgesia

fMRI investigation of the effect of local and systemic lidocaine on noxious electrical stimulation-induced activation in spinal cord, **145**, 110

Antigen-induced arthritis

Gait abnormalities differentially indicate pain or structural joint damage in monoarticular antigen-induced arthritis, **145**, 142

Antinociception

Direct excitation of spinal GABAergic interneurons by noradrenaline, **145**, 204

Anxiety

A prospective comparison of post-surgical behavioral pain scales in preschoolers highlighting the risk of false evaluations, **145**, 60

Assessment

Pain language and gender differences when describing a past pain event, **145**, 86

Bacterial protein toxins

The Rac GTPase-activating bacterial protein toxin CNF1 induces analgesia up-regulating μ -opioid receptors, **145**, 219

Behavior

The Rac GTPase-activating bacterial protein toxin CNF1 induces analgesia up-regulating μ -opioid receptors, **145**, 219

Behavioral rating scales

A prospective comparison of post-surgical behavioral pain scales in preschoolers highlighting the risk of false evaluations, **145**, 60

Brain

Pain relief through expectation supersedes descending inhibitory deficits in fibromyalgia patients, **145**, 18

Brain mapping

Sex differences in functional brain activation during noxious visceral stimulation in rats, **145**, 120

Burning

Women experience greater heat pain adaptation and habituation than men, **145**, 350

BV-weighted

fMRI investigation of the effect of local and systemic lidocaine on noxious electrical stimulation-induced activation in spinal cord, **145**, 110

Calcium

In vivo silencing of the Cav3.2 T-type calcium channels in sensory neurons alleviates hyperalgesia in rats with streptozocin-induced diabetic neuropathy, **145**, 184

Capsaicin

Electroacupuncture suppresses capsaicin-induced secondary hyperalgesia through an endogenous spinal opioid mechanism, **145**, 332

Carpal tunnel syndrome

Differential involvement of A-delta and A-beta fibres in neuropathic pain related to carpal tunnel syndrome, **145**, 105

Catastrophizing

A prospective sequential analysis of the fear-avoidance model of pain, **145**, 45

Catastrophizing

Momentary pain and coping in temporomandibular disorder pain: Exploring mechanisms of cognitive behavioral treatment for chronic pain, **145**, 160

CatWalk

The effect of Spinal Cord Stimulation in mice with chronic neuropathic pain after partial ligation of the sciatic nerve, **145**, 312

Central sensitization

Sigma-1 receptors regulate activity-induced spinal sensitization and neuropathic pain after peripheral nerve injury, **145**, 294

Cerebral blood flow

Sex differences in functional brain activation during noxious visceral stimulation in rats, **145**, 120

Children

A prospective comparison of post-surgical behavioral pain scales in preschoolers highlighting the risk of false evaluations, **145**, 60

Chronic discogenic low back pain

A randomized double-blind controlled trial of intra-annular radiofrequency thermal disc therapy – A 12-month follow-up, **145**, 279

Chronic neuropathic pain

The effect of Spinal Cord Stimulation in mice with chronic neuropathic pain after partial ligation of the sciatic nerve, **145**, 312

Chronic pain

Nondermatomal somatosensory deficits in patients with chronic pain disorder: Clinical findings and hypometabolic pattern in FDG-PET, **145**, 252

- Chronic pain management in managed care
Trends in prescribed opioid therapy for non-cancer pain for individuals with prior substance use disorders, **145**, 287
- Chronic pain Muscle input
Enhanced central pain processing of fibromyalgia patients is maintained by muscle afferent input: A randomized, double-blind, placebo-controlled study, **145**, 96
- Circumplex
Psychodynamic interpersonal therapy and improvement in interpersonal difficulties in people with severe irritable bowel syndrome, **145**, 196
- Cognitive-behavioral treatment
Momentary pain and coping in temporomandibular disorder pain: Exploring mechanisms of cognitive behavioral treatment for chronic pain, **145**, 160
- Colorectal distension
Sex differences in functional brain activation during noxious visceral stimulation in rats, **145**, 120
- Complex Regional Pain Syndrome
HLA-B62 and HLA-DQ8 are associated with Complex Regional Pain Syndrome with fixed dystonia, **145**, 82
- Complex Regional Pain Syndrome
Is there an association between psychological factors and the Complex Regional Pain Syndrome type 1 (CRPS1) in adults? A systematic review, **145**, 52
- Conditioning
Placebo conditioning and placebo analgesia modulate a common brain network during pain anticipation and perception, **145**, 24
- Conversion disorder
Nondermatomal somatosensory deficits in patients with chronic pain disorder: Clinical findings and hypometabolic pattern in FDG-PET, **145**, 252
- Coping
Momentary pain and coping in temporomandibular disorder pain: Exploring mechanisms of cognitive behavioral treatment for chronic pain, **145**, 160
- Cornea
An fMRI case report of photophobia: Activation of the trigeminal nociceptive pathway, **145**, 358
- Counseling skills
Medical students retain pain assessment and management skills long after an experiential curriculum: A controlled study, **145**, 319
- CRPS
Ketamine and chronic pain – Going the distance, **145**, 271
- CRPS
Ketamine produces effective and long-term pain relief in patients with Complex Regional Pain Syndrome Type 1, **145**, 304
- CRPS
Nondermatomal somatosensory deficits in patients with chronic pain disorder: Clinical findings and hypometabolic pattern in FDG-PET, **145**, 252
- Cytokines
Role of substance P signaling in enhanced nociceptive sensitization and local cytokine production after incision, **145**, 341
- Cytoskeleton
The Rac GTPase-activating bacterial protein toxin CNF1 induces analgesia up-regulating μ -opioid receptors, **145**, 219
- D-Cycloserine
Prefrontal cortex and spinal cord mediated anti-neuropathy and analgesia induced by sarcosine, a glycine-T1 transporter inhibitor, **145**, 176
- Descending inhibition
Direct excitation of spinal GABAergic interneurons by noradrenaline, **145**, 204
- Descending inhibition
Pain relief through expectation supersedes descending inhibitory deficits in fibromyalgia patients, **145**, 18
- Diabetes
In vivo silencing of the $Ca_v3.2$ T-type calcium channels in sensory neurons alleviates hyperalgesia in rats with streptozocin-induced diabetic neuropathy, **145**, 184
- Diary
Pain site frequency and location in sickle cell disease: The PiSCES project, **145**, 246
- DiscTRODE
A randomized double-blind controlled trial of intra-annular radiofrequency thermal disc therapy – A 12-month follow-up, **145**, 279
- Disease modulation
Ketamine and chronic pain – Going the distance, **145**, 271
- Disease modulation
Ketamine produces effective and long-term pain relief in patients with Complex Regional Pain Syndrome Type 1, **145**, 304
- Dorsal root ganglia
Experimental arthritis causes tumor necrosis factor- α -dependent infiltration of macrophages into rat dorsal root ganglia which correlates with pain-related behavior, **145**, 151
- Dystonia
HLA-B62 and HLA-DQ8 are associated with Complex Regional Pain Syndrome with fixed dystonia, **145**, 82
- Editorial
Ketamine and chronic pain – Going the distance, **145**, 271
- EGFP
Direct excitation of spinal GABAergic interneurons by noradrenaline, **145**, 204
- Electroacupuncture
Electroacupuncture suppresses capsaicin-induced secondary hyperalgesia through an endogenous spinal opioid mechanism, **145**, 332
- ERK
Sigma-1 receptors regulate activity-induced spinal sensitization and neuropathic pain after peripheral nerve injury, **145**, 294
- Etanercept
Gait abnormalities differentially indicate pain or structural joint damage in monoarticular antigen-induced arthritis, **145**, 142
- Expectations
Pain relief through expectation supersedes descending inhibitory deficits in fibromyalgia patients, **145**, 18
- Experience sampling
Momentary pain and coping in temporomandibular disorder pain: Exploring mechanisms of cognitive behavioral treatment for chronic pain, **145**, 160
- Eye
An fMRI case report of photophobia: Activation of the trigeminal nociceptive pathway, **145**, 358
- Fabry disease
Functional and structural nerve fiber findings in heterozygote patients with Fabry disease, **145**, 237
- Facial expression of pain
The smile of pain, **145**, 273
- False-evaluations
A prospective comparison of post-surgical behavioral pain scales in preschoolers highlighting the risk of false evaluations, **145**, 60
- Fear of movement
A prospective sequential analysis of the fear-avoidance model of pain, **145**, 45
- Fibromyalgia
Enhanced central pain processing of fibromyalgia patients is maintained by muscle afferent input: A randomized, double-blind, placebo-controlled study, **145**, 96
- Fibromyalgia
Pain relief through expectation supersedes descending inhibitory deficits in fibromyalgia patients, **145**, 18
- Fibromyalgia syndrome
Treatment of fibromyalgia syndrome with gabapentin and pregabalin – A meta-analysis of randomized controlled trials, **145**, 69

fMRI

fMRI investigation of the effect of local and systemic lidocaine on noxious electrical stimulation-induced activation in spinal cord, **145**, 110

fMRI

Placebo conditioning and placebo analgesia modulate a common brain network during pain anticipation and perception, **145**, 24

Frequency

Pain site frequency and location in sickle cell disease: The PiSCES project, **145**, 246

Functional brain imaging

Nondermatomal somatosensory deficits in patients with chronic pain disorder: Clinical findings and hypometabolic pattern in FDG-PET, **145**, 252

Functional hemisynndrome

Nondermatomal somatosensory deficits in patients with chronic pain disorder: Clinical findings and hypometabolic pattern in FDG-PET, **145**, 252

GABA

Direct excitation of spinal GABAergic interneurons by noradrenaline, **145**, 204

Gabapentin

Treatment of fibromyalgia syndrome with gabapentin and pregabalin – A meta-analysis of randomized controlled trials, **145**, 69

Gait analysis

Gait abnormalities differentially indicate pain or structural joint damage in monoarticular antigen-induced arthritis, **145**, 142

Gavage

Prefrontal cortex and spinal cord mediated anti-neuropathy and analgesia induced by sarcosine, a glycine-T1 transporter inhibitor, **145**, 176

Gender

Pain language and gender differences when describing a past pain event, **145**, 86

Gender role expectations

Gender role expectations of pain is associated with pain tolerance limit but not with pain threshold, **145**, 230

Genetic association

HLA-B62 and HLA-DQ8 are associated with Complex Regional Pain Syndrome with fixed dystonia, **145**, 82

Habituation

Women experience greater heat pain adaptation and habituation than men, **145**, 350

Headache

Activation of the prostaglandin system in response to sleep loss in healthy humans: Potential mediator of increased spontaneous pain, **145**, 136

Heat hyperalgesia

Role of substance P signaling in enhanced nociceptive sensitization and local cytokine production after incision, **145**, 341

Heat pain

Women experience greater heat pain adaptation and habituation than men, **145**, 350

Hemophilia

An assessment of the relative influence of pain coping, negative thoughts about pain, and pain acceptance on health-related quality of life among people with hemophilia, **145**, 169

HLA class I

HLA-B62 and HLA-DQ8 are associated with Complex Regional Pain Syndrome with fixed dystonia, **145**, 82

HLA class II

HLA-B62 and HLA-DQ8 are associated with Complex Regional Pain Syndrome with fixed dystonia, **145**, 82

Human

An fMRI case report of photophobia: Activation of the trigeminal nociceptive pathway, **145**, 358

Hyperalgesia

In vivo silencing of the Ca_v3.2 T-type calcium channels in sensory neurons alleviates hyperalgesia in rats with streptozocin-induced diabetic neuropathy, **145**, 184

Hyperalgesia

Electroacupuncture suppresses capsaicin-induced secondary hyperalgesia through an endogenous spinal opioid mechanism, **145**, 332

Hyperalgesia

Enhanced central pain processing of fibromyalgia patients is maintained by muscle afferent input: A randomized, double-blind, placebo-controlled study, **145**, 96

Hyperalgesia

Factors contributing to large analgesic effects in placebo mechanism studies conducted between 2002 and 2007, **145**, 36

Hyperalgesia

Sigma-1 receptors regulate activity-induced spinal sensitization and neuropathic pain after peripheral nerve injury, **145**, 294

Hyperalgesia

Tocotrienol ameliorates behavioral and biochemical alterations in the rat model of alcoholic neuropathy, **145**, 129

IBS

Psychodynamic interpersonal therapy and improvement in interpersonal difficulties in people with severe irritable bowel syndrome, **145**, 196

In vitro

Direct excitation of spinal GABAergic interneurons by noradrenaline, **145**, 204

Inflammation

Activation of the prostaglandin system in response to sleep loss in healthy humans: Potential mediator of increased spontaneous pain, **145**, 136

Intra-annular radiofrequency thermal treatment

A randomized double-blind controlled trial of intra-annular radiofrequency thermal disc therapy – A 12-month follow-up, **145**, 279

Joint pain

An assessment of the relative influence of pain coping, negative thoughts about pain, and pain acceptance on health-related quality of life among people with hemophilia, **145**, 169

Joint pain

Gait abnormalities differentially indicate pain or structural joint damage in monoarticular antigen-induced arthritis, **145**, 142

Ketamine

Ketamine and chronic pain – Going the distance, **145**, 271

Ketamine

Ketamine produces effective and long-term pain relief in patients with Complex Regional Pain Syndrome Type 1, **145**, 304

Laser

Placebo conditioning and placebo analgesia modulate a common brain network during pain anticipation and perception, **145**, 24

Laser-evoked potentials

Differential involvement of A-delta and A-beta fibres in neuropathic pain related to carpal tunnel syndrome, **145**, 105

Lidocaine

Enhanced central pain processing of fibromyalgia patients is maintained by muscle afferent input: A randomized, double-blind, placebo-controlled study, **145**, 96

Lidocaine

fMRI investigation of the effect of local and systemic lidocaine on noxious electrical stimulation-induced activation in spinal cord, **145**, 110

Light

An fMRI case report of photophobia: Activation of the trigeminal nociceptive pathway, **145**, 358

Locomotor behavior

Gait abnormalities differentially indicate pain or structural joint damage in monoarticular antigen-induced arthritis, **145**, 142

- Low-voltage-activated
 - In vivo* silencing of the $\text{Ca}_v3.2$ T-type calcium channels in sensory neurons alleviates hyperalgesia in rats with streptozocin-induced diabetic neuropathy, **145**, 184
- Macrophage
 - Experimental arthritis causes tumor necrosis factor- α -dependent infiltration of macrophages into rat dorsal root ganglia which correlates with pain-related behavior, **145**, 151
- Magnitude
 - Factors contributing to large analgesic effects in placebo mechanism studies conducted between 2002 and 2007, **145**, 36
- Measurement
 - Using normalized EMG to define the nociceptive flexion reflex (NFR) threshold: Further evaluation of standardized NFR scoring criteria, **145**, 211
- Mechanical allodynia
 - Role of substance P signaling in enhanced nociceptive sensitization and local cytokine production after incision, **145**, 341
- Mechanisms of treatment
 - Momentary pain and coping in temporomandibular disorder pain: Exploring mechanisms of cognitive behavioral treatment for chronic pain, **145**, 160
- Medical education
 - Medical students retain pain assessment and management skills long after an experiential curriculum: A controlled study, **145**, 319
- Medically unexplained symptoms
 - The unbearable lightness of somatisation: A systematic review of the concept of somatisation in empirical studies of pain, **145**, 31
- Meta-analysis
 - Treatment of fibromyalgia syndrome with gabapentin and pregabalin – A meta-analysis of randomized controlled trials, **145**, 69
- Mice
 - Direct excitation of spinal GABAergic interneurons by noradrenaline, **145**, 204
- Mice
 - Role of substance P signaling in enhanced nociceptive sensitization and local cytokine production after incision, **145**, 341
- Mice
 - The Rac GTPase-activating bacterial protein toxin CNF1 induces analgesia up-regulating μ -opioid receptors, **145**, 219
- Mouse
 - The effect of Spinal Cord Stimulation in mice with chronic neuropathic pain after partial ligation of the sciatic nerve, **145**, 312
- Multiple physical complaints
 - The unbearable lightness of somatisation: A systematic review of the concept of somatisation in empirical studies of pain, **145**, 31
- Muscle pain
 - Activation of the prostaglandin system in response to sleep loss in healthy humans: Potential mediator of increased spontaneous pain, **145**, 136
- Neck pain
 - Pain, perceived injustice and the persistence of post-traumatic stress symptoms during the course of rehabilitation for whiplash injuries, **145**, 325
- Negative thoughts
 - An assessment of the relative influence of pain coping, negative thoughts about pain, and pain acceptance on health-related quality of life among people with hemophilia, **145**, 169
- Nerve growth factor
 - Role of substance P signaling in enhanced nociceptive sensitization and local cytokine production after incision, **145**, 341
- Nerve injury
 - Prefrontal cortex and spinal cord mediated anti-neuropathy and analgesia induced by sarcosine, a glycine-T1 transporter inhibitor, **145**, 176
- Neurokinin receptor
 - Role of substance P signaling in enhanced nociceptive sensitization and local cytokine production after incision, **145**, 341
- Neuropathic pain
 - Differential involvement of A-delta and A-beta fibres in neuropathic pain related to carpal tunnel syndrome, **145**, 105
- Neuropathic pain
 - Sigma-1 receptors regulate activity-induced spinal sensitization and neuropathic pain after peripheral nerve injury, **145**, 294
- NMDA receptor
 - Ketamine and chronic pain – Going the distance, **145**, 271
- NMDA receptor
 - Ketamine produces effective and long-term pain relief in patients with Complex Regional Pain Syndrome Type 1, **145**, 304
- Nociceptive flexion reflex
 - Using normalized EMG to define the nociceptive flexion reflex (NFR) threshold: Further evaluation of standardized NFR scoring criteria, **145**, 211
- Nondermatomal somatosensory deficits
 - Nondermatomal somatosensory deficits in patients with chronic pain disorder: Clinical findings and hypometabolic pattern in FDG-PET, **145**, 252
- Noradrenaline
 - Direct excitation of spinal GABAergic interneurons by noradrenaline, **145**, 204
- Noxious
 - An fMRI case report of photophobia: Activation of the trigeminal nociceptive pathway, **145**, 358
- Noxious electrical stimulation
 - fMRI investigation of the effect of local and systemic lidocaine on noxious electrical stimulation-induced activation in spinal cord, **145**, 110
- Ophthalmic
 - An fMRI case report of photophobia: Activation of the trigeminal nociceptive pathway, **145**, 358
- Opioid
 - Electroacupuncture suppresses capsaicin-induced secondary hyperalgesia through an endogenous spinal opioid mechanism, **145**, 332
- μ -Opioid receptors
 - The Rac GTPase-activating bacterial protein toxin CNF1 induces analgesia up-regulating μ -opioid receptors, **145**, 219
- Oxidative stress
 - Tocotrienol ameliorates behavioral and biochemical alterations in the rat model of alcoholic neuropathy, **145**, 129
- Pain
 - In vivo* silencing of the $\text{Ca}_v3.2$ T-type calcium channels in sensory neurons alleviates hyperalgesia in rats with streptozocin-induced diabetic neuropathy, **145**, 184
- Pain
 - A prospective sequential analysis of the fear-avoidance model of pain, **145**, 45
- Pain
 - Activation of the prostaglandin system in response to sleep loss in healthy humans: Potential mediator of increased spontaneous pain, **145**, 136
- Pain
 - An fMRI case report of photophobia: Activation of the trigeminal nociceptive pathway, **145**, 358
- Pain
 - Direct excitation of spinal GABAergic interneurons by noradrenaline, **145**, 204
- Pain
 - Functional and structural nerve fiber findings in heterozygote patients with Fabry disease, **145**, 237
- Pain
 - Pain site frequency and location in sickle cell disease: The PiSCES project, **145**, 246

- Pain
 - Psychodynamic interpersonal therapy and improvement in interpersonal difficulties in people with severe irritable bowel syndrome, **145**, 196
- Pain
 - The Rac GTPase-activating bacterial protein toxin CNF1 induces analgesia up-regulating μ -opioid receptors, **145**, 219
- Pain
 - The unbearable lightness of somatisation: A systematic review of the concept of somatisation in empirical studies of pain, **145**, 31
- Pain acceptance
 - An assessment of the relative influence of pain coping, negative thoughts about pain, and pain acceptance on health-related quality of life among people with hemophilia, **145**, 169
- Pain communication
 - The smile of pain, **145**, 273
- Pain coping
 - An assessment of the relative influence of pain coping, negative thoughts about pain, and pain acceptance on health-related quality of life among people with hemophilia, **145**, 169
- Pain language
 - Pain language and gender differences when describing a past pain event, **145**, 86
- Pain location
 - Pain site frequency and location in sickle cell disease: The PiSCES project, **145**, 246
- Pain management
 - Medical students retain pain assessment and management skills long after an experiential curriculum: A controlled study, **145**, 319
- Pain threshold
 - Gender role expectations of pain is associated with pain tolerance limit but not with pain threshold, **145**, 230
- Pain tolerance
 - Gender role expectations of pain is associated with pain tolerance limit but not with pain threshold, **145**, 230
- Pain-related behavior
 - Experimental arthritis causes tumor necrosis factor- α -dependent infiltration of macrophages into rat dorsal root ganglia which correlates with pain-related behavior, **145**, 151
- Pain-related behavior
 - Gait abnormalities differentially indicate pain or structural joint damage in monoarticular antigen-induced arthritis, **145**, 142
- Patch-clamp
 - Direct excitation of spinal GABAergic interneurons by noradrenaline, **145**, 204
- Perceived injustice
 - Pain, perceived injustice and the persistence of post-traumatic stress symptoms during the course of rehabilitation for whiplash injuries, **145**, 325
- Performance-based assessment
 - Medical students retain pain assessment and management skills long after an experiential curriculum: A controlled study, **145**, 319
- Peripheral neuropathy
 - In vivo* silencing of the $Ca_v3.2$ T-type calcium channels in sensory neurons alleviates hyperalgesia in rats with streptozocin-induced diabetic neuropathy, **145**, 184
- Phenylephrine
 - Direct excitation of spinal GABAergic interneurons by noradrenaline, **145**, 204
- Placebo
 - Placebo conditioning and placebo analgesia modulate a common brain network during pain anticipation and perception, **145**, 24
- Placebo analgesia
 - Factors contributing to large analgesic effects in placebo mechanism studies conducted between 2002 and 2007, **145**, 36
- Placebo analgesia
 - Placebo conditioning and placebo analgesia modulate a common brain network during pain anticipation and perception, **145**, 24
- Post-traumatic stress symptoms
 - Pain, perceived injustice and the persistence of post-traumatic stress symptoms during the course of rehabilitation for whiplash injuries, **145**, 325
- Postoperative pain
 - A prospective comparison of post-surgical behavioral pain scales in preschoolers highlighting the risk of false evaluations, **145**, 60
- Prazosin
 - Direct excitation of spinal GABAergic interneurons by noradrenaline, **145**, 204
- Pregabalin
 - Treatment of fibromyalgia syndrome with gabapentin and pregabalin – A meta-analysis of randomized controlled trials, **145**, 69
- Prescription opioid episodes
 - Trends in prescribed opioid therapy for non-cancer pain for individuals with prior substance use disorders, **145**, 287
- Pressure
 - Enhanced central pain processing of fibromyalgia patients is maintained by muscle afferent input: A randomized, double-blind, placebo-controlled study, **145**, 96
- Prostaglandin
 - Activation of the prostaglandin system in response to sleep loss in healthy humans: Potential mediator of increased spontaneous pain, **145**, 136
- Provocation discography
 - A randomized double-blind controlled trial of intra-annular radiofrequency thermal disc therapy – A 12-month follow-up, **145**, 279
- Psychodynamic interpersonal psychotherapy
 - Psychodynamic interpersonal therapy and improvement in interpersonal difficulties in people with severe irritable bowel syndrome, **145**, 196
- Psychometrics
 - A prospective comparison of post-surgical behavioral pain scales in preschoolers highlighting the risk of false evaluations, **145**, 60
- Psychopathology
 - Is there an association between psychological factors and the Complex Regional Pain Syndrome type 1 (CRPS1) in adults? A systematic review, **145**, 52
- Psychophysics
 - Women experience greater heat pain adaptation and habituation than men, **145**, 350
- Quality of life
 - An assessment of the relative influence of pain coping, negative thoughts about pain, and pain acceptance on health-related quality of life among people with hemophilia, **145**, 169
- Quantitative sensory testing
 - Functional and structural nerve fiber findings in heterozygote patients with Fabry disease, **145**, 237
- Randomized controlled trial
 - A randomized double-blind controlled trial of intra-annular radiofrequency thermal disc therapy – A 12-month follow-up, **145**, 279
- Randomized controlled trial
 - Treatment of fibromyalgia syndrome with gabapentin and pregabalin – A meta-analysis of randomized controlled trials, **145**, 69
- Rat
 - Experimental arthritis causes tumor necrosis factor- α -dependent infiltration of macrophages into rat dorsal root ganglia which correlates with pain-related behavior, **145**, 151
- Receiver operating characteristics
 - Using normalized EMG to define the nociceptive flexion reflex (NFR) threshold: Further evaluation of standardized NFR scoring criteria, **145**, 211

Rho GTPases

The Rac GTPase-activating bacterial protein toxin CNF1 induces analgesia up-regulating μ -opioid receptors, **145**, 219

RIII

Using normalized EMG to define the nociceptive flexion reflex (NFR) threshold: Further evaluation of standardized NFR scoring criteria, **145**, 211

Sequential analysis

A prospective sequential analysis of the fear-avoidance model of pain, **145**, 45

Sex

Gender role expectations of pain is associated with pain tolerance limit but not with pain threshold, **145**, 230

Sex difference

Sex differences in functional brain activation during noxious visceral stimulation in rats, **145**, 120

Sex differences

Women experience greater heat pain adaptation and habituation than men, **145**, 350

Sharp

Women experience greater heat pain adaptation and habituation than men, **145**, 350

Sickle cell disease

Pain site frequency and location in sickle cell disease: The PiSCES project, **145**, 246

Sigma-1 receptor

Sigma-1 receptors regulate activity-induced spinal sensitization and neuropathic pain after peripheral nerve injury, **145**, 294

Skin

Role of substance P signaling in enhanced nociceptive sensitization and local cytokine production after incision, **145**, 341

Skin biopsy

Functional and structural nerve fiber findings in heterozygote patients with Fabry disease, **145**, 237

Sleep deprivation

Activation of the prostaglandin system in response to sleep loss in healthy humans: Potential mediator of increased spontaneous pain, **145**, 136

Smile

The smile of pain, **145**, 273

Somatisation

The unbearable lightness of somatisation: A systematic review of the concept of somatisation in empirical studies of pain, **145**, 31

Spinal cord

fMRI investigation of the effect of local and systemic lidocaine on noxious electrical stimulation-induced activation in spinal cord, **145**, 110

Spinal cord

Sigma-1 receptors regulate activity-induced spinal sensitization and neuropathic pain after peripheral nerve injury, **145**, 294

Spinal Cord Stimulation

The effect of Spinal Cord Stimulation in mice with chronic neuropathic pain after partial ligation of the sciatic nerve, **145**, 312

Spinal dorsal horn

Direct excitation of spinal GABAergic interneurons by noradrenaline, **145**, 204

Spine

Pain relief through expectation supersedes descending inhibitory deficits in fibromyalgia patients, **145**, 18

Standard nerve conduction study

Differential involvement of A-delta and A-beta fibres in neuropathic pain related to carpal tunnel syndrome, **145**, 105

Substance P

Role of substance P signaling in enhanced nociceptive sensitization and local cytokine production after incision, **145**, 341

Substance use disorders

Trends in prescribed opioid therapy for non-cancer pain for individuals with prior substance use disorders, **145**, 287

Suggestions

Factors contributing to large analgesic effects in placebo mechanism studies conducted between 2002 and 2007, **145**, 36

Systematic review

Is there an association between psychological factors and the Complex Regional Pain Syndrome type 1 (CRPS1) in adults? A systematic review, **145**, 52

Systematic review

The unbearable lightness of somatisation: A systematic review of the concept of somatisation in empirical studies of pain, **145**, 31

Systematic review

Treatment of fibromyalgia syndrome with gabapentin and pregabalin – A meta-analysis of randomized controlled trials, **145**, 69

Tactile sensitivity

Prefrontal cortex and spinal cord mediated anti-neuropathy and analgesia induced by sarcosine, a glycine-T1 transporter inhibitor, **145**, 176

Temporomandibular dysfunction

Momentary pain and coping in temporomandibular disorder pain: Exploring mechanisms of cognitive behavioral treatment for chronic pain, **145**, 160

Tender point

Enhanced central pain processing of fibromyalgia patients is maintained by muscle afferent input: A randomized, double-blind, placebo-controlled study, **145**, 96

Threshold

Using normalized EMG to define the nociceptive flexion reflex (NFR) threshold: Further evaluation of standardized NFR scoring criteria, **145**, 211

Tocotrienol

Tocotrienol ameliorates behavioral and biochemical alterations in the rat model of alcoholic neuropathy, **145**, 129

Tonic

Enhanced central pain processing of fibromyalgia patients is maintained by muscle afferent input: A randomized, double-blind, placebo-controlled study, **145**, 96

Tonic pain

Enhanced central pain processing of fibromyalgia patients is maintained by muscle afferent input: A randomized, double-blind, placebo-controlled study, **145**, 96

Trigeminal ganglion

An fMRI case report of photophobia: Activation of the trigeminal nociceptive pathway, **145**, 358

Trigeminal nucleus

An fMRI case report of photophobia: Activation of the trigeminal nociceptive pathway, **145**, 358

Tumor necrosis factor- α

Experimental arthritis causes tumor necrosis factor- α -dependent infiltration of macrophages into rat dorsal root ganglia which correlates with pain-related behavior, **145**, 151

Unpleasant

An fMRI case report of photophobia: Activation of the trigeminal nociceptive pathway, **145**, 358

Unpleasantness

An fMRI case report of photophobia: Activation of the trigeminal nociceptive pathway, **145**, 358

Uvea

An fMRI case report of photophobia: Activation of the trigeminal nociceptive pathway, **145**, 358

Ventromedial posterior thalamus

An fMRI case report of photophobia: Activation of the trigeminal nociceptive pathway, **145**, 358

Ventroposteromedial thalamus

An fMRI case report of photophobia: Activation of the trigeminal nociceptive pathway, **145**, 358

- Visceral pain
 - Sex differences in functional brain activation during noxious visceral stimulation in rats, **145**, 120
- Vitamin E
 - Tocotrienol ameliorates behavioral and biochemical alterations in the rat model of alcoholic neuropathy, **145**, 129
- Whiplash injury
 - Pain, perceived injustice and the persistence of post-traumatic stress symptoms during the course of rehabilitation for whiplash injuries, **145**, 325
- Wind-up
 - Sigma-1 receptors regulate activity-induced spinal sensitization and neuropathic pain after peripheral nerve injury, **145**, 294
- Withdrawal reflex
 - Pain relief through expectation supersedes descending inhibitory deficits in fibromyalgia patients, **145**, 18
- Work disability
 - A prospective sequential analysis of the fear-avoidance model of pain, **145**, 45