

DTQEM: A Phenomenological Model for Correlated Path-Information and Thermal Decoherence (v18.0-C)

Reddouane Berramdane^{1, *}

¹*Independent Researcher, Oujda, Morocco*
(Dated: June 12, 2026)

We present a phenomenological model for quantum decoherence in systems simultaneously subject to path-information extraction and thermal dephasing. The coherence function $C = C_0 \exp(-aI_{\text{path}} - b\Delta T/T_{\text{ref}} - cI_{\text{path}}\Delta T/T_{\text{ref}})$ extends the independent-bath baseline (v17.0-C, $c = 0$) by a bilinear cross-coupling term c , capturing the synergistic action of correlated environmental modes. Calibrated at $T_{\text{ref}} = 300$ K, the conservative ($c = 0$) limit achieves $R^2 = 0.9679$ and LOOCV $R^2 = 0.9356$; the full model reaches $R^2 = 0.9982$ (LOOCV = 0.9814) on the same synthetic dataset. Re-fitted to the eight-point dataset of Hornberger et al. (2003), the functional form yields $R^2 = 0.936$, demonstrating compatibility with molecule-interferometry data. A second-order cumulant analysis suggests strong inter-bath correlations, and an exploratory numerical scan reveals a thermal crossover at $T^* \approx 3.82$ K, flagged as a preliminary observation requiring low-temperature experimental confirmation. All code is openly available. A companion paper (Paper II) provides the full microscopic derivation.

I. INTRODUCTION

Quantum decoherence—the irreversible loss of phase coherence in quantum superpositions—limits every practical quantum technology [1, 2]. Two physically distinct channels often coexist: (i) which-path information leakage, reducing fringe visibility via the Englert–Greenberger–Yasin (EGY) relation [3]; and (ii) thermal dephasing from stochastic phase kicks.

In the simplest treatment, these channels add independently:

$$C_{\text{v17}} = C_0 \exp\left(-aI_{\text{path}} - b\frac{\Delta T}{T_{\text{ref}}}\right), \quad (1)$$

where $I_{\text{path}} \in [0, 1]$ is the EGY distinguishability. However, in molecular and solid-state platforms, the same phonon bath often mediates both channels, producing a cross term that Eq. (1) cannot capture. We formulate a minimal extension—DTQEM v18.0-C—adding a bilinear coupling $cI_{\text{path}}\Delta T/T_{\text{ref}}$, calibrate it, and test it against independent data. This paper focuses on the phenomenological model; a companion paper (Paper II) provides the full microscopic derivation from the Feynman–Vernon influence functional.

II. MODEL DEFINITION

A. Core equation

$$C(I_{\text{path}}, T) = C_0 \exp\left(-aI_{\text{path}} - b\frac{\Delta T}{T_{\text{ref}}} - cI_{\text{path}}\frac{\Delta T}{T_{\text{ref}}}\right), \quad (2)$$

with $\Delta T = T - T_{\text{ref}}$, $T_{\text{ref}} = 300$ K fixed, and four free parameters $\{C_0, a, b, c\}$. Setting $c = 0$ recovers Eq. (1).

B. Physical interpretation

- aI_{path} : EGY path-dephasing, consistent with Caldeira–Leggett weak coupling [2].
- $b\Delta T/T_{\text{ref}}$: linear thermal dephasing from the high- T Ohmic expansion $\coth(\hbar\omega/2k_B T) \approx 2k_B T/\hbar\omega$.
- $cI_{\text{path}}\Delta T/T_{\text{ref}}$: phenomenological cross-coupling for shared bath modes. Motivated by physical boundary conditions and a second-order cumulant argument (Appendix A); treated as a fit parameter.

C. Operational mapping of I_{path}

In circuit-QED, the mean intracavity photon number $\bar{n}$ gives [5]

$$I_{\text{path}}(\bar{n}) = \sqrt{1 - e^{-\kappa\bar{n}}}. \quad (3)$$

This is a testable proposal for future experiments; it is not used in the calibration below.

III. CALIBRATION

A. Dataset and procedure

Calibration used a synthetic grid $I_{\text{path}} \in [0, 1]$, $T \in [300, 550]$ K generated by the `dtqem_joint_bath_v18.py` simulator. Levenberg–Marquardt nonlinear least squares with leave-one-out cross-validation (LOOCV) was applied.

* reddoma@gmail.com

TABLE I. Calibrated parameters (DTQEM v18.0-C). Uncertainties: 1σ from the covariance matrix.

Parameter	Value	1σ	Role
C_0	0.3675	± 0.0052	Baseline coherence
a	1.6968	± 0.0245	Path-dephasing rate
b	0.8055	± 0.0148	Thermal-dephasing rate
c	0.5000	± 0.0201	Cross-coupling (empirical)
T_{ref}	300 K	fixed	Reference T

B. Performance metrics

TABLE II. Goodness-of-fit. LOOCV R^2 is the primary generalisation metric.

Model	R^2	LOOCV R^2	RMSE
v17.0-C ($c = 0$)	0.9679	0.9356	0.0202
v18.0-C (full)	0.9982	0.9814	0.0045

Note. Metrics computed on a synthetic dataset generated from the same functional form; LOOCV assesses overfitting, not experimental generalisation.

IV. VALIDATION ON HORNBERGER (2003) DATA

Hornberger et al. [4] reported C_{70} fullerene fringe visibility vs. collision rate. We extracted the eight published points, mapped collision rate to I_{path} via an auxiliary calibration, and re-fitted all four parameters independently. The re-fitted model achieves $R^2 = 0.936$.

The agreement demonstrates that Eq. (2) is compatible with independent molecule-interferometry data. It does not prove universality of the calibrated c . Statistical power of this $N = 8$ test is quantified in Appendix B.

V. EXPLORATORY TEMPERATURE SCAN: T^*

We define the effective thermal coefficient

$$b_{\text{eff}}(T) \equiv - \left. \frac{\partial \ln C}{\partial (\Delta T / T_{\text{ref}})} \right|_{I_{\text{path}}} \quad (4)$$

and evaluate it by numerically integrating the spin-boson dephasing functional with the Ohmic-Drude spectral density. Two regimes emerge (Fig. 1):

- $T > T^*$: $b_{\text{eff}} \propto T$ (classical Ohmic limit).
- $T < T^*$: b_{eff} saturates to $b_{\text{sat}} \approx 0.0103$, as $\coth(\hbar\omega_c/2k_B T) \rightarrow 1$.

The crossover temperature is

$$T^* = \frac{\hbar\omega_c}{2k_B} \approx 3.82 \text{ K} \quad (\omega_c = 10^{12} \text{ rad s}^{-1}). \quad (5)$$




FIG. 1. Effective thermal coefficient $b_{\text{eff}}(T)$. Blue dashed: classical extrapolation $b_0 T / T_{\text{ref}}$. Dotted: saturation $b_{\text{sat}} \approx 0.0103$. Dash-dotted: crossover $T^* \approx 3.82$ K.

Important caveat. Extrapolation below ~ 50 K violates the high- T Ohmic assumption underlying the $b\Delta T$ term. T^* is therefore a numerical observation from the chosen approximation, not a validated physical prediction. Experimental verification in the low-temperature regime is required.

VI. ASSUMPTIONS AND LIMITATIONS

1. **Markovian bath:** Memory effects are outside the model's scope.
2. **Pure dephasing dominance:** Longitudinal relaxation (T_1) is neglected.
3. **Room-temperature calibration:** The linear $b\Delta T$ term degrades below ~ 50 K.
4. **c is empirical:** Not derived microscopically in this work; specific to the calibration dataset. The full microscopic derivation is presented in Paper II.
5. **Small Hornberger dataset ($N = 8$):** $N \geq 36$ is required for reliable model selection (Appendix B).
6. **I_{path} mapping errors:** Propagate into all reported fits.
7. **T^* is exploratory:** The crossover temperature is a numerical observation under specific approximations, not a rigorous prediction.

VII. DISCUSSION

A. Contribution

DTQEM v18.0-C unifies two decoherence channels in one analytically tractable equation, predicts C with quantified uncertainty, and suggests a falsifiable crossover temperature T^* testable by measuring $\Gamma_\phi(T)/T$ over $0.5 \text{ K} \leq T \leq 100 \text{ K}$.

B. Relation to prior work

The bilinear form of Eq. (2) has not, to our knowledge, appeared in the open quantum systems literature. The closest antecedents—Hornberger [4] and Caldeira–Leggett [2]—both implicitly assume $c = 0$.

VIII. OUTLOOK

- **Microscopic derivation (Paper II):** The companion paper derives the cross term via the Feynman–Vernon influence functional for two correlated baths. The exact form is $\propto 2\rho\sqrt{ab}\sqrt{I_{\text{path}} \cdot \Delta T/T_{\text{ref}}}$, of which Eq. (2) is the leading-order polynomial approximation satisfying physical boundary conditions.
- **Low-temperature experimental test:** Confirm or refute the existence of a crossover near T^* using cold-atom or superconducting qubit platforms.
- **Larger validation datasets:** $N \geq 36$ configurations (fullerene beams or transmon qubits) to robustly identify c .

IX. CONCLUSIONS

DTQEM v18.0-C provides a four-parameter phenomenological model for joint-bath coherence loss. Key results:

- Conservative calibration: $R^2 = 0.9679$, LOOCV $R^2 = 0.9356$ ($c = 0$ limit).
- Independent validation: $R^2 = 0.936$ on Hornberger (2003), $N = 8$.
- Cumulant analysis suggests strong inter-bath correlation.
- Numerical crossover $T^* \approx 3.82$ K, flagged as exploratory and requiring experimental verification.

A companion paper (Paper II) deriving the exact cross-coupling from first principles is available alongside this work.

ACKNOWLEDGMENTS

The author thanks the open-source communities behind NumPy, SciPy, and Matplotlib. AI language models

(DeepSeek, Claude, and Qwen) were used as analytical and editorial tools under full human scientific supervision. Their contributions included critical analysis, code implementation, theoretical derivations, and manuscript organization. The author independently verified all derivations, numerical results, and physical claims; all scientific content and conclusions are the sole responsibility of the author. Code: <https://github.com/reddoma742/Davisson-Germer-DTQEM>. Archive: Zenodo [7].

Appendix A: Motivation for the Cross-Coupling Term

Consider a two-level system coupled to a path-measurement bath (operator $\hat{B}_1$) and a thermal bath ($\hat{B}_2$), with total bath Hamiltonian $\hat{H}_B = \hat{H}_{B_1} + \hat{H}_{B_2} + \hat{H}_{B_1 B_2}$. A second-order cumulant expansion yields

$$\Phi_{\text{total}} = \Phi_1 + \Phi_2 + 2\rho\sqrt{\Phi_1\Phi_2}, \quad (\text{A1})$$

where $\rho \in [0, 1]$ is the normalized cross-correlation coefficient. With $\Phi_1 \propto I_{\text{path}} \equiv u$ and $\Phi_2 \propto \Delta T/T_{\text{ref}} \equiv v$, the exact microscopic cross term is proportional to $\sqrt{uv}$.

Expanding $\sqrt{uv}$ around a nominal operating point (u_0, v_0) gives a bilinear term. Matching the coefficient of uv yields $c = \rho\sqrt{ab}/(2\sqrt{u_0v_0})$. Using the calibrated a, b, c from Table I with representative values $u_0 = v_0 = 0.5$ suggests a cross-correlation coefficient of order unity ($\rho \sim 1$), indicating strong inter-bath coupling. The estimate is sensitive to the choice of (u_0, v_0) ; the precise numerical value is presented in Paper II, which provides the full microscopic derivation and the rigorous expression for ρ .

Appendix B: AICc Model Selection Threshold

TABLE III. AICc model-selection probability ($c = 0.5$, $\sigma = 0.005$, 10^4 simulations per row). Reference: [6].

N	v17.0-C (%)	v18.0-C (%)	Reliability
8	89	11	Low
16	57	43	Low
36	21	79	Moderate
50	10	90	High
80	5	95	Very high

[1] H.-P. Breuer and F. Petruccione, *The Theory of Open Quantum Systems* (Oxford University Press, Oxford, 2002).

[2] U. Weiss, *Quantum Dissipative Systems*, 4th ed. (World Scientific, Singapore, 2012).

- [3] B.-G. Englert, Phys. Rev. Lett. **77**, 2154 (1996).
- [4] K. Hornberger, S. Uttenthaler, B. Brezger, L. Hackermüller, M. Arndt, and A. Zeilinger, Phys. Rev. Lett. **90**, 160401 (2003).
- [5] A. Blais, A. L. Grimsmo, S. M. Girvin, and A. Wallraff, Rev. Mod. Phys. **93**, 025005 (2021).
- [6] K. P. Burnham and D. R. Anderson, *Model Selection and Multimodel Inference*, 2nd ed. (Springer, New York, 2002).
- [7] R. Berramdane, DTQEM: A Phenomenological Model for Correlated Path-Information and Thermal Decoherence (v18.0-C), Zenodo (2026), [DOI:10.5281/zenodo.20670032](https://doi.org/10.5281/zenodo.20670032).