

RV COLLEGE OF ENGINEERING
(An Autonomous Institute Affiliated to VTU, Bangalore)
DEPARTMENT OF MATHEMATICS
FUNDAMENTALS OF LINEAR ALGEBRA, CALCULUS AND NUMERICAL METHODS (MATH214T)
Multivariable Functions and Partial Differentiation

TUTORIAL SHEET 1

1. If $z = r \cos \theta$, $y = r \sin \theta$, then $\left(\frac{\partial z}{\partial r}\right)^2 + \left(\frac{\partial z}{\partial \theta}\right)^2 = \dots$ Ans: r^2
2. If $z = x \sin y + y \sin x$, then $\frac{\partial^2 z}{\partial x \partial y} = \dots$ Ans: $\cos x + \cos y$
3. If $z = e^{x^2+y^2}$, then $\frac{\partial^2 z}{\partial x^2} = \dots$ Ans: $2x^2 e^{x^2+y^2}$
4. The steady state temperature of a metal sheet is $T(x, y) = x^3 - a^2 y^3$. The values of 'a' for which $T(x, y)$ satisfies the Laplace equation $T_{xx} + T_{yy} = 0$ are \dots Ans: ± 1
5. If $u = y \cos(xy)$ then $\frac{\partial^2 u}{\partial x^2}$ at the point $(1, \pi)$ is \dots Ans: -1
6. If $V = f(x-ct) + g(x+ct)$ where f and g are arbitrary functions of $x-ct$ and $x+ct$ respectively and c is a constant, then show that $\frac{\partial^2 V}{\partial x^2} = c^2 \frac{\partial^2 V}{\partial t^2}$
7. If $u = ae^{ax} \sin(ax - g \cdot x)$, where a, g and n are positive constants and $\frac{\partial^2 u}{\partial x^2} = \mu \frac{\partial^2 u}{\partial t^2}$, show that $g = \frac{a}{\mu}$
8. If V is the volume and S is the total surface area of rectangular box of length x , breadth y and height z , find
 - (i) the rate of change of V with respect to x if $y=4$ and $z=12$.
 - (ii) the rate of change of S with respect to x if $y=3$ and $z=4$.
9. If $u = \log(x^2 + y^2 + z^2 - 3xyz)$, then show that $\left(\frac{\partial u}{\partial x}\right)^2 + \left(\frac{\partial u}{\partial y}\right)^2 + \left(\frac{\partial u}{\partial z}\right)^2 = -\frac{9}{(x+y+z)^2}$

$$a^3 + b^3 + c^3 = (a+b+c)(a^2 + b^2 + c^2 - ab - bc - ca)$$

2 | Page

RV COLLEGE OF ENGINEERING
(An Autonomous Institute Affiliated to VTU, Bangalore)
DEPARTMENT OF MATHEMATICS
FUNDAMENTALS OF LINEAR ALGEBRA, CALCULUS AND NUMERICAL METHODS (MATH214T)
Multivariable Functions and Partial Differentiation

TUTORIAL SHEET 2

1. Given $z = xy^2 + x^2 y$ where x and y are functions of t with $x(1) = 1, y(1) = 2, x'(1) = 3$ and $y'(1) = 4$. The value of $\frac{dz}{dt}$ at $t=1$ is \dots Ans: 16
2. For the implicit function $\cos(x^2) = (\sin y)^3$, $\frac{\partial^2 z}{\partial x^2} = \dots$
3. Given $u = x^2 y^2$ represents time and $u = x^2 - y^2, x = \frac{1}{t}, y = \frac{1}{t}$ then the rate of change of u with respect to t is \dots Ans: $-\frac{2}{t^3}$
4. For the implicit function $e^x - e^y = 2xy$, $\frac{\partial^2 u}{\partial x^2} = \dots$ Ans: $\frac{e^x - 2y}{(e^x - 2y)^2}$
5. Given $x^2 + y^2 + 3z^2 = 1$ and $z = y + x$, then $\frac{dz}{dx} = \dots$ Ans: $\frac{2x - y}{2}$
6. If $u = x^2 y^2 z^2$, $x = e^t, y = e^t, z = e^t$ then $\frac{du}{dt} = \dots$ Ans: $6e^{3t}$
7. If $u = xyz$ where $x = e^t, y = e^t, z = e^t$ then $\frac{du}{dt} = \dots$ Ans: $3e^{3t}$
8. For the implicit function $e^x - e^y = 2xy$, find $\frac{\partial^2 u}{\partial x^2} = \dots$ Ans: $2(x + e^x) + 2(x + 4e^{3x})3e^{3x}$
9. If z is a function of x and y and if $z = e^x \sin y, y = e^x \cos x$, prove that $\frac{\partial^2 z}{\partial x^2} + \frac{\partial^2 z}{\partial y^2} = 0$
10. If $u = x^2 \tan^{-1}\left(\frac{y}{x}\right) - y^2 \tan^{-1}\left(\frac{x}{y}\right)$, determine $\frac{\partial^2 u}{\partial x^2} = \dots$

2 | Page

RV COLLEGE OF ENGINEERING
(An Autonomous Institute Affiliated to VTU, Bangalore)
DEPARTMENT OF MATHEMATICS
FUNDAMENTALS OF LINEAR ALGEBRA, CALCULUS AND NUMERICAL METHODS (MATH214T)
Multivariable Functions and Partial Differentiation

TUTORIAL SHEET 3

1. Match the following:

(i) If $z = e^{x^2+y^2}, y = e^x \cos x$ then $\frac{\partial z}{\partial x} = \dots$	a) $\frac{1}{4\sqrt{2}\ln 2}$
(ii) If (a,b) is a stationary point of $f(x,y)$ and $f_{xx}(a,b) = 2$ and $f_{yy}(a,b) = 2$ at this point then the nature of (a,b) is \dots	b) minimum
(iii) The nature of the point $(1, -1)$ to the function $f(x,y) = x^2 - 3xy^2 + y^4$ is \dots	c) $e^{2x^2}(\sin^2 y - \sin^4 y)$
(iv) If $\frac{\partial^2 f}{\partial x^2} = \sin 2x$ and $\frac{\partial^2 f}{\partial y^2} = \frac{1}{4}$ then $\frac{\partial^2 f}{\partial x^2} = \dots$	d) $\sin 2x$
2. Find the extreme values of $\sin x + \sin y + \sin(x+y)$. Ans: Maximum value at $\left(\frac{\pi}{2}, \frac{\pi}{2}\right)$
3. Find the volume of largest rectangular parallelepiped that can be inscribed in an ellipsoid $\frac{x^2}{a^2} + \frac{y^2}{b^2} + \frac{z^2}{c^2} = 1$. Ans: Maximum volume $\frac{8abc}{3}$
4. Find the maximum and minimum distances of the point $(1, 2, 3)$ from the sphere $x^2 + y^2 + z^2 = 56$ using Lagrange's Method of undetermined multipliers. Ans: Minimum distance at $(2, 4, 6)$; Maximum distance at $(-2, -4, -6)$
5. Show that $u = \frac{x^2+y^2+z^2}{x^2+y^2+z^2}$ are functionally dependent and find the relation between them. Ans: $x^2 + y^2 + z^2 = 1$
6. For $u = xyz, x = yz + xz, y = xz + xy, z = xy + xz$, find $\frac{\partial u}{\partial x} = \dots$ Ans: $(y-z)(x-z)(x-y)$
7. If $u = e^x \sec u, y = e^x \tan u$, then verify that $f'' = 1$.

2 | Page

$$\textcircled{1} y = 8 \sin \theta ; \left(\frac{dy}{d\theta}\right)^2 + \left(\frac{dx}{d\theta}\right)^2 = 64$$

$$\frac{dy}{d\theta} = 8 \cos \theta = 8^2 \cos^2 \theta + 8^2 \sin^2 \theta = 64$$

$$x = 8 \cos \theta \Rightarrow \frac{dx}{d\theta} = -8 \sin \theta$$

$$\textcircled{2} z = x \sin y + y \sin x$$

$$\frac{\partial}{\partial x} \left(\frac{\partial z}{\partial y} \right) = \frac{\partial}{\partial x} (x \cos y + \sin x)$$

$$= \cos y + \cos x$$

$$\textcircled{3} u = y \cos(xy)$$

$$\frac{\partial u}{\partial y} = (1) \cos(xy) + y(\sin(xy)) \cdot x$$

$$\frac{\partial^2 u}{\partial x^2} = -y \sin(xy) = -1$$

$$\textcircled{6} v = f(x-ct) + g(x+ct)$$

$$\frac{dv}{dt} = f'(x-ct) \cdot (-c) + g'(x+ct) \cdot c$$

$$\frac{d^2 v}{dt^2} = c^2 f''(x-ct) + c^2 g''(x+ct)$$

$$\Rightarrow \frac{d^2 v}{dt^2} = c^2 \frac{d^2 v}{dx^2}$$

$$\textcircled{8} u = ae^{-gx} \sin(nt-gx)$$

$$\frac{\partial u}{\partial t} = ae^{-gx} \cos(nt-gx) \cdot n$$

$$\frac{\partial^2 u}{\partial x^2} = \frac{\partial}{\partial x} \left(a(-e^{-gx}) \cos(nt-gx) \cdot (-g) \right)$$

$$= \frac{\partial}{\partial x} \left(-age^{-gx} (\cos(nt-gx) + \sin(nt-gx)) \right)$$

$$= -ag \left[-e^{-gx} (-g) (\cos X + \sin X) + e^{-gx} (-\sin X (-g) + \cos X (-g)) \right]$$

$$= ag^2 e^{-gx} (\cos X + \sin X + \sin X + \cos X)$$

$$= 2ag^2 e^{-gx} \cos(nt-gx)$$

$$\frac{du}{dt} = M \frac{d^2 u}{dx^2}$$

$$\frac{d}{dx} \left(\frac{1}{e^{-gx}} \cos(nt-gx) \right) \cdot n = M \cdot \frac{2}{e^{-gx}} \cos(nt-gx)$$

$$n = M(2g^2)$$

$$g = \sqrt{\frac{n}{2M}}$$

$$\text{III } z = xy^2 + x^2 y \quad z = f_1 + f_2$$

$$\frac{dz}{dt} = \frac{\partial z}{\partial x} \frac{dx}{dt} + \frac{\partial z}{\partial y} \frac{dy}{dt}$$

$$= (y^2 + 2xy^2)(x'(t)) + (2xy)(y'(t))$$

$$at \ t=1, x=1, y=2, x'(1)=3, y'(1)=4$$

$$\frac{dz}{dt} = (2^2 + 3(1)^2(2))(3) + (2(1)(2))(4)$$

$$= 30 + 20 = 50$$

$$\textcircled{2} (\cos x)^y = (\sin y)^x \rightarrow \text{implicit}$$

$$b = (\cos x)^y = (\sin y)^x$$

$$\frac{dy}{dx} = \frac{-fx}{fy} \quad \langle a^x = a^x \log a \rangle$$

$$= - \frac{y(\cos x)^{y-1} x(-\sin x) - \sin y^x \log(\sin y)}{(\cos x)^y \log(\cos x) - x(\sin y)^{x-1} \cos y}$$

$$\textcircled{3} u = x^2 - y^2 ; x = \frac{1}{t}, y = e^t$$

$$\textcircled{2} z = e^{2x^2+xy}$$

$$\frac{dz}{dy} = xe^{2x^2+xy}$$

$$f(x,y) = x^2 - a^2 y^2 = y$$

$$fx + fy = 0$$

$$\frac{dx}{dy} + \frac{dy}{dy} = 0$$

$$\frac{dx}{dy} + 1 = 0$$

$$2 - 2a^2 = 0$$

$$a^2 = 1 \Rightarrow a = \pm 1$$

$$\textcircled{7} u = \frac{x+y}{x-y}$$

$$\frac{d^2 u}{dx dy} = \frac{d}{dx} \left(\frac{du}{dy} \right)$$

$$= \frac{d}{dx} \left(\frac{(1)(x-y) - (x+y)(-1)}{(x-y)^2} \right)$$

$$= \frac{d}{dx} \left(\frac{x-y+x+y}{(x-y)^2} \right)$$

$$= \frac{d}{dx} \left(\frac{2x}{(x-y)^2} \right)$$

$$= 2 \left(\frac{(1)(x-y)^2 - x(2(x-y)(-1))}{(x-y)^4} \right)$$

$$= 2 \left(\frac{x^2 + y^2 - 2xy - 2x^2 + 2xy}{(x-y)^4} \right)$$

$$= 2 \left(\frac{y^2 - x^2}{(x-y)^4} \right) = 2 \left(\frac{y^2 - x^2}{(x-y)^4} \right) = 2 \left(\frac{y^2 - x^2}{(x-y)^4} \right)$$

$$\textcircled{9} V = xyz \quad S = 2(xy + yz + xz)$$

$$(i) \frac{\partial V}{\partial x} \bigg|_{y=4, z=12} = yz = 4 \times 12 = 48$$

$$(ii) \frac{\partial S}{\partial z} \bigg|_{x=3, y=4} = 2(y+x) = 2(4+3) = 14$$

$$\textcircled{10} M = \log(x^3 + y^3 + z^3 - 3xyz)$$

$$\left(\frac{\partial}{\partial x} + \frac{\partial}{\partial y} + \frac{\partial}{\partial z} \right)^2 M = \frac{-9}{(x+y+z)^2}$$

$$\rightarrow \left(\frac{\partial}{\partial x} + \frac{\partial}{\partial y} + \frac{\partial}{\partial z} \right) \left(\frac{1}{(x+y+z)^2} \right) \left(\frac{\partial u}{\partial x} + \frac{\partial u}{\partial y} + \frac{\partial u}{\partial z} \right)$$

$$\rightarrow \frac{\partial u}{\partial x} = \frac{1}{x^3 + y^3 + z^3 - 3xyz} \times (3x^2 - 3yz)$$

$$\rightarrow \frac{\partial u}{\partial y} = \frac{1}{x^3 + y^3 + z^3 - 3xyz} \times (3y^2 - 3xz)$$

$$\rightarrow \frac{\partial u}{\partial z} = \frac{1}{x^3 + y^3 + z^3 - 3xyz} \times (3z^2 - 3xy)$$

$$\left(\frac{\partial}{\partial x} + \frac{\partial}{\partial y} + \frac{\partial}{\partial z} \right) \left(\frac{1}{x^3 + y^3 + z^3 - 3xyz} \right) \left(\frac{3x^2 + y^2 + z^2 - xy - yz - xz}{x^3 + y^3 + z^3 - 3xyz} \right)$$

$$\left(\frac{3(11)}{(x+y+z)(1)} \right)$$

$$= \left(\frac{\partial}{\partial x} + \frac{\partial}{\partial y} + \frac{\partial}{\partial z} \right) \left(\frac{3}{x+y+z} \right)$$

$$= \frac{\partial}{\partial x} \left(\frac{3}{x+y+z} \right) + \dots$$

$$= 3 \left(\frac{-1}{(x+y+z)^2} \right) + \dots$$

$$= -\frac{3}{(x+y+z)^2} + \frac{-3}{(x+y+z)^2} + \dots$$

$$= -\frac{9}{(x+y+z)^2}$$

$$\textcircled{4} e^x - e^y = 2xy$$

$$f: e^x - e^y - 2xy = 0$$

$$\frac{dy}{dx} = \frac{-fx}{fy} = \frac{-(e^x - 2y)}{-e^y - 2x}$$

$$= \frac{e^x - 2y}{e^y + 2x}$$

$$\textcircled{5} x^2 + y^2 + 3xz = 1$$

$$2x + 2y \frac{dy}{dx} + 3(x \frac{dz}{dx} + z) = 0$$

$$\frac{dx}{dx} + \frac{dy}{dx} = 0$$

$$du = -1$$

$$\begin{aligned} \text{[5]} \quad x^2 + y^2 + 3xz &= 1 \\ 2x + 2y \frac{dy}{dx} + 3(x \frac{dz}{dx} + z) &= 0 \\ 2x - 2y + 3x \frac{dz}{dx} + 3z &= 0 \\ 3x \frac{dz}{dx} &= 2y - 2x - 3z \\ \frac{dz}{dx} &= \frac{2y}{3x} - \frac{2}{3} - \frac{3z}{3x} \\ &= \frac{2y}{3x} - \frac{2}{3} - \frac{z}{x} \end{aligned}$$

$$\begin{aligned} \frac{x^2 + y^2}{x^2} + \frac{3xz}{x^2} &= \frac{1}{x^2} \\ 1 + \frac{y^2}{x^2} + \frac{3z}{x} &= \frac{1}{x^2} \\ 3\frac{z}{x} &= \frac{1}{x^2} - \frac{y^2}{x^2} - 1 \end{aligned}$$

$$\begin{aligned} -\frac{z}{x} &= \frac{1}{3} - \frac{(1-y^2)}{3x^2} \\ \frac{2y}{3x} - \frac{2}{3} - \frac{z}{x} &= \frac{1}{3} - \frac{(1-y^2)}{3x^2} \\ \frac{2y}{3x} - \frac{2}{3} - \frac{z}{x} &= \frac{1}{3} - \frac{(1-y^2)}{3x^2} \end{aligned}$$

$$\text{[9]} \quad z = \begin{cases} x = u \\ y = v \end{cases}$$

$$\begin{aligned} \text{PTs:} \\ \text{(i)} \quad \frac{\partial z}{\partial u} &= x \frac{\partial z}{\partial x} + y \frac{\partial z}{\partial y} \\ \rightarrow \frac{\partial z}{\partial u} &= \frac{\partial z}{\partial x} \cdot \frac{\partial x}{\partial u} + \frac{\partial z}{\partial y} \cdot \frac{\partial y}{\partial u} \\ &= \frac{\partial z}{\partial x} \cdot e^u \sin v + \frac{\partial z}{\partial y} \cdot e^u \cos v \\ \frac{\partial z}{\partial u} &= \frac{\partial z}{\partial x} x + \frac{\partial z}{\partial y} y \quad \text{proved} \end{aligned}$$

$$\begin{aligned} \text{ii)} \quad \frac{\partial z}{\partial x} &= e^{-u} \left(\sin v \frac{\partial z}{\partial u} + \cos v \frac{\partial z}{\partial v} \right) \\ \frac{\partial z}{\partial v} &= \frac{\partial z}{\partial x} \frac{\partial x}{\partial v} + \frac{\partial z}{\partial y} \frac{\partial y}{\partial v} \\ &= \frac{\partial z}{\partial x} \cdot e^u \cos v + \frac{\partial z}{\partial y} \cdot e^u (\sin v) \\ &= e^u \left(\frac{\partial z}{\partial x} \cos v + \frac{\partial z}{\partial y} \sin v \right) \\ \frac{\partial z}{\partial v} &= \frac{\partial z}{\partial x} y + \frac{\partial z}{\partial y} (x) \\ \frac{\partial z}{\partial u} \frac{\partial z}{\partial v} &= \frac{\partial z}{\partial x} (x+y) + \frac{\partial z}{\partial y} (y-x) \\ &= \frac{\partial z}{\partial x} = \text{LHS} \end{aligned}$$

$$\begin{aligned} \frac{dy}{dz} &= \frac{-fx}{fy} = -\frac{[e^x - 2y]}{-e^y - 2x} \\ &= \frac{e^x - 2y}{e^y + 2x} \end{aligned}$$

$$\begin{aligned} \text{[7]} \quad M &= xyz \\ x &= e^{-t}, y = e^t \sin t, z = \sin t \\ u &= \begin{cases} x \\ y \\ z \end{cases} t \end{aligned}$$

$$\begin{aligned} \frac{du}{dt} &= \frac{\partial u}{\partial x} \frac{\partial x}{\partial t} + \frac{\partial u}{\partial y} \frac{\partial y}{\partial t} + \frac{\partial u}{\partial z} \frac{\partial z}{\partial t} \\ &= yz(-e^{-t}) + xz(e^{-t} \sin t + e^{-t} (\sin t \cos t - \cos t)) \\ &= -e^{-t} \sin^2 t + e^{-t} \sin t (2 \cos t - \sin t) \\ &= -e^{-t} \sin^2 t + e^{-t} \sin t (2 \cos t - \sin t) \\ &= e^{-t} \sin^2 t (3 \cos t - 2 \sin t) \end{aligned}$$

$$\begin{aligned} \text{[10]} \quad u &= f(x-y, y-z, z-x) \\ r &= x-y, s = y-z, t = z-x \\ u &= \begin{cases} r \\ s \\ t \end{cases} \end{aligned}$$

$$\begin{aligned} \rightarrow \frac{\partial u}{\partial x} &= \frac{\partial u}{\partial r} \frac{\partial r}{\partial x} + \frac{\partial u}{\partial s} \frac{\partial s}{\partial x} + \frac{\partial u}{\partial t} \frac{\partial t}{\partial x} \\ &= \frac{\partial u}{\partial r} (1) + \frac{\partial u}{\partial s} (0) + \frac{\partial u}{\partial t} (-1) \end{aligned}$$

$$\begin{aligned} \frac{\partial u}{\partial x} &= \frac{\partial u}{\partial r} - \frac{\partial u}{\partial t} \\ \frac{\partial u}{\partial y} &= \frac{\partial u}{\partial r} \frac{\partial r}{\partial y} + \frac{\partial u}{\partial s} \frac{\partial s}{\partial y} + \frac{\partial u}{\partial t} \frac{\partial t}{\partial y} \\ &= \frac{\partial u}{\partial r} (-1) + \frac{\partial u}{\partial s} (1) + \frac{\partial u}{\partial t} (0) \end{aligned}$$

$$\begin{aligned} \frac{\partial u}{\partial y} &= \frac{\partial u}{\partial s} - \frac{\partial u}{\partial r} \\ \frac{\partial u}{\partial z} &= \frac{\partial u}{\partial r} \frac{\partial r}{\partial z} + \frac{\partial u}{\partial s} \frac{\partial s}{\partial z} + \frac{\partial u}{\partial t} \frac{\partial t}{\partial z} \\ &= \frac{\partial u}{\partial r} (0) + \frac{\partial u}{\partial s} (-1) + \frac{\partial u}{\partial t} (1) \end{aligned}$$

$$\begin{aligned} \frac{\partial u}{\partial z} &= \frac{\partial u}{\partial t} - \frac{\partial u}{\partial s} \\ \rightarrow \frac{\partial u}{\partial x} \frac{\partial u}{\partial y} + \frac{\partial u}{\partial y} \frac{\partial u}{\partial z} &= \frac{\partial u}{\partial r} \frac{\partial u}{\partial s} - \frac{\partial u}{\partial s} \frac{\partial u}{\partial r} + \frac{\partial u}{\partial s} \frac{\partial u}{\partial t} - \frac{\partial u}{\partial t} \frac{\partial u}{\partial s} \\ &= 0 \end{aligned}$$

$$\begin{aligned} \text{[8]} \quad Z &= x^2 + 2xy + 4y^2, y = e^{3x} \\ \frac{dz}{dx} &= 2x + 2(x \frac{dy}{dx} + y) + 4(2y) \frac{dy}{dx} \\ &= 2x + 2(x(3e^{3x}) + e^{3x}) + 8(e^{3x})(3e^{3x}) \\ &= 2x + 2e^{3x}(3x+1) + 24e^{6x} \\ &= 2x + 6xe^{3x} + 2e^{3x} + 24e^{6x} \\ &= 2(x + e^{3x}) + 6e^{3x}(x + 4e^{3x}) \end{aligned}$$

$$\text{[11]} \quad u = x^2 \tan^{-1}\left(\frac{y}{x}\right) - y^2 \tan^{-1}\left(\frac{x}{y}\right)$$

$$\begin{aligned} \frac{\partial u}{\partial x \partial y} &= \frac{\partial}{\partial x} \left(\frac{\partial u}{\partial y} \right) \\ &= \frac{\partial}{\partial x} \left(x^2 \cdot \frac{1}{1 + \frac{y^2}{x^2}} \cdot \left(\frac{1}{x} \right) - 2y \cdot \frac{1}{1 + \frac{x^2}{y^2}} \cdot \left(\frac{1}{y} \right) \right) \\ &= \frac{\partial}{\partial x} \left(\frac{x^3}{x^2 + y^2} + \frac{2xy}{x^2 + y^2} \right) \\ &= \frac{\partial}{\partial x} \left(\frac{x^3 + 2xy}{x^2 + y^2} \right) \\ &= \frac{(3x^2 + 2y)(x^2 + y^2) - (x^3 + 2xy)(2x + y)}{(x^2 + y^2)^2} \\ &= \frac{3x^4 + 3x^2y^2 + 2x^2y + 2y^3 - 2x^4 - x^3y^2 - 4x^2y^2 - 2xy^3}{(x^2 + y^2)^2} \\ &= \frac{x^4 + 3x^2y^2 - 2x^2y + 2y^3 - x^3y^2 - 2xy^3}{(x^2 + y^2)^2} \end{aligned}$$