



unit3_tutorial

RV COLLEGE OF ENGINEERING
(An Autonomous Institute Affiliated to VTU, Bangalore)
DEPARTMENT OF MATHEMATICS
FUNDAMENTALS OF LINEAR ALGEBRA, CALCULUS AND NUMERICAL METHODS (MATH1AT)
Multivariable Functions and Partial Differentiation

TUTORIAL SHEET 1

1. If $x = r \cos \theta$, $y = r \sin \theta$, then $\left(\frac{\partial}{\partial x}\right)^2 + \left(\frac{\partial}{\partial y}\right)^2 = \frac{\partial^2}{\partial r^2}$. Ans: ?
2. If $z = x \sin y + y \sin x$, then $\frac{\partial^2 z}{\partial x^2} = \sin y$. Ans: $\cos y + \sin x$
3. For steady state temperature of a semi disc $\nabla^2 u = 0$, $u = 0$ on $r = 1$. The value of u for which $\nabla u(x, y)$ satisfies the Laplace equation $\nabla^2 u = 0$ on $r = 1$ is $\frac{1}{2}$. Ans: ?
4. If $u = y \cos(xy)$ then $\frac{\partial^2 u}{\partial x^2}$ at the point $(1, 1)$ is -1 . Ans: -1
5. If $V = f(x - ct) + g(x + ct)$ where f and g are arbitrary functions of $x - ct$ and $x + ct$ respectively and c is a constant, then show that $\frac{\partial^2 V}{\partial x^2} = c^2 \frac{\partial^2 V}{\partial t^2}$.
6. If $u = \cos^{-1} \sin(xt - yz)$, where x, y and z are positive constants and $\frac{\partial^2 u}{\partial x^2} = \frac{\partial^2 u}{\partial y^2}$, show that $g = \frac{1}{2}$.
7. If V is the volume and S is the total surface area of rectangular box of length x , breadth y and height z , find
- (i) the rate of change of V with respect to x if $y = 4$ and $z = 12$.
- (ii) the rate of change of S with respect to x if $y = 3$ and $z = 4$.
- Ans: (i) 48, (ii) 44
8. If $u = \log(x^2 + y^2 + z^2 - 3xyz)$, then show that $\left(\frac{\partial}{\partial x} + \frac{\partial}{\partial y} + \frac{\partial}{\partial z}\right)u = -\frac{3}{(x+y+z)}$.

2 | Page

RV COLLEGE OF ENGINEERING
(An Autonomous Institute Affiliated to VTU, Bangalore)
DEPARTMENT OF MATHEMATICS
FUNDAMENTALS OF LINEAR ALGEBRA, CALCULUS AND NUMERICAL METHODS (MATH1AT)
Multivariable Functions and Partial Differentiation

TUTORIAL SHEET 2

1. Given $u = xy^2 + x^2y$ where x and y are functions of t with $x(1) = 1$, $y(1) = 2$, $x'(1) = 3$ and $y'(1) = 4$. The value of $\frac{du}{dt}$ at $t = 1$ is 70 . Ans: 16?
2. For the implicit function $\cos(y) = \sin(x)$, $\frac{dy}{dx} = \frac{1}{\cos(x)}$.
3. Given r represents time and $u = x^2 - y^2$, $x = e^t$, $y = e^{2t}$ then the rate of change of u with respect to t is $-2e^{3t}$. Ans: $\frac{e^{3t}-2e^{2t}}{e^{3t}+1}$
4. For the implicit function $e^x - e^y = 2xy$, $\frac{dy}{dx} = \frac{e^x - 2y}{e^y - 2x}$. Ans: $\frac{e^{2x}-2y}{e^{2y}-2x}$
5. Given, $e^x + y^2 + 3xz = 1$ and $x + y = 1$, then $\frac{dz}{dx} = \frac{1}{2}$. Ans: $\frac{1}{2}$
6. If $u = z(x, y)$, $x = e^t \sin t$, $y = e^t \cos t$, then $\frac{du}{dt} = e^t \sin t$.
7. If $u = xyz$ where $x = e^t$, $y = e^{2t} \sin t$, $z = \sin t$, then find $\frac{du}{dt}$.
8. If $u = e^x + y + 2xy + 4yz$ and $v = e^{2x}$, find $\frac{du}{dv}$. Ans: $2(x + e^{2x}) + 2(x + 4e^{2x})e^{2x}$
9. If u is a function of x and y and if $z = e^x \sin y$, $y = e^x \cos x$, prove that $\frac{du}{dz} = e^x \left(\frac{\partial u}{\partial x} + \frac{\partial u}{\partial y} \right)$.
10. If $u = f(x - y, y - z, z - x)$, show that $\frac{\partial u}{\partial x} + \frac{\partial u}{\partial y} + \frac{\partial u}{\partial z} = 0$.
11. If $u = x^2 \tan^{-1} \left(\frac{y}{x} \right) - y^2 \tan^{-1} \left(\frac{x}{y} \right)$, determine $\frac{\partial^2 u}{\partial x^2}$.

2 | Page

RV COLLEGE OF ENGINEERING
(An Autonomous Institute Affiliated to VTU, Bangalore)
DEPARTMENT OF MATHEMATICS
FUNDAMENTALS OF LINEAR ALGEBRA, CALCULUS AND NUMERICAL METHODS (MATH1AT)
Multivariable Functions and Partial Differentiation

TUTORIAL SHEET 3

1. Match the following:
- | | |
|--|---------------------|
| (i) If $x = e^{\cos y}$, $y = e^{\cos x}$ then $\frac{dy}{dx} = \frac{1}{\sin(2x)}$ | (ii) minimum |
| (iii) If (u, v) is a stationary point of $f(x, y)$ and $f_{xx} > 0$, $f_{yy} > 0$ and $f_{xy} = 0$ at this point then the nature of (u, v) is | (iv) maximum |
| (iv) The nature of the point $(1, -1)$ to the function $f(x, y) = x^2 + 3xy - 3y^2 - 7y + 2$ is | (v) saddle point |
| (v) If $\frac{\partial^2 f}{\partial x^2} = \sin 2x$ and $\frac{\partial^2 f}{\partial y^2} = \frac{1}{2}$ then $\frac{\partial^2 f}{\partial x^2} = \frac{\partial^2 f}{\partial y^2}$ | (vi) maximum |
| (vi) If $\frac{\partial^2 f}{\partial x^2} = \sin 2x$ and $\frac{\partial^2 f}{\partial y^2} = \frac{1}{2}$ then $\frac{\partial^2 f}{\partial x^2} = \frac{\partial^2 f}{\partial y^2}$ | (vii) minimum |
| (vii) If $\frac{\partial^2 f}{\partial x^2} = \sin 2x$ and $\frac{\partial^2 f}{\partial y^2} = \frac{1}{2}$ then $\frac{\partial^2 f}{\partial x^2} = \frac{\partial^2 f}{\partial y^2}$ | (viii) saddle point |
| (viii) If $\frac{\partial^2 f}{\partial x^2} = \sin 2x$ and $\frac{\partial^2 f}{\partial y^2} = \frac{1}{2}$ then $\frac{\partial^2 f}{\partial x^2} = \frac{\partial^2 f}{\partial y^2}$ | (ix) maximum |
| (ix) If $\frac{\partial^2 f}{\partial x^2} = \sin 2x$ and $\frac{\partial^2 f}{\partial y^2} = \frac{1}{2}$ then $\frac{\partial^2 f}{\partial x^2} = \frac{\partial^2 f}{\partial y^2}$ | (x) minimum |
- Ans: (i) - (ii) (iii) - (iv) (v) - (vi) (vii) - (viii) (ix) - (ix) (x) - (x)

2. Find the extreme values of $\sin x + \sin y + \sin(x + y)$. Ans: Maximum value at $\left(\frac{\pi}{2}, \frac{\pi}{2}\right)$, minimum value is $\frac{3}{2}$.
3. Find the volume of largest rectangular parallelepiped that can be inscribed in an ellipsoid $\frac{x^2}{a^2} + \frac{y^2}{b^2} + \frac{z^2}{c^2} = 1$. Ans: Maximum volume = $\frac{8abc}{3}$.
4. Find the maximum and minimum distances of the point $(1, 2, 3)$ from the sphere $x^2 + y^2 + z^2 = 16$ using Lagrange's Method of undetermined multipliers. Ans: Maximum distance at $(2, 4, 6)$, $\sqrt{55}$; Minimum distance at $(-2, -4, -6)$, minimum distance = $\sqrt{55}$.
5. Show that $u = \frac{1}{\sqrt{x^2+y^2+z^2}}$ and $v = \frac{1}{\sqrt{x^2+y^2+z^2}}$ are functionally dependent and find the relation between them. Ans: $u^2 + v^2 = 1$.
6. For $u = xyz$, $v = yz + xz + xy$, $w = yz + xz$, find $\frac{\partial u}{\partial x}$. Ans: $(y - z)(x - y)$.
7. If $u = e^x \sec y$, $y = e^x \tan x$, then verify that $f = 1$.

2 | Page

$$\textcircled{1} y = 8 \sin \theta ; \left(\frac{dy}{d\theta} \right)^2 + \left(\frac{dx}{d\theta} \right)^2 = 9^2 \cos^2 \theta + x^2 \sin^2 \theta$$

$$x = 8 \cos \theta = 9^2$$

$$\frac{dx}{d\theta} = -8 \sin \theta$$

$$\textcircled{2} z = x \sin y + y \sin x$$

$$\frac{\partial}{\partial x} \left(\frac{dz}{dy} \right) = \frac{\partial}{\partial x} (x \cos y + \sin x)$$

$$= \cos y + \cos x$$

$$\textcircled{3} u = y \cos(xy)$$

$$\frac{\partial u}{\partial y} = (1) \cos(xy) + y(\sin(xy)) \cdot x$$

$$\frac{\partial^2 u}{\partial x \partial y} = -y \sin(xy) + \cos(xy)$$

$$\textcircled{4} u = x + y$$

$$\frac{\partial^2 u}{\partial x \partial y} = \frac{\partial}{\partial x} \left(\frac{\partial u}{\partial y} \right) = \frac{\partial}{\partial x} (1) = 0$$

$$\textcircled{5} v = f(x-ct) + g(x+ct)$$

$$\rightarrow \frac{dv}{dt} = f'(x-ct) \cdot (-c) + g'(x+ct) \cdot c$$

$$\frac{d^2 v}{dt^2} = c^2 f''(x-ct) + c^2 g''(x+ct)$$

$$\rightarrow \frac{dv}{dx} = f'(x-ct) + g'(x+ct)$$

$$\frac{d^2 v}{dx^2} = f''(x-ct) + g''(x+ct)$$

$$\rightarrow \frac{d^2 v}{dt^2} = c^2 \frac{d^2 v}{dx^2}$$

$$\textcircled{6} u = ae^{-gx} \sin(nt-gx)$$

$$\rightarrow \frac{\partial u}{\partial t} = ae^{-gx} \cos(nt-gx) \cdot n$$

$$\rightarrow \frac{\partial^2 u}{\partial x^2} = \frac{\partial}{\partial x} \left(\frac{\partial u}{\partial x} \right)$$

$$= \frac{\partial}{\partial x} \left(a(e^{-gx} \cos(nt-gx) - g \sin(nt-gx)) \right)$$

$$= \frac{\partial}{\partial x} (-age^{-gx} \cos(nt-gx) - g^2 \sin(nt-gx))$$

$$= -ag \left[e^{-gx} \cos(nt-gx) + g \sin(nt-gx) \right]$$

$$= -ag^2 e^{-gx} (\cos X - \sin X) = -ag^2 e^{-gx} \sin X$$

$$= -2ag^2 e^{-gx} \cos(nt-gx)$$

$$\frac{du}{dt} = M \frac{dz}{dt}$$

$$\frac{du}{dx} = M \frac{dz}{dx}$$

$$n = M \frac{dz}{dx}$$

$$g = \frac{n}{\partial M}$$

$$\textcircled{7} z = xy^2 + x^2y$$

$$\frac{dz}{dt} = \frac{\partial z}{\partial x} \frac{dx}{dt} + \frac{\partial z}{\partial y} \frac{dy}{dt}$$

$$= (y^2 + 2xy) \frac{dx}{dt} + (2xy + x^2) \frac{dy}{dt}$$

$$\text{at } t=1, x=1, y=2$$

$$\frac{dz}{dt} = (2^2 + 3(2)(1)) \cdot 3 + (2(1)(2) + 1^2) \cdot 4$$

$$= 30 + 20 = 50$$

$$\textcircled{8} (\cos x)^4 = (\sin y)^4 \rightarrow \text{implicit}$$

$$6 = (\cos x)^4 = (\sin y)^4$$

$$\frac{dy}{dx} = \frac{-\frac{1}{2} \cos x}{\frac{1}{2} \sin x} = -\frac{\cos x}{\sin x} = -\cot x$$

$$= -\frac{1}{\tan x} = -\frac{1}{\cot x} = -\tan x$$

$$\textcircled{9} u = x^2 - y^2 ; x = \frac{1}{t}, y = e^t$$

$$\frac{\partial u}{\partial x} = 2x = \frac{2}{t}$$

$$\frac{\partial u}{\partial y} = -2y = -2e^t$$

$$\textcircled{10} x^2 + y^2 + 3xz = 1$$

$$2x + 2y \frac{dy}{dx} + 3(x \frac{dz}{dx} + z) = 0$$

$$2x - 2y + 3x \frac{dz}{dx} + 3z = 0$$

$$3x \frac{dz}{dx} = 2y - 2x - 3z$$

$$\frac{dz}{dx} = \frac{2y}{3x} - \frac{2}{3} - \frac{3z}{3x}$$

$$= \frac{2y}{3x} - \frac{2}{3} - \frac{3z}{3x}$$

$$\textcircled{11} z = e^{2x+xy}$$

$$\frac{dz}{dy} = x e^{2x+xy}$$

$$\frac{\partial^2 z}{\partial x \partial y} = \frac{\partial}{\partial x} (x e^{2x+xy})$$

$$= e^{2x+xy} + x^2 e^{2x+xy}$$

$$= e^{2x+xy} (1 + x^2)$$

$$\textcircled{12} u = x + y$$

$$\frac{\partial^2 u}{\partial x \partial y} = \frac{\partial}{\partial x} \left(\frac{\partial u}{\partial y} \right) = \frac{\partial}{\partial x} (1) = 0$$

$$\textcircled{13} v = f(x-ct) + g(x+ct)$$

$$\frac{dv}{dt} = f'(x-ct) \cdot (-c) + g'(x+ct) \cdot c$$

$$\frac{d^2 v}{dt^2} = c^2 f''(x-ct) + c^2 g''(x+ct)$$

$$\rightarrow \frac{dv}{dx} = f'(x-ct) + g'(x+ct)$$

$$\frac{d^2 v}{dx^2} = f''(x-ct) + g''(x+ct)$$

$$\rightarrow \frac{d^2 v}{dt^2} = c^2 \frac{d^2 v}{dx^2}$$

$$\textcircled{14} V = xyz$$

$$\frac{\partial V}{\partial x} = yz = 4 \times 12 = 48$$

$$\textcircled{15} M = \log(x^3 + y^3 + z^3 - 3xyz)$$

$$\left(\frac{\partial}{\partial x} + \frac{\partial}{\partial y} + \frac{\partial}{\partial z} \right) M = \frac{-9}{(x+y+z)^2}$$

$$\rightarrow \left(\frac{\partial}{\partial x} + \frac{\partial}{\partial y} + \frac{\partial}{\partial z} \right) \left(\frac{3x^2 + 3y^2 + 3z^2 - 3xyz}{x^3 + y^3 + z^3 - 3xyz} \right)$$

$$\rightarrow \frac{\partial u}{\partial x} = \frac{1}{x^3 + y^3 + z^3 - 3xyz} \times (3x^2 - 3yz)$$

$$\rightarrow \frac{\partial u}{\partial y} = \frac{1}{x^3 + y^3 + z^3 - 3xyz} \times (3y^2 - 3xz)$$

$$\rightarrow \frac{\partial u}{\partial z} = \frac{1}{x^3 + y^3 + z^3 - 3xyz} \times (3z^2 - 3xy)$$

$$\left(\frac{\partial}{\partial x} + \frac{\partial}{\partial y} + \frac{\partial}{\partial z} \right) \left(\frac{3x^2 + 3y^2 + 3z^2 - 3xyz}{x^3 + y^3 + z^3 - 3xyz} \right)$$

$$= \left(\frac{\partial}{\partial x} + \frac{\partial}{\partial y} + \frac{\partial}{\partial z} \right) \left(\frac{3}{(x+y+z)^2} \right)$$

$$= \frac{\partial}{\partial x} \left(\frac{3}{(x+y+z)^2} \right) + \dots$$

$$= 3 \left(\frac{-1}{(x+y+z)^3} \right) + \dots$$

$$= -\frac{3}{(x+y+z)^2} + \dots$$

$$= -\frac{9}{(x+y+z)^2}$$

$$\textcircled{16} e^x - e^y = 2xy$$

$$f: e^x - e^y - 2xy = 0$$

$$\frac{dy}{dx} = -\frac{f_x}{f_y} = -\frac{e^x - 2y}{-e^y - 2x}$$

$$= \frac{e^x - 2y}{e^y + 2x}$$

$$\textcircled{17} M = xyz$$

$$x = e^t, y = e^t \sin t, z = \sin t$$

$$u = \frac{x}{y} = \frac{e^t}{e^t \sin t} = \frac{1}{\sin t}$$

$$\textcircled{18} z = \frac{x}{y}$$

$$\frac{dz}{dx} = \frac{1}{y}$$

$$\frac{dz}{dy} = -\frac{x}{y^2}$$

$$\frac{d^2 z}{dx dy} = \frac{\partial}{\partial x} \left(-\frac{x}{y^2} \right) = -\frac{1}{y^2}$$

$$\textcircled{19} x^2 + y^2 + 3xz = 1$$

$$2x + 2y \frac{dy}{dx} + 3(x \frac{dz}{dx} + z) = 0$$

$$2x - 2y + 3x \frac{dz}{dx} + 3z = 0$$

$$3x \frac{dz}{dx} = 2y - 2x - 3z$$

$$\frac{dz}{dx} = \frac{2y}{3x} - \frac{2}{3} - \frac{3z}{3x}$$

$$= \frac{2y}{3x} - \frac{2}{3} - \frac{3z}{3x}$$

$$\textcircled{20} z = \frac{x}{y}$$

$$\frac{dz}{dx} = \frac{1}{y}$$

$$\frac{dz}{dy} = -\frac{x}{y^2}$$

$$\frac{d^2 z}{dx dy} = \frac{\partial}{\partial x} \left(-\frac{x}{y^2} \right) = -\frac{1}{y^2}$$



$= \frac{\alpha - \alpha e}{t^2}$

6 $z \begin{cases} x \leq u \\ y \leq u \\ y \leq v \end{cases}$

$x = \rho^u \sin v \quad y = \rho^v \cos v$

$$u \begin{matrix} \nearrow x \\ \rightarrow y \\ \searrow z \end{matrix} \begin{matrix} \nwarrow \\ \rightarrow \\ \nearrow \end{matrix} t$$

$$\begin{aligned} \frac{dz}{du} &= \frac{\partial z}{\partial x} \frac{dx}{du} + \frac{\partial z}{\partial y} \frac{dy}{du} \\ &= \frac{\partial z}{\partial x} \cdot e^u \sin u + \frac{\partial z}{\partial y} (10) \\ &= \frac{\partial z}{\partial x} (\sin u e^u) \end{aligned}$$

$$\begin{aligned} \frac{du}{dt} &= \frac{\partial u}{\partial x} \frac{dx}{dt} + \frac{\partial u}{\partial y} \frac{dy}{dt} + \frac{\partial u}{\partial t} \frac{dt}{dt} \\ &= \frac{1}{2}(-e^{-t}) + \frac{1}{2}(-e^{-t} \sin^2 t + e^{-t} \sin 2t) \\ &\quad + xy \cos t \\ &= -e^{-t} \sin^2 t + e^{-t} \sin t (2 \cos t - \sin t) \\ &\quad + e^{-t} \sin^2 t \cos t \\ &= e^{-t} \sin^2 t (-\sin t + 2 \cos t - \sin t + \cos t) \\ \frac{du}{dt} &= e^{-t} \sin^2 t (3 \cos t - 2 \sin t) \end{aligned}$$

$$\frac{24}{3x} - \frac{1}{3} - \frac{(2x-x^2)}{3x^2} - \frac{1}{3}$$

8 $z = x^2 + 2xy + 4y^2, y = e^{3x}$

$$\begin{aligned} \frac{dz}{dx} &= 2x + 2\left(x \frac{dy}{dx} + y\right) + 4(2y) \frac{dy}{dx} \\ &= 2x + 2\left(x(2e^{3x}) + e^{3x}\right) \\ &\quad + 8(e^{3x})(2e^{3x}) \\ &= 2x + 2e^{3x}(3x+1) + 24e^{6x} \\ &= \underline{2x} + \underline{6xe^{3x}} + \underline{2e^{3x}} + \underline{24e^{6x}} \\ &= 2(x + e^{3x}) + 6e^{3x}(x + 4e^{3x}) \end{aligned}$$

$$\frac{dy}{dx} = 3e^{3x}$$

9) $z = \begin{matrix} x \\ y \end{matrix} = \begin{matrix} e^{i\theta} \cos \theta \\ e^{i\theta} \sin \theta \end{matrix}$

$PT_2: \quad y = e^{\cos x}$
 (i) $\frac{\partial z}{\partial u} = x \frac{\partial z}{\partial x} + y \frac{\partial z}{\partial y}$
 $\rightarrow \frac{\partial z}{\partial u} = \frac{\partial z}{\partial x} \cdot \frac{\partial x}{\partial u} + \frac{\partial z}{\partial y} \cdot \frac{\partial y}{\partial u}$
 $= \frac{\partial z}{\partial x} \cdot e^u \sin u + \frac{\partial z}{\partial y} \cdot e^u \cos u$
 $\frac{\partial z}{\partial u} = \frac{\partial z}{\partial x} \cdot x + \frac{\partial z}{\partial y} \cdot y$ ✓ proved

$$\begin{aligned} \frac{\partial z}{\partial u} &= \frac{\partial z}{\partial x} \frac{\partial x}{\partial u} + \frac{\partial z}{\partial y} \frac{\partial y}{\partial u} \\ \frac{\partial z}{\partial v} &= \frac{\partial z}{\partial x} \frac{\partial x}{\partial v} + \frac{\partial z}{\partial y} \frac{\partial y}{\partial v} \\ \frac{\partial z}{\partial u} &= \frac{\partial z}{\partial x} e^u \cos v + \frac{\partial z}{\partial y} e^u \sin v \\ \frac{\partial z}{\partial v} &= -\frac{\partial z}{\partial x} e^u \sin v + \frac{\partial z}{\partial y} e^u \cos v \end{aligned}$$

10 $u = f(x-y, y-z, z-x)$

$$q = x - y, \quad s = y - z, \quad t = z - x$$

u — 9 — x
— 5 — y — y
— 1 — 2 — 2
— 1 — x
— 2

$$\rightarrow \frac{\partial u}{\partial x} = \frac{\partial u}{\partial y} \frac{\partial y}{\partial x} + \frac{\partial u}{\partial s} \frac{\partial s}{\partial x} + \frac{\partial u}{\partial t} \frac{\partial t}{\partial x}$$

$$= \frac{\partial u}{\partial y} (1) + \frac{\partial u}{\partial s} (0) + \frac{\partial u}{\partial t} (-1)$$

$$\frac{\partial u}{\partial x} = \frac{\partial u}{\partial \lambda} - \frac{\partial u}{\partial t}$$

$$\begin{aligned} \rightarrow \frac{\partial u}{\partial y} &= \frac{\partial u}{\partial x} \frac{\partial x}{\partial y} + \frac{\partial u}{\partial s} \frac{\partial s}{\partial y} + \frac{\partial u}{\partial t} \frac{\partial t}{\partial y} \\ &= \frac{\partial u}{\partial x} (1) + \frac{\partial u}{\partial s} (1) + \frac{\partial u}{\partial t} (0) \end{aligned}$$

$$\frac{\partial u}{\partial y} = \frac{\partial u}{\partial s} - \frac{\partial u}{\partial r}$$

$$\rightarrow \frac{\partial u}{\partial z} = \frac{\partial u}{\partial r} \cdot \frac{\partial r}{\partial z} + \frac{\partial u}{\partial s} \cdot \frac{\partial s}{\partial z} + \frac{\partial u}{\partial t} \cdot \frac{\partial t}{\partial z}$$

$$= \frac{\partial u}{\partial r}(0) + \frac{\partial u}{\partial s}(-1) + \frac{\partial u}{\partial t}(1)$$

$$\frac{\partial u}{\partial z} = \frac{\partial u}{\partial t} - \frac{\partial u}{\partial s}$$

$$\begin{aligned} \rightarrow \frac{2x}{2x} + \frac{2x}{2x} + \frac{2x}{2x} &= \frac{2x}{2x} - \frac{2x}{2x} + \frac{2x}{2x} - \frac{2x}{2x} \\ &= 0 \end{aligned}$$

11) $u = x^2 \tan^{-1}\left(\frac{y}{x}\right) - y^2 \tan^{-1}\left(\frac{x}{y}\right)$

$$\begin{aligned} \frac{\partial^2 u}{\partial x \partial y} &= \frac{\partial}{\partial x} \left(\frac{\partial u}{\partial y} \right) \\ &= \frac{\partial}{\partial x} \left(x^2 \cdot \frac{1}{1+y^2} \cdot \left(\frac{1}{x} \right) - 2y \cdot \frac{1}{1+y^2} \cdot \frac{-x}{y^2} \right) \\ &= \frac{\partial}{\partial x} \left(\frac{x^3}{x^4+y^2} + \frac{2xy}{x^4+y^2} \right) \\ &= \frac{\partial}{\partial x} \left(\frac{x^3+2xy}{x^4+y^2} \right) \\ &= \frac{(3x^2+2y)(x^4+y^2) - (x^3+2xy)(2x+y^2)}{(x^4+y^2)^2} \\ &= \frac{3x^4+3x^2y^2+2x^2y+2y^3 - 2x^4-x^3y^2-4x^2y-2xy^3}{(x^4+y^2)^2} \\ &= \frac{x^4+3x^2y^2-2x^3y+2y^3-x^3y^2-2xy^3}{(x^4+y^2)^2} \end{aligned}$$

$$\frac{\partial z}{\partial y} = e^u \cdot \frac{\partial u}{\partial y}$$