

Title: Binary classification

URL: https://en.wikipedia.org/wiki/Binary_classification

PageID: 205393

Categories: Category:Machine learning, Category:Statistical classification

Source: Wikipedia (CC BY-SA 4.0). Content may require attribution.

Binary classification is the task of classifying the elements of a set into one of two groups (each called class). Typical binary classification problems include:

Medical testing to determine if a patient has a certain disease or not;

Quality control in industry, deciding whether a specification has been met;

In information retrieval, deciding whether a page should be in the result set of a search or not

In administration, deciding whether someone should be issued with a driving licence or not

In cognition, deciding whether an object is food or not food.

When measuring the accuracy of a binary classifier, the simplest way is to count the errors. But in the real world often one of the two classes is more important, so that the number of both of the different types of errors is of interest. For example, in medical testing, detecting a disease when it is not present (a false positive) is considered differently from not detecting a disease when it is present (a false negative).

Four outcomes

Given a classification of a specific data set, there are four basic combinations of actual data category and assigned category: true positives TP (correct positive assignments), true negatives TN (correct negative assignments), false positives FP (incorrect positive assignments), and false negatives FN (incorrect negative assignments).

These can be arranged into a 2x2 contingency table, with rows corresponding to actual value – condition positive or condition negative – and columns corresponding to classification value – test outcome positive or test outcome negative.

Evaluation

From tallies of the four basic outcomes, there are many approaches that can be used to measure the accuracy of a classifier or predictor. Different fields have different preferences.

The eight basic ratios

A common approach to evaluation is to begin by computing two ratios of a standard pattern. There are eight basic ratios of this form that one can compute from the contingency table, which come in four complementary pairs (each pair summing to 1). These are obtained by dividing each of the four numbers by the sum of its row or column, yielding eight numbers, which can be referred to generically in the form "true positive row ratio" or "false negative column ratio".

There are thus two pairs of column ratios and two pairs of row ratios, and one can summarize these with four numbers by choosing one ratio from each pair – the other four numbers are the complements.

The row ratios are:

true positive rate (TPR) = $(TP/(TP+FN))$, aka sensitivity or recall. These are the proportion of the population with the condition for which the test is correct. with complement the false negative rate (FNR) = $(FN/(TP+FN))$

with complement the false negative rate (FNR) = $(FN/(TP+FN))$

true negative rate (TNR) = $(TN/(TN+FP))$, aka specificity (SPC), with complement false positive rate (FPR) = $(FP/(TN+FP))$, also called independent of prevalence

with complement false positive rate (FPR) = $(FP/(TN+FP))$, also called independent of prevalence

The column ratios are:

positive predictive value (PPV, aka precision) $(TP/(TP+FP))$. These are the proportion of the population with a given test result for which the test is correct. with complement the false discovery rate (FDR) $(FP/(TP+FP))$

with complement the false discovery rate (FDR) $(FP/(TP+FP))$

negative predictive value (NPV) $(TN/(TN+FN))$ with complement the false omission rate (FOR) $(FN/(TN+FN))$, also called dependence on prevalence.

with complement the false omission rate (FOR) $(FN/(TN+FN))$, also called dependence on prevalence.

In diagnostic testing, the main ratios used are the true column ratios – true positive rate and true negative rate – where they are known as sensitivity and specificity . In informational retrieval, the main ratios are the true positive ratios (row and column) – positive predictive value and true positive rate – where they are known as precision and recall .

Cullerne Bown has suggested a flow chart for determining which pair of indicators should be used when. Otherwise, there is no general rule for deciding. There is also no general agreement on how the pair of indicators should be used to decide on concrete questions, such as when to prefer one classifier over another.

One can take ratios of a complementary pair of ratios, yielding four likelihood ratios (two column ratio of ratios, two row ratio of ratios). This is primarily done for the column (condition) ratios, yielding likelihood ratios in diagnostic testing . Taking the ratio of one of these groups of ratios yields a final ratio, the diagnostic odds ratio (DOR). This can also be defined directly as $(TP \times TN)/(FP \times FN) = (TP/FN)/(FP/TN)$; this has a useful interpretation – as an odds ratio – and is prevalence-independent.

Other metrics

There are a number of other metrics, most simply the accuracy or Fraction Correct (FC), which measures the fraction of all instances that are correctly categorized; the complement is the Fraction Incorrect (FiC). The F-score combines precision and recall into one number via a choice of weighing, most simply equal weighing, as the balanced F-score (F1 score). Some metrics come from regression coefficients : the markedness and the informedness , and their geometric mean , the Matthews correlation coefficient . Other metrics include Youden's J statistic , the uncertainty coefficient , the phi coefficient , and Cohen's kappa .

Statistical binary classification

Statistical classification is a problem studied in machine learning in which the classification is performed on the basis of a classification rule . It is a type of supervised learning , a method of machine learning where the categories are predefined, and is used to categorize new probabilistic observations into said categories. When there are only two categories the problem is known as statistical binary classification.

Some of the methods commonly used for binary classification are:

Decision trees

Random forests

Bayesian networks

Support vector machines

Neural networks

Logistic regression

Probit model

Genetic Programming

Multi expression programming

Linear genetic programming

Each classifier is best in only a select domain based upon the number of observations, the dimensionality of the feature vector , the noise in the data and many other factors. For example, random forests perform better than SVM classifiers for 3D point clouds.

Converting continuous values to binary

Binary classification may be a form of dichotomization in which a continuous function is transformed into a binary variable. Tests whose results are of continuous values, such as most blood values , can artificially be made binary by defining a cutoff value , with test results being designated as positive or negative depending on whether the resultant value is higher or lower than the cutoff.

However, such conversion causes a loss of information, as the resultant binary classification does not tell how much above or below the cutoff a value is. As a result, when converting a continuous value that is close to the cutoff to a binary one, the resultant positive or negative predictive value is generally higher than the predictive value given directly from the continuous value. In such cases, the designation of the test of being either positive or negative gives the appearance of an inappropriately high certainty, while the value is in fact in an interval of uncertainty. For example, with the urine concentration of hCG as a continuous value, a urine pregnancy test that measured 52 mIU/ml of hCG may show as "positive" with 50 mIU/ml as cutoff, but is in fact in an interval of uncertainty, which may be apparent only by knowing the original continuous value. On the other hand, a test result very far from the cutoff generally has a resultant positive or negative predictive value that is lower than the predictive value given from the continuous value. For example, a urine hCG value of 200,000 mIU/ml confers a very high probability of pregnancy, but conversion to binary values results in that it shows just as "positive" as the one of 52 mIU/ml.

See also

Mathematics portal

Approximate membership query filter

Examples of Bayesian inference

Classification rule

Confusion matrix

Detection theory

Kernel methods

Multiclass classification

Multi-label classification

One-class classification

Prosecutor's fallacy

Receiver operating characteristic

Thresholding (image processing)

Uncertainty coefficient , aka proficiency

Qualitative property

Precision and recall (equivalent classification schema)

References

Bibliography

Nello Cristianini and John Shawe-Taylor . An Introduction to Support Vector Machines and other kernel-based learning methods . Cambridge University Press, 2000. ISBN 0-521-78019-5 ([1] SVM Book)

John Shawe-Taylor and Nello Cristianini. Kernel Methods for Pattern Analysis . Cambridge University Press, 2004. ISBN 0-521-81397-2 (Website for the book)

Bernhard Schölkopf and A. J. Smola: Learning with Kernels . MIT Press, Cambridge, Massachusetts, 2002. ISBN 0-262-19475-9

v

t

e

Outline

Index

Mean Arithmetic Arithmetic-Geometric Contraharmonic Cubic Generalized/power Geometric Harmonic Heronian Heinz Lehmer

Arithmetic

Arithmetic-Geometric

Contraharmonic

Cubic

Generalized/power

Geometric

Harmonic

Heronian

Heinz

Lehmer

Median

Mode

Average absolute deviation

Coefficient of variation

Interquartile range

Percentile

Range

Standard deviation

Variance

Central limit theorem

Moments Kurtosis L-moments Skewness

Kurtosis

L-moments

Skewness

Index of dispersion

Contingency table
Frequency distribution
Grouped data
Partial correlation
Pearson product-moment correlation
Rank correlation Kendall's τ Spearman's ρ
Kendall's τ
Spearman's ρ
Scatter plot
Bar chart
Biplot
Box plot
Control chart
Correlogram
Fan chart
Forest plot
Histogram
Pie chart
Q–Q plot
Radar chart
Run chart
Scatter plot
Stem-and-leaf display
Violin plot
Effect size
Missing data
Optimal design
Population
Replication
Sample size determination
Statistic
Statistical power
Sampling Cluster Stratified
Cluster
Stratified
Opinion poll
Questionnaire
Standard error

Blocking
Factorial experiment
Interaction
Random assignment
Randomized controlled trial
Randomized experiment
Scientific control
Adaptive clinical trial
Stochastic approximation
Up-and-down designs
Cohort study
Cross-sectional study
Natural experiment
Quasi-experiment
Population
Statistic
Probability distribution
Sampling distribution Order statistic
Order statistic
Empirical distribution Density estimation
Density estimation
Statistical model Model specification L_p space
Model specification
 L_p space
Parameter location scale shape
location
scale
shape
Parametric family Likelihood (monotone) Location–scale family Exponential family
Likelihood (monotone)
Location–scale family
Exponential family
Completeness
Sufficiency
Statistical functional Bootstrap U V
Bootstrap
 U
 V

Optimal decision loss function

loss function

Efficiency

Statistical distance divergence

divergence

Asymptotics

Robustness

Estimating equations Maximum likelihood Method of moments M-estimator Minimum distance

Maximum likelihood

Method of moments

M-estimator

Minimum distance

Unbiased estimators Mean-unbiased minimum-variance Rao–Blackwellization Lehmann–Scheffé theorem Median unbiased

Mean-unbiased minimum-variance Rao–Blackwellization Lehmann–Scheffé theorem

Rao–Blackwellization

Lehmann–Scheffé theorem

Median unbiased

Plug-in

Confidence interval

Pivot

Likelihood interval

Prediction interval

Tolerance interval

Resampling Bootstrap Jackknife

Bootstrap

Jackknife

1- & 2-tails

Power Uniformly most powerful test

Uniformly most powerful test

Permutation test Randomization test

Randomization test

Multiple comparisons

Likelihood-ratio

Score/Lagrange multiplier

Wald

Z -test (normal)

Student's t -test

F -test
Chi-squared
G -test
Kolmogorov–Smirnov
Anderson–Darling
Lilliefors
Jarque–Bera
Normality (Shapiro–Wilk)
Likelihood-ratio test
Model selection Cross validation AIC BIC
Cross validation
AIC
BIC
Sign Sample median
Sample median
Signed rank (Wilcoxon) Hodges–Lehmann estimator
Hodges–Lehmann estimator
Rank sum (Mann–Whitney)
Nonparametric anova 1-way (Kruskal–Wallis) 2-way (Friedman) Ordered alternative (Jonckheere–Terpstra)
1-way (Kruskal–Wallis)
2-way (Friedman)
Ordered alternative (Jonckheere–Terpstra)
Van der Waerden test
Bayesian probability prior posterior
prior
posterior
Credible interval
Bayes factor
Bayesian estimator Maximum posterior estimator
Maximum posterior estimator
Correlation
Regression analysis
Pearson product-moment
Partial correlation
Confounding variable
Coefficient of determination
Errors and residuals

Regression validation
Mixed effects models
Simultaneous equations models
Multivariate adaptive regression splines (MARS)
Simple linear regression
Ordinary least squares
General linear model
Bayesian regression
Nonlinear regression
Nonparametric
Semiparametric
Isotonic
Robust
Homoscedasticity and Heteroscedasticity
Exponential families
Logistic (Bernoulli) / Binomial / Poisson regressions
Analysis of variance (ANOVA, anova)
Analysis of covariance
Multivariate ANOVA
Degrees of freedom
Cohen's kappa
Contingency table
Graphical model
Log-linear model
McNemar's test
Cochran–Mantel–Haenszel statistics
Regression
Manova
Principal components
Canonical correlation
Discriminant analysis
Cluster analysis
Classification
Structural equation model Factor analysis
Factor analysis
Multivariate distributions Elliptical distributions Normal
Elliptical distributions Normal
Normal

Decomposition
Trend
Stationarity
Seasonal adjustment
Exponential smoothing
Cointegration
Structural break
Granger causality
Dickey–Fuller
Johansen
Q-statistic (Ljung–Box)
Durbin–Watson
Breusch–Godfrey
Autocorrelation (ACF) partial (PACF)
partial (PACF)
Cross-correlation (XCF)
ARMA model
ARIMA model (Box–Jenkins)
Autoregressive conditional heteroskedasticity (ARCH)
Vector autoregression (VAR) (Autoregressive model (AR))
Spectral density estimation
Fourier analysis
Least-squares spectral analysis
Wavelet
Whittle likelihood
Kaplan–Meier estimator (product limit)
Proportional hazards models
Accelerated failure time (AFT) model
First hitting time
Nelson–Aalen estimator
Log-rank test
Bioinformatics
Clinical trials / studies
Epidemiology
Medical statistics
Chemometrics
Methods engineering
Probabilistic design

Process / quality control
Reliability
System identification
Actuarial science
Census
Crime statistics
Demography
Econometrics
Jurimetrics
National accounts
Official statistics
Population statistics
Psychometrics
Cartography
Environmental statistics
Geographic information system
Geostatistics
Kriging
Category
Mathematics portal
Commons
WikiProject