

Dysregulation of DNA Methylation in Human Intestinal Epithelial Organoids with Increasing Passage

Rachel Edgar

May 18, 2020

Diagnosis	Passage										Total
	1	2	3	4	6	7	8	11	14	16	
Control	5	27	8	5	5	5	1	0	0	0	56
Other GI	2	1	0	1	0	0	0	0	0	0	4
IBD	0	7	1	1	0	2	2	2	4	1	20
Total	7	35	9	7	5	7	3	2	4	1	80




Figure 1: Sampling site of origin and organoid passage number are associated to the main components of DNAm variation. (a) PC1 and PC2 are plotted for each sample. Samples are coloured by the sampling site of origin. (b) PC2 and PC3 are plotted for each sample. Samples are coloured by passage number. Lines connect samples derived from the same patient, but where organoids were cultured to a different number of passages. The histogram in the legend shows the distribution of passage numbers across the cohort.




Figure 2: The DNAm beta value distribution for high passage samples is trimodal while for low passage samples it is bimodal. All beta distributions displayed are for the 51,545 most variable CpGs. (a) Beta value distributions for all samples at a given passage number. Curves are coloured by the passage number. (b) The percent of samples considered trimodal. Points are coloured by passage and the bars around each point represent the confidence interval for that percent. (c) Beta value distributions for samples derived from the same patient but cultured to a different number of passages. Plots are labelled with the patient ID number, sampling site of origin and the passage number of each sample derived from that patient and sampling site. Curves are coloured by high or low passage relative to the other sample(s).

