

6.1 Rate of Reaction

- Reaction rates vary. Why? There are several factors. Section 6.2
- The study of reaction rates is called Reaction (or Chemical) Kinetics.
- The study of determining if a reaction will occur is called Thermodynamics

Describing Reaction Rates

- Reaction rates can be measured by changes of mass, gas production, concentration changes, conductivity or color change
- Most reaction rates are studied by changes in concentration
- The rate of a chemical reaction is defined as the change in concentration of a specific reactant or a specific product per unit time.
- Standard Unit = $\text{mol/L}\cdot\text{s}$ or $\text{mol}\cdot\text{L}^{-1}\cdot\text{s}^{-1}$

E.g. $\text{A} + \text{B} \rightarrow \text{AB}$


$$\text{rate} = \frac{\text{decrease in } [\text{A}]}{\text{time}}$$

square brackets mean “concentration of”

Note: Rate of reaction is always positive.

$$r = \frac{\text{change in concentration}}{\text{elapsed time}} = \frac{\Delta c}{\Delta t}$$

- A reaction rate with respect to a reactant will be negative
- A reaction rate with respect to a product will be positive
- A reaction rate is connected to the coefficients of a balanced chemical equation




If rate of the formation of A_2B_3 is $+ 3 \text{ mol/L}\cdot\text{s}$


Then the rate of decrease of A would be _____

And the rate of decrease of B would be _____

- Graphing examples of common rates – (looks like displacement / velocity / acceleration)



- Rates are usually determined at the beginning of a reaction due to the maximum amount of reactant present (max. collisions/sec.) Collision Theory is very important for understanding rates. (chapter 6.4)
- Most reaction rates change over time
- Rates are determined experimentally using many observations creating a graph and then calculating the slope at particular instances



Measuring Reaction Rates

- Reaction rates can be measured by directly measuring changes in concentration of the components or by measuring changes in concentration-related properties such as colour, density, electric conductivity, volume and pressure.
- Atomic absorption spectrometers, spectrophotometers, conductivity meters, and gas chromatographs are the favourite tools for measuring concentration.

Homework: Practice 1,2,3,4,5,6,7,8,11 and Questions 1,2

6.2 Factors Affecting Reaction Rates

- Nature of Reactants: Reactions between simple ions are almost instantaneous; reactions between more complex ions take longer.
- Concentration of the Reactants: The rate of reaction increases as the concentrations of the reactants increase.
- Temperature: Reactions occur faster at higher temperatures. In general a 10°C increase in temperature doubles the reaction rate.
- Catalysts: A catalyst is any reagent that increases the rate of reaction but is not consumed during the reaction.
- Surface Area: Increasing the surface area of the solid phase of a heterogeneous reaction increases the rate of the reaction.

Homework: Practice 1,2,3,4 Questions 1,2