

2.10 LIMITING AND EXCESS REAGENTS

PRACTICE

(Page 153)

Understanding Concepts

1. (a) $m_{\text{H}_2} = 10.0 \text{ g H}_2$
 $N_{\text{O}_2} = 32.00 \text{ g O}_2$

Balanced equation	$2 \text{ H}_{2(\text{g})}$	+	$\text{O}_{2(\text{g})}$	\rightarrow	$2 \text{ H}_2\text{O}_{(\text{l})}$
Before reaction	10.0 g		32.00 g		0 g
Reaction according to balanced equation	2 mol 4.04 g		1 mol 32.00 g		2 mol
After reaction	5.96 g		0 g		? g

According to the chart, oxygen is the limiting reagent, and hydrogen is the excess reagent.

(b)

$$n_{\text{O}_2} = 32.00 \text{ g O}_2 \times \frac{1 \text{ mol O}_2}{32.00 \text{ g O}_2}$$

$$n_{\text{O}_2} = 1.000 \text{ mol O}_2$$

$$n_{\text{H}_2\text{O}} = 1.000 \text{ mol O}_2 \times \frac{2 \text{ mol H}_2\text{O}}{1 \text{ mol O}_2}$$

$$n_{\text{H}_2\text{O}} = 2.000 \text{ mol H}_2\text{O}$$

$$m_{\text{H}_2\text{O}} = 2.000 \text{ mol H}_2\text{O} \times \frac{18.02 \text{ g H}_2\text{O}}{1 \text{ mol H}_2\text{O}}$$

$$m_{\text{H}_2\text{O}} = 36.04 \text{ g H}_2\text{O}$$

Therefore, 36.04 g of water is produced.

The combined calculation is as follows:

$$m_{\text{H}_2\text{O}} = 32.00 \text{ g O}_2 \times \frac{1 \text{ mol O}_2}{32.00 \text{ g O}_2} \times \frac{2 \text{ mol H}_2\text{O}}{1 \text{ mol O}_2} \times \frac{18.02 \text{ g H}_2\text{O}}{1 \text{ mol H}_2\text{O}}$$

$$m_{\text{H}_2\text{O}} = 36.04 \text{ g H}_2\text{O}$$

Therefore, 36.04 g of water is produced.

2. $m_{\text{CH}_4} = 32.1 \text{ g CH}_4$
 $m_{\text{O}_2} = 160.0 \text{ g O}_2$

Balanced equation	$2 \text{CH}_{4(g)} + 3 \text{O}_{2(g)} \rightarrow 2 \text{CO}_{(g)} + 4 \text{H}_2\text{O}_{(g)}$			
Before reaction	32.1 g	160.0 g	0 g	0 g
Reaction according to balanced equation	2 mol (32.10 g)	3 mol (96.00 g)		
After reaction	0 g	64.0 g	?	

According to the chart, methane is the limiting reagent and oxygen is the excess reagent.

$$n_{\text{CH}_4} = 32.1 \text{ g CH}_4 \times \frac{1 \text{ mol CH}_4}{16.05 \text{ g CH}_4}$$

$$n_{\text{CH}_4} = 2.00 \text{ mol CH}_4$$

$$n_{\text{CO}} = 2.00 \text{ mol CH}_4 \times \frac{2 \text{ mol CO}}{2 \text{ mol CH}_4}$$

$$n_{\text{CO}} = 2.00 \text{ mol CO}$$

$$m_{\text{CO}} = 2.00 \text{ mol CO} \times \frac{28.01 \text{ g CO}}{1 \text{ mol CO}}$$

$$m_{\text{CO}} = 56.0 \text{ g CO}$$

Therefore, 56.0 g of carbon monoxide is produced during incomplete combustion.

The combined calculation is as follows:

$$m_{\text{CO}} = 32.1 \text{ g CH}_4 \times \frac{1 \text{ mol CH}_4}{16.05 \text{ g CH}_4} \times \frac{2 \text{ mol CO}}{2 \text{ mol CH}_4} \times \frac{28.01 \text{ g CO}}{1 \text{ mol CO}}$$

$$m_{\text{CO}} = 56.0 \text{ g CO}$$

Therefore, 56.0 g of carbon monoxide is produced during incomplete combustion.

3. (a) $m_{\text{SO}_2} = 192.18 \text{ g SO}_2$

$$m_{\text{O}_2} = 32.00 \text{ g O}_2$$

Balanced equation	$2 \text{SO}_{2(g)} + \text{O}_{2(g)} \rightarrow 2 \text{SO}_{3(g)}$		
Before reaction	192.18 g	32.00 g	0 g
Reaction according to balanced equation	2 mol (128.00 g)	1 mol (32.00 g)	
After reaction	64.18 g	0 g	?

According to the chart, oxygen is the limiting reagent and sulfur dioxide is the excess reagent.

(b)

$$n_{\text{O}_2} = 32.00 \text{ g O}_2 \times \frac{1 \text{ mol O}_2}{32.00 \text{ g O}_2}$$

$$n_{\text{O}_2} = 1.000 \text{ mol O}_2$$

$$n_{\text{SO}_3} = 1.000 \text{ mol O}_2 \times \frac{2 \text{ mol SO}_3}{1 \text{ mol O}_2}$$

$$n_{\text{SO}_3} = 2.00 \text{ mol SO}_3$$

Therefore, 2.00 mol sulfur trioxide is produced.

$$(c) \quad m_{\text{SO}_3} = 2.00 \cancel{\text{ mol SO}_3} \times \frac{80.06 \text{ g SO}_3}{1 \cancel{\text{ mol SO}_3}}$$

$$m_{\text{SO}_3} = 160.1 \text{ g SO}_3$$

Therefore, 160.1 g of sulfur trioxide is produced.

4. $m_{\text{P}_4} = 123.88 \text{ g P}_4$

$$m_{\text{Cl}_2} = 1.00 \text{ kg Cl}_2 \times \frac{1000 \text{ g Cl}_2}{1 \text{ kg Cl}_2}$$

$$m_{\text{Cl}_2} = 1.00 \times 10^3 \text{ g Cl}_2$$

Balanced equation	$\text{P}_{4(s)}$	+	$10 \text{ Cl}_{2(g)}$	\rightarrow	$4 \text{ PCl}_{5(s)}$
Before reaction	123.88 g		354.5 g		0 g
Reaction according to balanced equation	1 mol (123.88 g)		10 mol (709.0 g)		
After reaction	0 g		354.5 g		?

$$n_{\text{P}_4} = 123.88 \cancel{\text{ g P}_4} \times \frac{1 \text{ mol P}_4}{123.88 \cancel{\text{ g P}_4}}$$

$$n_{\text{P}_4} = 1.000 \text{ mol P}_4$$

$$n_{\text{PCl}_5} = 1.000 \cancel{\text{ mol P}_4} \times \frac{4 \text{ mol PCl}_5}{1 \cancel{\text{ mol P}_4}}$$

$$n_{\text{PCl}_5} = 4.000 \text{ mol PCl}_5$$

$$m_{\text{PCl}_5} = 4.000 \cancel{\text{ mol PCl}_5} \times \frac{208.22 \text{ g PCl}_5}{1 \cancel{\text{ mol PCl}_5}}$$

$$m_{\text{PCl}_5} = 832.9 \text{ g PCl}_5$$

The mass of phosphorus pentachloride produced from the reaction is 832.9 g.

The combined calculation is as follows:

$$m_{\text{PCl}_5} = 123.88 \cancel{\text{ g P}_4} \times \frac{1 \cancel{\text{ mol P}_4}}{123.88 \cancel{\text{ g P}_4}} \times \frac{4 \cancel{\text{ mol PCl}_5}}{1 \cancel{\text{ mol P}_4}} \times \frac{208.22 \text{ g PCl}_5}{1 \cancel{\text{ mol PCl}_5}}$$

$$m_{\text{PCl}_5} = 832.9 \text{ g PCl}_5$$

The mass of phosphorus pentachloride produced from the reaction is 832.9 g.

SECTION 2.10 QUESTIONS

(Page 153)

Understanding Concepts

1. The excess reagent is the reactant that is not completely consumed in a chemical reaction. The limiting reagent is the reactant that is completely consumed in a chemical reaction. The limiting reagent limits the amount of product that can be formed.
2. No, there does not need to be a limiting reagent in a chemical reaction. If reactants are mixed in amounts equal to, or proportional to, the coefficients in the balanced reaction equation, then all reagents will be completely consumed.

- $$m_{\text{O}_2} = ?$$

$$m_{\text{CO}} = ?$$

From (a):

$$n_{\text{CH}_4} = 0.40 \text{ mol CH}_4$$

$$n_{\text{CO}} = 0.40 \text{ mol CH}_4 \times \frac{2 \text{ mol CO}}{2 \text{ mol CH}_4}$$

$$n_{\text{CO}} = 0.40 \text{ mol CO}$$

$$m_{\text{CO}} = 0.40 \text{ mol CO} \times \frac{28.01 \text{ g CO}}{1 \text{ mol CO}}$$

$$m_{\text{CO}} = 11 \text{ g CO}$$

The mass of carbon monoxide formed is 11 g.

The combined calculation is as follows:

$$m_{\text{CO}} = 6.4 \text{ g CH}_4 \times \frac{1 \text{ mol CH}_4}{16.05 \text{ g CH}_4} \times \frac{2 \text{ mol CO}}{2 \text{ mol CH}_4} \times \frac{28.01 \text{ g CO}}{1 \text{ mol CO}}$$

$$m_{\text{CO}} = 11 \text{ g CO}$$

The mass of carbon monoxide formed is 11 g.

5. $m_{\text{Al}} = 6.71 \text{ g Al}$

$$m_{\text{Al}_2(\text{SO}_4)_3} = ?$$

Balanced chemical equation	$2 \text{ Al}_{(s)} + 3 \text{ H}_2\text{SO}_{4(aq)} \rightarrow \text{Al}_2(\text{SO}_4)_{3(aq)} + 3 \text{ H}_{2(g)}$			
Given mass (g)	6.71 g		?	
Molar mass (g/mol)	26.98 g/mol		342.14 g/mol	

$$n_{\text{Al}} = 6.71 \text{ g Al} \times \frac{1 \text{ mol Al}}{26.98 \text{ g Al}}$$

$$n_{\text{Al}} = 0.2487 \text{ mol Al}$$

$$n_{\text{Al}_2(\text{SO}_4)_3} = 0.2487 \text{ mol Al} \times \frac{1 \text{ mol Al}_2(\text{SO}_4)_3}{2 \text{ mol Al}}$$

$$n_{\text{Al}_2(\text{SO}_4)_3} = 0.1244 \text{ mol Al}_2(\text{SO}_4)_3$$

$$m_{\text{Al}_2(\text{SO}_4)_3} = 0.1244 \text{ mol Al}_2(\text{SO}_4)_3 \times \frac{342.14 \text{ g Al}_2(\text{SO}_4)_3}{1 \text{ mol Al}_2(\text{SO}_4)_3}$$

$$m_{\text{Al}_2(\text{SO}_4)_3} = 42.5 \text{ g Al}_2(\text{SO}_4)_3$$

The mass of aluminum sulfate formed is 42.5 g.

The combined calculation is as follows:

$$m_{\text{Al}_2(\text{SO}_4)_3} = 6.71 \text{ g Al} \times \frac{1 \text{ mol Al}}{26.98 \text{ g Al}} \times \frac{1 \text{ mol Al}_2(\text{SO}_4)_3}{2 \text{ mol Al}} \times \frac{342.14 \text{ g Al}_2(\text{SO}_4)_3}{1 \text{ mol Al}_2(\text{SO}_4)_3}$$

$$m_{\text{Al}_2(\text{SO}_4)_3} = 42.5 \text{ g Al}_2(\text{SO}_4)_3$$

The mass of aluminum sulfate formed is 42.5 g.

6. $m_{\text{Cu}} = ?$

$$m_{\text{Cu}_2\text{O}} = 286.2 \text{ g Cu}_2\text{O}$$

$$m_{\text{Cu}_2\text{S}} = 286.2 \text{ g Cu}_2\text{S}$$

Balanced chemical equation	$2 \text{ Cu}_2\text{O}_{(\text{s})} + \text{Cu}_2\text{S}_{(\text{s})} \rightarrow 6 \text{ Cu}_{(\text{s})} + \text{SO}_{2(\text{g})}$			
Before reaction	286.2 g	286.2 g	0 g	0 g
Reaction according to balanced chemical equation	2 mol (286.20 g)	1 mol (159.16 g)	6 mol (381.30 g)	1 mol (64.06 g)
After reaction	0 g	127.0 g	?	

From the table, the limiting reagent is $\text{Cu}_2\text{O}_{(\text{s})}$ and the excess reagent is $\text{Cu}_2\text{S}_{(\text{s})}$.

$$n_{\text{Cu}_2\text{O}} = 286.2 \text{ g Cu}_2\text{O} \times \frac{1 \text{ mol Cu}_2\text{O}}{143.1 \text{ g Cu}_2\text{O}}$$

$$n_{\text{Cu}_2\text{O}} = 2.000 \text{ mol Cu}_2\text{O}$$

$$n_{\text{Cu}} = 2.000 \text{ mol Cu}_2\text{O} \times \frac{6 \text{ mol Cu}}{2 \text{ mol Cu}_2\text{O}}$$

$$n_{\text{Cu}} = 6.000 \text{ mol Cu}$$

$$m_{\text{Cu}} = 6.000 \text{ mol Cu} \times \frac{63.55 \text{ g Cu}}{1 \text{ mol Cu}}$$

$$m_{\text{Cu}} = 381.3 \text{ g Cu}$$

The mass of copper obtained from the reaction is 381.3 g.

The combined calculation is as follows:

$$m_{\text{Cu}} = 286.2 \text{ g Cu}_2\text{O} \times \frac{1 \text{ mol Cu}_2\text{O}}{143.1 \text{ g Cu}_2\text{O}} \times \frac{6 \text{ mol Cu}}{2 \text{ mol Cu}_2\text{O}} \times \frac{63.55 \text{ g Cu}}{1 \text{ mol Cu}}$$

$$m_{\text{Cu}} = 381.3 \text{ g Cu}$$

The mass of copper obtained from the reaction is 381.3 g.

2.11 INVESTIGATION: THE LIMITING REAGENT IN A CHEMICAL REACTION

(Page 154)

Prediction

- (a) According to the stoichiometric method, 2.32 g of strontium sulfate precipitate will be produced from the reaction of 2.00 g of strontium chloride with excess copper(II) sulfate. The reaction equation is shown below.