

Homework 2

Due: Monday, April 12, at 11:00 PM Seattle time on Gradescope

1 Inner Products

Identify which of the following are inner products. For the ones that are *not* inner products, show a property of inner products that they do not satisfy (i.e. **symmetric, linear in the first argument, or positive definite**). As an example, consider \langle, \rangle_0 defined on $C([0, 1])$ by

$$\langle f, g \rangle_0 = f(1/2)g(1/2)$$

Then \langle, \rangle_0 is symmetric, and linear in the first argument (can you show this?) but it is *not* positive definite. Let $f(x) = x - 1/2$. Then f is not identically zero, but $\langle f, f \rangle_0 = 0$.

A. Let $A = \begin{pmatrix} 1 & 2 \\ 3 & 4 \end{pmatrix}$. Define \langle, \rangle_1 on \mathbb{R}^2 by

$$\langle v, w \rangle_1 = (Av) \bullet (Aw)$$

B. Let $B = \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix}$. Define \langle, \rangle_2 on \mathbb{R}^2 by

$$\langle v, w \rangle_2 = (Bv) \bullet (Bw)$$

C. Define \langle, \rangle_3 on $C^1([0, 1])$ by

$$\langle f, g \rangle_3 = \int_0^1 f'(s)g'(s)ds$$

D. Define \langle, \rangle_4 on $C([0, 1])$ by

$$\langle f, g \rangle_4 = \int_0^1 (1 + s^2)f(s)g(s)ds$$

E. Define \langle, \rangle_5 on $C([0, 1])$ by

$$\langle f, g \rangle_5 = \int_0^1 f(s)g(s)ds + f(1/2)$$

2 Initial Value Problems

For each of the following initial value problems

- Solve the IVP
- Compute $\lim_{t \rightarrow \infty} |x(t)|$, for the solution $|x(t)|$.

You may use a computer to diagonalize matrices and solve for coefficients, once you are confident with your skills. In particular, the coefficients in (D.) and (F.) are tricky to solve for by hand.

A.

$$\begin{cases} x' &= \begin{pmatrix} 11 & -6 \\ 18 & -10 \end{pmatrix} \\ x(0) &= (4, 7) \end{cases}$$

B.

$$\begin{cases} x' &= \begin{pmatrix} 4 & 2 \\ -1 & 1 \end{pmatrix} \\ x(0) &= (3, -2) \end{cases}$$

C.

$$\begin{cases} x' &= \begin{pmatrix} 8 & 10 \\ -5 & -7 \end{pmatrix} \\ x(0) &= (1, -1) \end{cases}$$

D.

$$\begin{cases} x' &= \begin{pmatrix} 8 & 10 \\ -5 & -7 \end{pmatrix} \\ x(0) &= (1.01, -1) \end{cases}$$

E.

$$\begin{cases} x' &= \begin{pmatrix} -4 & -2 \\ 1 & -1 \end{pmatrix} \\ x(0) &= (2, -1) \end{cases}$$

F.

$$\begin{cases} x' &= \begin{pmatrix} -4 & -2 \\ 1 & -1 \end{pmatrix} \\ x(0) &= (2.01, -1) \end{cases}$$