

## Database Questions and Answers – Tuple Relational Calculus and Domain Relational Calculus

[« Prev](#)[Next »](#)

This set of Database test focuses on “Tuple Relational Calculus and Domain Relational Calculus”.

1. Find the ID, name, dept name, salary for instructors whose salary is greater than \$80,000 .

- a)  $\{t \mid t \in \text{instructor} \wedge t[\text{salary}] > 80000\}$
- b)  $\exists t \in r (Q(t))$
- c)  $\{t \mid \exists s \in \text{instructor} (t[\text{ID}] = s[\text{ID}] \wedge s[\text{salary}] > 80000)\}$
- d) None of the mentioned

[View Answer](#)

Answer: a

Explanation: This expression is in tuple relational format.

2. A query in the tuple relational calculus is expressed as:

- a)  $\{t \mid P() \mid t\}$
- b)  $\{P(t) \mid t\}$
- c)  $\{t \mid P(t)\}$
- d) All of the mentioned

[View Answer](#)

^

Answer: c

Explanation: The tuple relational calculus, is a nonprocedural query language. It describes the desired information without giving a specific procedure for obtaining that information.

advertisement

3.

```
{t | ∃ s ∈ instructor (t[name] = s[name]
∧ ∃ u ∈ department (u[dept name] = s[dept name]
∧ u[building] = "Watson"))}
```

Which of the following best describes the query?

- a) Finds the names of all instructors whose department is in the Watson building
- b) Finds the names of all department is in the Watson building
- c) Finds the name of the department whose instructor and building is Watson
- d) Returns the building name of all the departments

[View Answer](#)

Answer: a

Explanation: This query has two "there exists" clauses in our tuple-relational-calculus expression, connected by and ( $\wedge$ ).

advertisement



4. Which of the following symbol is used in the place of except?

- a)  $\wedge$
- b)  $\vee$
- c)  $\neg$
- d)  $\sim$

[View Answer](#)

Answer: c

Explanation: The query  $\neg P$  negates the value of P.

5. "Find all students who have taken all courses offered in the Biology department." The expressions that matches this sentence is :

- a)  $\exists t \in r(Q(t))$
- b)  $\forall t \in r(Q(t))$
- c)  $\neg t \in r(Q(t))$
- d)  $\sim t \in r(Q(t))$

[View Answer](#)

Answer: b

Explanation:  $\forall$  is used denote "for all" in SQL.

6. Which of the following is the comparison operator in tuple relational calculus

- a)  $\Rightarrow$
- b)  $=$
- c)  $\epsilon$
- d) All of the mentioned

[View Answer](#)

Answer: b

Explanation: The comparison operators are ( $<$ ,  $\leq$ ,  $=$ ,  $\neq$ ,  $>$ ,  $\geq$ ).

7. An expression in the domain relational calculus is of the form

- a)  $\{P(x_1, x_2, \dots, x_n) \mid \langle x_1, x_2, \dots, x_n \rangle\}$
- b)  $\{x_1, x_2, \dots, x_n \mid \langle x_1, x_2, \dots, x_n \rangle\}$
- c)  $\{x_1, x_2, \dots, x_n \mid x_1, x_2, \dots, x_n\}$
- d)  $\{\langle x_1, x_2, \dots, x_n \rangle \mid P(x_1, x_2, \dots, x_n)\}$

[View Answer](#)

Answer: d

Explanation: Here  $x_1, x_2, \dots, x_n$  represent domain variables.  $P$  represents a formula composed of atoms, as was the case in the tuple relational calculus.

advertisement

8. Find the names of all instructors in the Physics department together with the course id of all courses they teach:

a)

$$\{ \langle c \rangle \mid \exists s (\langle c, a, s, y, b, r, t \rangle \in \text{section} \wedge s = \text{"Fall"} \wedge y = \text{"2009"} \vee \exists u (\langle c, a, s, y, b, r, t \rangle \in \text{section} \wedge s = \text{"Spring"} \wedge y = \text{"2010"})) \}$$

b)

$$\{ \langle n, c \rangle \mid \exists i, a (\langle i, c, a, s, y \rangle \in \text{teaches} \wedge \exists d, s (\langle i, n, d, s \rangle \in \text{instructor} \wedge d = \text{"Physics"})) \}$$

c)

$$\{ \langle n \rangle \mid \exists i, d, s (\langle i, n, d, s \rangle \in \text{instructor} \wedge s > 80000) \}$$

d)

$$\{ \langle i, n, d, s \rangle \mid \langle i, n, d, s \rangle \in \text{instructor} \wedge s > 80000 \}$$

[View Answer](#)

Answer: b

Explanation: None.

9. In domain relational calculus "there exist" can be expressed as

- a)  $(P1(x))$
- b)  $(P1(x)) \exists x$
- c)  $\forall x (P1(x))$
- d)  $\exists x (P1(x))$

[View Answer](#)

Answer: d

Explanation:  $\exists$  is used to denote "some" values in relational calculus.

advertisement

10. A set of possible data values is called

- a) Attribute
- b) Degree
- c) Tuple
- d) Domain

[View Answer](#)

Answer: d

Explanation: None.



**Sanfoundry Global Education & Learning Series – Database Management System.**